
4NCG (X-ray, 2.58 Å) vs 7Z2G (cryo-EM, 3.65 Å, + DNA): DOR heavy-atom RMSD 0.62 Å in the NNIBP frame
a where the two structures differ
Cα RMSD, palm frame (Å)
1.83
1.21
1.62
0.77
1.37
2.23
0.97
fingers
palm
thumb
connection
RNase H
p51
NNIBP lining
b closest approach of each residue to DOR
4NCG (X-ray)
7Z2G (cryo-EM)
r = 0.974
median |Δ| = 0.16 Å
largest |Δ| = 0.41 Å
Lys103
Gly190
Val189
Tyr181
Tyr188
Lys101
Trp229
Val179
Val106
Leu234
Tyr318
Phe227
Leu100
Val108
Thr107
Glu138
minimum heavy-atom distance to DOR (Å)
c NNIBP side chains
median 0.80 Å
side-chain RMSD, NNIBP frame (Å)
Leu100
Lys101
Lys103
Val106
Thr107
Val108
Val179
Tyr181
Tyr188
Val189
Phe227
Trp229
Leu234
Tyr318
Glu138
d DOR dihedrals: WT simulated ensemble (n = 860 frames)
dihedral (deg)
4NCG
7Z2G
τ1
τ2
τ3
τ4