

A stochastic approximation method for chance-constrained nonlinear programs

Rohit Kannan

UW-Madison

December 10, 2018

Joint work with Jim Luedtke

Outline

1 Introduction

Formulation

Central hypothesis

Main idea

2 Related approaches

3 Proposed Approach

4 Computational results

5 Summary and Open questions

Formulation

$$\begin{aligned} \nu^* &:= \min_{x \in X} f(x) \\ \text{s.t. } &\mathbb{P}\{g(x, \xi) \leq 0\} \geq 1 - \alpha. \end{aligned} \tag{CCP}$$

- Assume:

- ▶ $X \subset \mathbb{R}^n$ is nonempty, compact, and convex
- ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and quasiconvex
- ▶ $g : \mathbb{R}^n \times \mathbb{R}^d \rightarrow \mathbb{R}^m$ is continuously differentiable
- ▶ $\xi \sim \mathcal{P}$ is continuous with support $\Xi \subset \mathbb{R}^d$
- ▶ Other relatively mild technical assumptions ...

Formulation

$$\begin{aligned} \nu^* &:= \min_{x \in X} f(x) && \text{(CCP)} \\ \text{s.t. } &\mathbb{P}\{g(x, \xi) \leq 0\} \geq 1 - \alpha. \end{aligned}$$

- Assume:
 - ▶ $X \subset \mathbb{R}^n$ is nonempty, compact, and convex
 - ▶ $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous and quasiconvex
 - ▶ $g : \mathbb{R}^n \times \mathbb{R}^d \rightarrow \mathbb{R}^m$ is continuously differentiable
 - ▶ $\xi \sim \mathcal{P}$ is continuous with support $\Xi \subset \mathbb{R}^d$
 - ▶ Other relatively mild technical assumptions ...
- Can model joint chance constraints, deterministic nonconvex constraints, and some recourse structures

Central hypothesis

Main idea #1

In many cases, decision makers are interested in generating the efficient frontier of optimal objective function value (ν^*) versus risk level (α) rather than simply solving (CCP) for a single prespecified risk level so that they can make a more informed decision.

— Rengarajan and Morton [2009], Luedtke [2014]

Central hypothesis

Main idea #1

In many cases, decision makers are interested in generating the efficient frontier of optimal objective function value (ν^*) versus risk level (α) rather than simply solving (CCP) for a single prespecified risk level so that they can make a more informed decision.

— Rengarajan and Morton [2009], Luedtke [2014]

- Efficient frontier of (CCP) can be recovered by solving the following stochastic optimization problem:

$$\begin{aligned} \min_{x \in X} \mathbb{P}\{g(x, \xi) \not\leq 0\} &\quad \equiv \quad \min_{x \in X} \mathbb{E}[\max[1, [g(x, \xi)]]] & \text{(SP)} \\ \text{s.t. } f(x) \leq \nu. & & \text{s.t. } f(x) \leq \nu. \end{aligned}$$

► $\mathbb{1}[\cdot]$ denotes the l.s.c. step function

Main idea #2

- Approximate the efficient frontier of (CCP) using:

$$\begin{array}{ll} \min_{x \in X} \mathbb{E} [\max [1 [g(x, \xi)]]] & \approx \min_{x \in X} \mathbb{E} [\max [\phi_k (g(x, \xi))]] \quad (\text{APP}_k) \\ \text{s.t. } f(x) \leq \nu. & \text{s.t. } f(x) \leq \nu. \end{array}$$

$\{\phi_k\}$ is a 'convergent' seq. of smooth approx. of the step function.

Main idea #2

- Approximate the efficient frontier of (CCP) using:

$$\begin{array}{ll} \min_{x \in X} \mathbb{E} [\max [1 [g(x, \xi)]]] & \approx \min_{x \in X} \mathbb{E} [\max [\phi_k (g(x, \xi))]] \quad (\text{APP}_k) \\ \text{s.t. } f(x) \leq \nu. & \text{s.t. } f(x) \leq \nu. \end{array}$$

$\{\phi_k\}$ is a 'convergent' seq. of smooth approx. of the step function.

- (APP_k) can be solved using stochastic gradient methods.

Main idea #2

- Approximate the efficient frontier of (CCP) using:


$$\min_{x \in X} \mathbb{E} [\max [1 [g(x, \xi)]]] \approx \min_{x \in X} \mathbb{E} [\max [\phi_k (g(x, \xi))]] \quad (\text{APP}_k)$$

$$\text{s.t. } f(x) \leq \nu.$$

$$\text{s.t. } f(x) \leq \nu.$$

$\{\phi_k\}$ is a 'convergent' seq. of smooth approx. of the step function.

- (APP_k) can be solved using stochastic gradient methods.



Outline

1 Introduction

2 Related approaches

- Scenario approximation

- Smoothing-based approaches

- Motivation for proposed approach

3 Proposed Approach

4 Computational results

5 Summary and Open questions

Scenario approximation

- Sample N random realizations of ξ from \mathcal{P}
- Solve the following scenario problem to local optimality using a cutting-plane approach:

$$\begin{aligned} \hat{x}_N \in \arg \min_{x \in X} f(x) \\ \text{s.t. } g(x, \xi_j) \leq 0, \quad \forall j \in \{1, \dots, N\}. \end{aligned}$$

- Estimate the risk level $\hat{\alpha}_N := \mathbb{P}\{g(\hat{x}_N, \xi) \not\leq 0\}$ using an independent Monte Carlo sample to determine the point $(\hat{\alpha}_N, f(\hat{x}_N))$ on the approximation of the efficient frontier
- References: Calafiore and Campi [2005], Campi and Garatti [2011]

Existing smoothing-based approaches

- Approximate the solution of (CCP) using:

$$\begin{array}{ll} \min_{x \in X} f(x) & \approx \min_{x \in X} f(x) \\ \text{s.t. } \mathbb{E} [\max [1 [g(x, \xi)]]] \leq \alpha. & \text{s.t. } \mathbb{E} [\max [\phi_k (g(x, \xi))]] \leq \alpha. \end{array}$$

where $\{\phi_k\}$ is a 'convergent' sequence of conservative smooth approximations of the step function.

- Solution of each approximation yields a feasible solution to the original chance-constrained program

Existing smoothing-based approaches

- Approximate the solution of (CCP) using:

$$\begin{array}{ll} \min_{x \in X} f(x) & \approx \min_{x \in X} f(x) \\ \text{s.t. } \mathbb{E} [\max [1 [g(x, \xi)]]] \leq \alpha. & \text{s.t. } \mathbb{E} [\max [\phi_k (g(x, \xi))]] \leq \alpha. \end{array}$$

where $\{\phi_k\}$ is a 'convergent' sequence of conservative smooth approximations of the step function.

- Solution of each approximation yields a feasible solution to the original chance-constrained program
- Currently, the only known approach for solving optimization problems with nonconvex expectation constraints is via sample average approximation
- References: Hong et al. [2011], Shan et al. [2014, 2016], Geletu et al. [2017], Cao and Zavala [2017], Adam et al. [2018]

Pros and cons of exterior sampling approaches

Pros

- ▶ Can use off-the-shelf solvers
- ▶ Possess strong theoretical guarantees

Cons

- ▶ May find spurious local optima
- ▶ May need to solve large-scale NLPs


- We propose the first interior sampling approach for solving chance-constrained problems

Sampling may lead to spurious local minima

- Consider the following chance constraint [Curtis et al., 2018]:

$$\mathbb{P} \left\{ 0.25x_1^4 - \frac{1}{3}x_1^3 - x_1^2 + 0.2x_1 - 19.5 + \xi_1x_1 + \xi_1\xi_0 \leq x_2 \right\} \geq 0.95,$$

where $\xi_1 \sim U(-3, 3)$ and $\xi_0 \sim U(-12, 12)$ are independent.



Outline

- 1 Introduction
- 2 Related approaches
- 3 Proposed Approach**
 - Main theoretical results
 - Approximating the efficient frontier
- 4 Computational results
- 5 Summary and Open questions

Main theoretical results

- Let $\{\phi_k\}$ be a ‘convergent’ sequence of smooth approximations of the step function.

$$\begin{array}{ll} \min_{x \in X} \mathbb{E} [\max [\mathbb{1} [g(x, \xi)]]] & \text{(SP)} \\ \text{s.t. } f(x) \leq \nu. & \end{array} \quad \begin{array}{ll} \min_{x \in X} \mathbb{E} [\max [\phi_k (g(x, \xi))]] & \text{(APP}_k\text{)} \\ \text{s.t. } f(x) \leq \nu. & \end{array}$$

Theorem

Every limit point of a sequence of global solutions to the approximations (APP_k) is a global solution to Problem (SP)

Tentative Theorem

Every limit point of a sequence of stationary solutions to the approximations (APP_k) is a stationary solution to Problem (SP)

Proposal for approximating the efficient frontier

- **Input:** initial guess $\hat{x}^0 \in X$, sequence of objective bounds $\{\bar{\nu}^k\}$ (determined by solving a scenario approximation problem), and lower bound on risk level $\alpha_{low} \in (0, 1)$
- **Output:** pairs $\{(\bar{\nu}^i, \bar{\alpha}^i)\}$ of objective values and risk levels used to approximate the efficient frontier (+ solutions $\{\bar{x}^i\}$)

Proposal for approximating the efficient frontier

- **Input:** initial guess $\hat{x}^0 \in X$, sequence of objective bounds $\{\bar{\nu}^k\}$ (determined by solving a scenario approximation problem), and lower bound on risk level $\alpha_{low} \in (0, 1)$
- **Output:** pairs $\{(\bar{\nu}^i, \bar{\alpha}^i)\}$ of objective values and risk levels used to approximate the efficient frontier (+ solutions $\{\bar{x}^i\}$)
- **Solve:** sequence of problems (APP_k) with objective bound $\bar{\nu}^k$ using the projected stochastic subgradient method [Davis and Drusvyatskiy, 2018]
- **Estimate:** risk level $\bar{\alpha}^k$ of the best found solution using an independent Monte Carlo sample to determine the point $(\bar{\nu}^k, \bar{\alpha}^k)$ on our approximation of the efficient frontier
- **Terminate:** when $\bar{\alpha}^k \leq \alpha_{low}$


Outline

- 1 Introduction
- 2 Related approaches
- 3 Proposed Approach
- 4 Computational results
 - Portfolio optimization
 - Resource planning
- 5 Summary and Open questions

Portfolio optimization [Ben-Tal et al., 2009]

$$\begin{aligned} \max_{t, x \in X} \quad & t \\ \text{s.t.} \quad & \mathbb{P} \left\{ \xi^T x \geq t \right\} \geq 1 - \alpha, \end{aligned}$$

where $X := \{x \in \mathbb{R}_+^{1000} : \sum_i x_i = 1\}$, ξ_i are independent Normal.



Resource planning [Luedtke, 2014]


$$\begin{aligned} \min_{x \in \mathbb{R}_+^{20}} \quad & c^\top x \\ \text{s.t.} \quad & \mathbb{P}\{x \in R(\lambda, \rho)\} \geq 1 - \alpha, \end{aligned}$$

where

$$R(\lambda, \rho) = \left\{ x \in \mathbb{R}_+^{20} : \exists y \in \mathbb{R}_+^{20 \times 30} \text{ s.t. } \sum_{j=1}^{30} y_{ij} \leq \rho_i x_i^2, \forall i \in \{1, \dots, 20\}, \right. \\ \left. \sum_{i=1}^{20} \mu_{ij} y_{ij} \geq \lambda_j, \forall j \in \{1, \dots, 30\} \right\}.$$

- ▶ x_i : quantity of resource i , c_i : unit cost of resource i
- ▶ y_{ij} : amount of resource i allocated to customer type j
- ▶ $\rho_i \in (0, 1]$: **random** yield of resource i
- ▶ $\lambda_j \geq 0$: **random** demand of customer type j
- ▶ $\mu_{ij} \geq 0$: service rate of resource i for customer type j

Resource planning



Outline

- 1 Introduction
- 2 Related approaches
- 3 Proposed Approach
- 4 Computational results
- 5 Summary and Open questions

Summary and Open questions

- Proposed a **stochastic approximation approach** for generating the efficient frontier of **chance-constrained NLPs**
- Computational results indicate that **our proposal consistently determines better approximations of the efficient frontier** than existing approaches in reasonable computation times
- Open questions
 - Extension to multiple sets of joint chance constraints
 - Handle deterministic nonconvex constraints more naturally
 - Theory for randomized constraint projection techniques to reduce effort spent on projections
 - Extension to distributionally robust chance constraints

References I

- Lukáš Adam, Martin Branda, Holger Heitsch, and René Henrion. Solving joint chance constrained problems using regularization and Benders decomposition. *Annals of Operations Research*, pages 1–27, 2018. doi: 10.1007/s10479-018-3091-9.
- Aharon Ben-Tal, Laurent El Ghaoui, and Arkadi Nemirovski. *Robust optimization*. Princeton University Press, 2009.
- Giuseppe Calafiore and Marco C Campi. Uncertain convex programs: randomized solutions and confidence levels. *Mathematical Programming*, 102(1):25–46, 2005.
- Marco C Campi and Simone Garatti. A sampling-and-discarding approach to chance-constrained optimization: feasibility and optimality. *Journal of Optimization Theory and Applications*, 148(2):257–280, 2011.
- Yankai Cao and Victor Zavala. A sigmoidal approximation for chance-constrained nonlinear programs.
http://www.optimization-online.org/DB_FILE/2017/10/6236.pdf, 2017.
- Frank E Curtis, Andreas Wachter, and Victor M Zavala. A sequential algorithm for solving nonlinear optimization problems with chance constraints. *SIAM Journal on Optimization*, 28(1):930–958, 2018.
- Damek Davis and Dmitriy Drusvyatskiy. Stochastic subgradient method converges at the rate $O(k^{-1/4})$ on weakly convex functions. *arXiv preprint arXiv:1802.02988*, 2018.

References II

- Abebe Geletu, Armin Hoffmann, Michael Kloppel, and Pu Li. An inner-outer approximation approach to chance constrained optimization. *SIAM Journal on Optimization*, 27(3):1834–1857, 2017.
- L Jeff Hong, Yi Yang, and Liwei Zhang. Sequential convex approximations to joint chance constrained programs: A Monte Carlo approach. *Operations Research*, 59(3):617–630, 2011.
- James Luedtke. A branch-and-cut decomposition algorithm for solving chance-constrained mathematical programs with finite support. *Mathematical Programming*, 146(1-2):219–244, 2014.
- Tara Rengarajan and David P Morton. Estimating the efficient frontier of a probabilistic bicriteria model. In *Winter Simulation Conference*, pages 494–504, 2009.
- F Shan, XT Xiao, and LW Zhang. Convergence analysis on a smoothing approach to joint chance constrained programs. *Optimization*, 65(12):2171–2193, 2016.
- Feng Shan, Liwei Zhang, and Xiantao Xiao. A smoothing function approach to joint chance-constrained programs. *Journal of Optimization Theory and Applications*, 163(1):181–199, 2014.