

- 1.2.1 (a) Prove that $\sqrt{3}$ is irrational. Does a similar argument work to show $\sqrt{6}$ is irrational?

Proof. Assume for the sake of contradiction that $\sqrt{3}$ is rational, that is there exist coprime integers p and q such that $3 = (\frac{p}{q})^2$. Then, $3q^2 = p^2$. Since p^2 is divisible by 3, p can be represented as $p = 3r$ for some integer r . So, $3q^2 = (3r)^2$ which simplifies to $q^2 = 3r^2$. Since, q^2 is divisible by 3, we have shown that both p and q are divisible by 3. However, this is a contradiction to the original claim that p and q are coprime integers. Thus, $\sqrt{3}$ is irrational. \square

Yes, a similar argument can be made to show that $\sqrt{6}$ is irrational.

- (b) Where does the proof of Theorem 1.1.1 break down if we try to use it to prove $\sqrt{4}$ is irrational?

It breaks down when we have that $4q^2 = p^2$ where p and q are coprime integers. It breaks down because $p = 2q$ and so p is not always divisible by 4 which means that we cannot represent p as $p = 4r$ for some $r \in \mathbb{Z}$. Thus, the proof fails.

- 1.2.2 Decide which of the following represent true statements about the nature of sets. For any that are false, provide a specific example where the statement in question does not hold.

- (a) If $A_1 \supseteq A_2 \supseteq A_3 \supseteq A_4 \dots$ are all sets containing an infinite number of elements, then the intersection $\bigcap_{n=1}^{\infty} A_n$ is infinite as well.

False. Let $A_n = \mathbb{N}_{\geq n}$ where $n \in \mathbb{N}$. Then, $\bigcap_{n=1}^{\infty} A_n = \emptyset$ which is a size of 0.

- (b) If $A_1 \supseteq A_2 \supseteq A_3 \supseteq A_4 \dots$ are all finite, nonempty sets of real numbers, then the intersection $\bigcap_{n=1}^{\infty} A_n$ is finite and nonempty.

True. Since $\forall n \in \mathbb{N}$, $A_n \subseteq A_1$ and A_n is finite and nonempty, there exists an x such that $x \in A_n$ which means $x \in A_1$, and so x is in $\bigcap_{n=1}^{\infty} A_n$. Thus, $\bigcap_{n=1}^{\infty} A_n$ is nonempty. Also, $\forall n \in \mathbb{N}$, A_n is finite which means $\bigcap_{n=1}^{\infty} A_n$ is finite since the intersection of finite sets must be finite. Thus, $\bigcap_{n=1}^{\infty} A_n$ is finite and nonempty.

- (c) $A \cap (B \cup C) = (A \cap B) \cup C$

False, let $A = \{0\}$, $B = \{1\}$, $C = \{2\}$. Then, $A \cap (B \cup C) = \{0\} \cap (\{1\} \cup \{2\}) = \{0\} \cap \{1, 2\} = \emptyset$. But, $(A \cap B) \cup C = (\{0\} \cap \{1\}) \cup \{2\} = \emptyset \cup \{2\} = \{2\}$. $\emptyset \neq \{2\}$.

- (d) $A \cap (B \cap C) = (A \cap B) \cap C$

True, because set intersection is associative.

Proof. By the definition of set intersection, $A \cap (B \cap C)$ is equivalent to $(x \in A) \wedge (x \in B \wedge x \in C)$. Since conjunction is associative, this becomes $(x \in A \wedge x \in B) \wedge x \in C$ which is equivalent to $(A \cap B) \cap C$. \square

(e) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

True, because set intersection is distributive over set union.

Proof. By the definition of set union and set intersection, $A \cap (B \cup C)$ is equivalent to $(x \in A) \wedge (x \in B \vee x \in C)$. By the distributive rule for conjunction over disjunction, this becomes $(x \in A \wedge x \in B) \vee (x \in A \wedge x \in C)$ which is equivalent to $(A \cap B) \cup (A \cap C)$. \square

1.2.10 Let $y_1 = 1$, and for each $n \in \mathbb{N}$ define $y_{n+1} = \frac{3y_n+4}{4}$.

(a) Use induction to prove that the sequence satisfies $y_n < 4$ for all $n \in \mathbb{N}$.

Proof.

Base Case ($n = 1$): $y_{n+1} = y_2 = \frac{3(1)+4}{4} = \frac{7}{4} < 4$.

Induction Step:

- * Suppose $k \in \mathbb{N}$ such that $k \geq 2$.
- * Assume that for all natural numbers $i < k$, $y_{i+1} < 4$.
- * Need to prove that $y_k < 4$, that is $\frac{3y_{k-1}+4}{4} = \frac{3}{4}y_{k-1} + 1 < 4$. By the induction hypothesis, $y_{k-1} < 4$. So, $\frac{3}{4}y_{k-1} < 3$ which means $\frac{3}{4}y_{k-1} + 1 < 4$. So, $y_k < 4$.

Hence, by the principle of complete induction, $\forall n \in \mathbb{N}, y_n < 4$. \square

(b) Use another induction argument to show the sequence (y_1, y_2, y_3, \dots) is increasing.

Proof. We'll write $p(n)$ to denote the statement " $y_n \leq y_{n+1}$ ". Need to prove that $\forall n \in \mathbb{N}, p(n)$.

Base Case ($n = 1$): Then, $y_n = y_1 = 1$ and $y_{n+1} = y_2 = \frac{7}{4}$. Clearly, $1 \leq \frac{7}{4}$.

Induction Step:

- * Suppose $k \in \mathbb{N}$ such that $k \geq 2$.
- * Assume that for all natural numbers $i < k$, $p(i)$ is true.
- * Need to prove that $p(k)$ holds true, that is $y_k \leq y_{k+1}$. By the definition of y , $y_k = \frac{3y_{k-1}+4}{4}$ and $y_{k+1} = \frac{3y_k+4}{4}$. Need to show that $\frac{3y_{k-1}+4}{4} \leq \frac{3y_k+4}{4}$, or in more simplified terms $y_{k-1} \leq y_k$. By the induction hypothesis, $p(k-1)$ is true, that is $y_{k-1} \leq y_k$. So, $\frac{3y_{k-1}+4}{4} \leq \frac{3y_k+4}{4}$ which means $y_k \leq y_{k+1}$. Thus, $p(k)$ holds true.

Hence, by the principle of complete induction, $\forall n \in \mathbb{N}, p(n)$ is true. \square

1.3.2 (a) Write a formal definition in the style of Definition 1.3.2 for the infimum or greatest lower bound of a set.

A real number s is an infimum for a set $A \subseteq \mathbb{R}$ if it meets these two criteria:

- (i) s is a lower bound for A
- (ii) if b is any lower bound for A , then $s \geq b$

- (b) Now, state and prove a version of Lemma 1.3.7 for greatest lower bounds.

Assume $s \in \mathbb{R}$ is a lower bound for the set $A \subseteq \mathbb{R}$. Then, $s = \inf A$ iff for every choice of $\epsilon > 0$, there exists an element $a \in A$ satisfying $s + \epsilon > a$.

Proof.

(\Rightarrow) Assume $s = \inf A$. Need to show that $\forall \epsilon > 0, \exists a \in A$ such that $s + \epsilon > a$. Assume $\epsilon > 0$. Since s is an infimum for A and $s + \epsilon > s$, $s + \epsilon$ cannot be a lower bound for A which means that $\exists a \in A$ such that $a < s + \epsilon$ (because otherwise $s + \epsilon$ would be a lower bound). Thus, the claim is satisfied.

(\Leftarrow) Assume s is a lower bound for A such that $\forall \epsilon > 0, s + \epsilon$ is not a lower bound for A . So, for all lower bounds $l, l \leq s$. This satisfies both parts of the definition of $s = \inf A$, that is s is a lower bound and for all lower bounds $l, l \leq s$. \square

- 1.3.8 If $\sup A < \sup B$, then show that there exists an element $b \in B$ that is an upper bound for A .

Proof. Assume $\sup A < \sup B$. Need to show that $\exists b \in B$ such that b is an upper bound for A . Let $\epsilon = \sup B - \sup A > 0$. Then, we know that $\exists b \in B$ such that $b > \sup B - \epsilon = \sup A$. Thus, since $b > \sup A$, b is an upper bound for A . \square

- 1.3.9 Without worrying about formal proofs for the moment, decide if the following statements about suprema and infima are true or false. For any that are false, supply an example where the claim in question does not appear to hold.

- (a) A finite, nonempty set always contains its supremum.

True, the last element of the set is the supremum.

- (b) If $a < L$ for every element a in the set A , then $\sup A < L$.

False, let $A = (0, 1)$ which means $\sup A = 1$. Let $L = 1$. Then, $\forall a \in A, a < L$, but $\sup A \not< L$.

- (c) If A and B are sets with the property that $a < b$ for every $a \in A$ and every $b \in B$, then it follows that $\sup A < \inf B$.

False, let $A = (0, 1)$ and $B = (1, 2)$. Then, $\forall a \in A$ and $\forall b \in B, a < b$, but $\sup A = 1 = \inf B$.

- (d) If $\sup A = s$ and $\sup B = t$, then $\sup(A + B) = s + t$. The set $A + B$ is defined as $A + B = \{a + b : a \in A \text{ and } b \in B\}$.

True

- (e) If $\sup A \leq \sup B$ then there exists a $b \in B$ that is an upper bound for A .

False, let $A = [0, 1]$ and $B = (0, 1)$. Then, $\sup A = \sup B = 1$. But, there is no $b \in B$ such that b is an upper bound for A .

1.4.2 Recall that \mathbb{I} stands for the set of irrational numbers.

- (a) Show that if $a, b \in \mathbb{Q}$, then ab and $a + b$ are elements of \mathbb{Q} as well.

Proof. Assume $a, b \in \mathbb{Q}$. Then, a can be represented as $\frac{p}{q}$ for some $p, q \in \mathbb{Z}$ and b can be represented as $\frac{r}{s}$ for some $r, s \in \mathbb{Z}$. Then, $a + b = \frac{p}{q} + \frac{r}{s} = \frac{ps+qr}{qs}$. Since \mathbb{Z} is closed under addition and multiplication, $ps + qr \in \mathbb{Z}$ and $qs \in \mathbb{Z}$, which means that $\frac{ps+qr}{qs} \in \mathbb{Q}$. Similarly, $ab = \frac{p}{q} * \frac{r}{s} = \frac{pr}{qs} \in \mathbb{Q}$. Thus, $a + b \in \mathbb{Q}$ and $ab \in \mathbb{Q}$. \square

- (b) Show that if $a \in \mathbb{Q}$ and $t \in \mathbb{I}$, then $a + t \in \mathbb{I}$ and $at \in \mathbb{I}$ as long as $a \neq 0$.

Proof. Suppose $a \in \mathbb{Q}$ and $t \in \mathbb{I}$. Assume for the sake of contradiction that $a + t \in \mathbb{Q}$. Then, $t = (-a) + (a + t)$. Since \mathbb{Q} is closed under addition, this means $t \in \mathbb{Q}$. However, this contradicts the initial claim $t \in \mathbb{I}$. Thus, $a + t \in \mathbb{I}$. \square

Proof. Suppose $a \in \mathbb{Q}$ such that $a \neq 0$ and $t \in \mathbb{I}$. Assume for the sake of contradiction that for $at \in \mathbb{Q}$. Then, $t = (\frac{1}{a}) * (at)$. Since \mathbb{Q} is closed under multiplication, this means that $t \in \mathbb{Q}$. However, this contradicts the initial claim $t \in \mathbb{I}$. Thus, $at \in \mathbb{I}$. \square

- (c) Part (a) can be summarized by saying that \mathbb{Q} is closed under addition and multiplication. Is \mathbb{I} closed under addition and multiplication? Given two irrational numbers s and t , what can we say about $s + t$ and st ?

\mathbb{I} is not closed under addition nor is it closed under multiplication. Given $s, t \in \mathbb{I}$, $s + t$ and st can be in \mathbb{I} or not in \mathbb{I} depending on the specific values of s and t . For example, let $s = \sqrt{2}$ and $t = -\sqrt{2}$. Then, $s + t = 0 \notin \mathbb{I}$ and $st = -2 \notin \mathbb{I}$. But now let $s = \sqrt{2}$ and $t = \sqrt{3}$. Then, $s + t = (\sqrt{2} + \sqrt{3}) \in \mathbb{I}$ and $st = \sqrt{6} \in \mathbb{I}$.

- 1.4.6 (a) Finish the proof of Theorem 1.4.5 by showing that the assumption $\alpha^2 > 2$ leads to a contradiction of the fact that $\alpha = \sup T$.

Proof. Assume for the sake of contradiction that $\alpha^2 > 2$. Then, we get

$$\begin{aligned} \left(\alpha - \frac{1}{n}\right)^2 &= \alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} \\ &> \alpha^2 - \frac{2\alpha}{n}. \end{aligned}$$

Choose $n_0 \in \mathbb{N}$ such that $\alpha^2 - \frac{2\alpha}{n_0} > 2$ (this is possible by the Archimedean Property and since $\alpha^2 > 2$). So, $(\alpha - \frac{1}{n_0})^2 > \alpha^2 - \frac{2\alpha}{n_0} > 2$. This means $\alpha - \frac{1}{n_0}$ must be an upper bound for T . However, $\alpha - \frac{1}{n_0} < \alpha$ which contradicts the initial claim that $\alpha = \sup T$. So, $\alpha^2 \not> 2$. Since, part 1 of the proof showed that $\alpha^2 \not< 2$, the only conclusion left is that $\alpha^2 = 2$ which means $\alpha = \sqrt{2} \in \mathbb{R}$. \square

(b) Modify this argument to prove the existence of \sqrt{b} for any real number $b \geq 0$.

Proof. Suppose $b \in \mathbb{R}_{\geq 0}$. Let $T = \{t \in \mathbb{R} : t^2 < b\}$. Need to prove that $(\sup T)^2 = b$, that is $(\sup T)^2 \not< b$ and $(\sup T)^2 \not> b$.

Assume for the sake of contradiction that $\alpha = \sup T$ and $\alpha^2 < b$. In search of an element of T that is greater than α , we get

$$\begin{aligned} \left(\alpha + \frac{1}{n}\right)^2 &= \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n^2} \\ &< \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n} \\ &= \alpha^2 + \frac{2\alpha + 1}{n}. \end{aligned}$$

Choose $n_0 \in \mathbb{N}$ such that $\alpha^2 + \frac{2\alpha+1}{n_0} < b$. So, $(\alpha + \frac{1}{n_0})^2 < \alpha^2 + \frac{2\alpha+1}{n_0} < b$. This means $\alpha + \frac{1}{n_0} \in T$ and is an upper bound for T . However, $\alpha + \frac{1}{n_0} > \alpha$ which contradicts the initial claim that $\alpha = \sup T$ since a supremum of a set should be larger than all elements within that set. So, $\alpha^2 \not< b$.

Assume for the sake of contradiction that $\alpha = \sup T$ and $\alpha^2 > b$. Then, we get

$$\begin{aligned} \left(\alpha - \frac{1}{n}\right)^2 &= \alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} \\ &> \alpha^2 - \frac{2\alpha}{n}. \end{aligned}$$

Choose $n_0 \in \mathbb{N}$ such that $\alpha^2 - \frac{2\alpha}{n_0} > b$ (this is possible by the Archimedean Property and since $\alpha^2 > 2$). So, $(\alpha - \frac{1}{n_0})^2 > \alpha^2 - \frac{2\alpha}{n_0} > b$. This means $\alpha - \frac{1}{n_0}$ must be an upper bound for T . However, $\alpha - \frac{1}{n_0} < \alpha$ which contradicts the initial claim that $\alpha = \sup T$ since a supremum of a set should be the smallest element amongst all the upper bounds for that set. So, $\alpha^2 \not> b$.

Since $(\sup T)^2 \not< b$ and $(\sup T)^2 \not> b$, $(\sup T)^2 = b$ and so $\sup T = \sqrt{b} \in \mathbb{R}$. □

1.4.8 Use the following outline to supply proofs for the statements in Theorem 1.4.13.

- (a) First, prove statement (i) for two countable sets, A_1 and A_2 . Example 1.4.8 (ii) may be a useful reference. Some technicalities can be avoided by first replacing A_2 with the set $B_2 = A_2 - A_1 = \{x \in A_2 : x \notin A_1\}$. The point of this is that the union $A_1 \cup B_2$ is equal to $A_1 \cup A_2$ and the sets A_1 and B_2 are disjoint. (What happens if B_2 is finite?)

- (i) If A_1 and A_2 are countable sets, then the union $A_1 \cup A_2$ is countable.

Proof. Assume A_1 and A_2 are countable sets. Let $B_2 = A_2 - A_1 = \{x \in A_2 : x \notin A_1\}$ which means that $A_1 \cup A_2 = A_1 \cup B_2$ and the sets A_1 and B_2 are disjoint. We need to show that $A_1 \cup B_2$ is countable. Since, A_1 is countable, there exists a bijective function $f : \mathbb{N} \rightarrow A_1$. For B_2 , there are 3 cases to consider: $B_2 = \emptyset$, B_2 is a nonempty and finite set, and B_2 is an infinite set.

Case 1: Assume $B_2 = \emptyset$. Then, $A_1 \cup B_2 = A_1$ and since A_1 is countable, it would mean that $A_1 \cup B_2$ is countable.

Case 2: Assume B_2 is a nonempty, finite set defined as $B_2 = \{b_1, b_2, \dots, b_i\}$ for some $i \in \mathbb{N}$. Need to prove that there exists a bijective function $g : \mathbb{N} \rightarrow (A_1 \cup B_2)$. Let $g : \mathbb{N} \rightarrow (A_1 \cup B_2)$ be defined in the following manner: $\forall n \in \mathbb{N}$, $n \leq i \implies g(n) = b_n$ and $n > i \implies g(n) = f(n - i)$. Since f is a bijection and A_1 and B_2 are disjoint, g must also be a bijection.

Case 3: Assume B_2 is an infinite set. Since, $B_2 \subseteq A_2$ and A_2 is a countable set, B_2 must be a countably infinite set (since its not finite or nonempty) which means that there exists a bijective function $g : \mathbb{N} \rightarrow B_2$. We need to prove that there exists a bijective function $h : \mathbb{N} \rightarrow (A_1 \cup B_2)$. Let $h : \mathbb{N} \rightarrow (A_1 \cup B_2)$ be defined in the following manner: $\forall n \in \mathbb{N}$, n is odd $\implies h(n) = f(\frac{n+1}{2})$ and n is even $\implies h(n) = g(\frac{n}{2})$. Then, since both f and g are bijective functions and A_1 and B_2 are disjoint, h must also be a bijective function. \square

Now, explain how the more general statement in (i) follows.

To prove the more general statement about the union of m countable sets, we would need to apply induction. The proof above would serve as the base case. Then, we would need to assume that the union of $m - 1$ countable sets is countable and show that the union of m countable sets is also countable.

- (b) Explain why induction cannot be used to prove part (ii) of Theorem 1.4.13 from part (i).

Induction cannot be used to prove part (ii) of Theorem 1.4.13 from part (i) because it can only be used on $n \in \mathbb{N}$ and $\infty \notin \mathbb{N}$.

(c) Show how arranging \mathbb{N} into the two-dimensional array

1	3	6	10	15	...
2	5	9	14	...	
4	8	13	...		
7	12	...			
11	...				
...					

leads to a proof of Theorem 1.4.13 (ii).

Proof. Assume A_n is a countable set for each $n \in \mathbb{N}$. Need to prove $\bigcup_{n=1}^{\infty} A_n$ is countable. In order to ensure that the sets in the union are disjoint, let $B_1 = A_1$, $B_2 = A_2 - A_1$, $B_3 = A_3 - (A_1 \cup A_2)$, $B_4 = A_4 - (A_1 \cup A_2 \cup A_3)$, ..., $B_n = A_n - \bigcup_{i=1}^{n-1} A_i$ where $n \in \mathbb{N}$. Then, organize the elements of $\bigcup_{i=1}^{\infty} B_i$ into a two-dimensional array similar to above where b_{rc} is the element at the r 'th row and c 'th column. So,

$B_1 = b_{11}$	b_{12}	b_{13}	b_{14}	b_{15}	...
$B_2 = b_{21}$	b_{22}	b_{23}	b_{24}	...	
$B_3 = b_{31}$	b_{32}	b_{33}	...		
$B_4 = b_{41}$	b_{42}	...			
$B_5 = b_{51}$...				
...					

Now, it is clear to see that every element in both arrays will be uniquely mapped, that is there exists a bijective function $f : \mathbb{N} \rightarrow \bigcup_{i=1}^{\infty} B_i$. Specifically, for any $x \in \mathbb{N}$, $f(x) = b_{pq}$ where p and q are the row and column (respectively) of $x \in \mathbb{N}$ in the diagram for \mathbb{N} above. Thus, $\mathbb{N} \sim \bigcup_{i=1}^{\infty} B_i$ so $\bigcup_{i=1}^{\infty} B_i$ is countable which means $\bigcup_{n=1}^{\infty} A_n$ is countable. \square