

Test Results

Patient: Grace Wilson | DOB: 1963-12-19 | Patient ID: P005
Collected: 2025-06-20 | Source: Patient portal sample export


COMPREHENSIVE METABOLIC PANEL (CMP)

Results

Sodium

[View trends](#)


Normal range: 135 - 145 mmol/L



Potassium

[View trends](#)


Normal range: 3.5 - 5.0 mmol/L



Chloride

[View trends](#)

Normal range: 98 - 108 mmol/L



Carbon Dioxide (CO2)


[View trends](#)

Normal range: 21 - 30 mmol/L

Carbon Dioxide (CO2)

[View trends](#)

Normal range: 21 - 30 mmol/L



Urea Nitrogen (BUN)


[View trends](#)

Normal range: 7 - 20 mg/dL

Urea Nitrogen (BUN)

[View trends](#)

Normal range: 7 - 20 mg/dL



Creatinine


[View trends](#)

Normal range: 0.4 - 1.0 mg/dL

Creatinine

[View trends](#)

Normal range: 0.4 - 1.0 mg/dL



Glucose


[View trends](#)

Normal range: 70 - 140 mg/dL

Glucose

[View trends](#)

Normal range: 70 - 140 mg/dL



Interpretive Data:

Above is the NONFASTING reference range.

Below are the FASTING reference ranges:

NORMAL: 70-99 mg/dL


PREDIABETES: 100-125 mg/dL

DIABETES: > 125 mg/dL

Calcium

[View trends](#)


Normal range: 8.7 - 10.2 mg/dL



AST (Aspartate Aminotransferase)

[View trends](#)


Normal range: 15 - 41 U/L



ALT (Alanine Aminotransferase)

[View trends](#)

Normal range: 10 - 39 U/L



Bilirubin, Total


[View trends](#)

Normal range: 0.4 - 1.5 mg/dL

Bilirubin, Total

[View trends](#)

Normal range: 0.4 - 1.5 mg/dL



Alk Phos (Alkaline Phosphatase)


[View trends](#)

Normal range: 24 - 110 U/L

Alk Phos (Alkaline Phosphatase)

[View trends](#)

Normal range: 24 - 110 U/L



Albumin


[View trends](#)

Normal range: 3.5 - 4.8 g/dL

Albumin

[View trends](#)

Normal range: 3.5 - 4.8 g/dL



Protein, Total


[View trends](#)

Normal range: 6.2 - 8.1 g/dL

Protein, Total

[View trends](#)

Normal range: 6.2 - 8.1 g/dL



Anion Gap


[View trends](#)

Normal range: 3 - 12 mmol/L

Anion Gap

[View trends](#)

Normal range: 3 - 12 mmol/L



BUN/CREA Ratio


[View trends](#)

Normal range: 6 - 27

BUN/CREA Ratio

[View trends](#)

Normal range: 6 - 27



Glomerular Filtration Rate (eGFR)


[View trends](#)

Normal range: > 60 mL/min/1.73 sq m

Glomerular Filtration Rate (eGFR)

[View trends](#)

Normal range: > 60 mL/min/1.73 sq m



CKD-EPI (2021) does not include patient's race in the calculation of eGFR. Monitoring changes of plasma creatinine and eGFR over time is useful for monitoring kidney function. This change was made on 3/1/2022.

Interpretive Ranges for eGFR(CKD-EPI 2021):

eGFR: > 60 mL/min/1.73 sq m - Normal

eGFR: 30 - 59 mL/min/1.73 sq m - Moderately Decreased

eGFR: 15 - 29 mL/min/1.73 sq m - Severely Decreased

eGFR: < 15 mL/min/1.73 sq m - Kidney Failure

Note: These eGFR calculations do not apply in acute situations when eGFR is changing rapidly or in patients on dialysis.


Lipid panel:

Results

Cholesterol, Total

[View trends](#)

Normal range: below 200 mg/dL



Comment: The significance of total cholesterol depends on the values of individual components including HDL, LDL, non-HDL, and triglycerides. A fasting or non-fasting lipid panel, including total, HDL, and LDL cholesterol, and triglycerides, is recommended to screen and monitor lipid-related risk.

Desirable: <200 mg/dL

Borderline High: 200-239 mg/dL


High: ≥ 240 mg/dL

Ref: NCEP guidelines

LDL Calculated

[View trends](#)

Normal range: below 100 mg/dL



<70 mg/dL Desired target for prior heart disease, stroke, and those at high-risk. Even lower levels may be recommended to decrease the risk of heart attack and stroke.

70-159 mg/dL Comprehensive cardiovascular risk assessment is recommended. Statin therapy may be advised based on risk factors.

Desirable: <100 mg/dL

Above desirable: 100-129 mg/dL

Borderline high: 130-159 mg/dL

160-189 mg/dL Moderately elevated LDL level. Statin therapy is recommended if other risk factors are present.

≥ 190 mg/dL Severely elevated LDL level. High long-term risk of heart disease and stroke. High-intensity statin therapy is recommended for most people. Consider specialist referral.

*A healthy diet and exercise are recommended for all to reduce heart disease risk. Statin choice should be based on patient preference after patient-provider discussions.

Ref: 2018 ACC/AHA and the NCEP guidelines

HDL


[View trends](#)

Normal range: 40 - 80 mg/dL

HDL

[View trends](#)

Normal range: 40 - 80 mg/dL



People with low HDL levels are at increased risk of heart disease:

<50 mg/dL for Women

<40 mg/dL for Men

Triglyceride


[View trends](#)

Normal range: below 150 mg/dL

Triglyceride

[View trends](#)

Normal range: below 150 mg/dL



<150 mg/dL Normal

150-499 mg/dL High Triglycerides. Risk of heart disease may be increased. Address reversible causes (eg sugar in foods and beverages, alcohol, and diabetes control). Medication may be appropriate based on other clinical factors.

>=500 mg/dL Very High Triglycerides. Risk of heart disease and pancreatitis increased. Address reversible causes as above. Medication to lower triglycerides usually advised.

*Ranges provided for adults, pediatric guidelines vary.

VITAMIN D 25 HYDROXY

Results

Bone and Mineral Metabolism: Vitamin D

25 OH Vitamin D Status

<10 ng/mL Deficiency

10-30 ng/mL Insufficiency


30-100 ng/mL Sufficiency

>100 ng/mL Toxicity

Vitamin D Total, 25OH

[View trends](#)

Normal range: 30 - 100 ng/ml



HEMOGLOBIN A1C

Results

Between 5.7% and 6.4% is suggestive of Pre-Diabetes or controlled Diabetes.
Greater than or equal to 6.5% is suggestive of Diabetes, and if more than one value, diagnostic.

Accuracy may be reduced by anemia, hemoglobinopathy, recent transfusion, sickle cell, artificial heart valve, dialysis, TIPS, severe hyperglycemia, etc.

Hemoglobin A1C

[View trends](#)


Normal range: below 5.7 %



Average Blood Glucose (Calculated From HgBA1c Level)

[View trends](#)

Normal range: 70 - 117 mg/dL



Thyroid Hormone Tests


Collected on Jun 17, 2024 9:38 AM

Results

Thyroid Stimulating Hormone (TSH)

[View trends](#)


Normal range: 0.34 - 5.66 uIU/mL



Thyroxine, Free (FT4)

[View trends](#)

Normal range: 0.52 - 1.21 ng/dL



VITAMIN B12


Collected on Jun 17, 2024 9:38 AM

Results

Vitamin B12

[View trends](#)

Normal range: 180 - 914 pg/mL



IRON AND TOTAL IRON BINDING CAPACITY (TIBC)


Collected on Jun 17, 2024 9:38 AM

Results

Iron

[View trends](#)


Normal range: 28 - 170 mcg/dL



Total Iron Binding Capacity (TIBC)

[View trends](#)


Normal range: 261 - 478 mcg/dL



Percent Transferrin Saturation

[View trends](#)

Normal range: 15 - 55 %



Additional information

Synthetic sample data for software testing only. Not a real medical record.