

# Supplementary Materials: Derivation of a Dynamic Model for Palmitate-induced NF $\kappa$ B Signaling Pathway through Systems Biology Approach

Dongheon Lee, Yufang Ding, Arul Jayaraman, and Joseph Sang-Il Kwon

TABLE I: Model Equations for the AMPK-SIRT1 regulation model

$$\begin{aligned}
 \frac{d NAD^+}{dt} &= k_{gn} + k_{gne} \frac{pAMPK}{pAMPK + K_{gne}} - 7k_{bo} \cdot pAMPK \frac{Pal}{Pal + K_{bo}} - k_{c1} \cdot NAD^+ + k_{c2} \cdot NADH - NAD^+ \frac{v_{dnad}}{NAD^+ + K_{dnad}} - k_{dn} \cdot NAD^+ \\
 \frac{d NADH}{dt} &= 7k_{bo} \cdot pAMPK \frac{Pal}{Pal + K_{bo}} + k_{c1} \cdot NAD^+ - k_{c2} \cdot NADH \\
 \frac{d Pal_o}{dt} &= -k_{tran} \cdot Pal_o \\
 \frac{d Pal}{dt} &= -k_{tran} \cdot Pal_o - k_{bo} \cdot pAMPK \frac{pAMPK}{pAMPK + K_{bo}} - k_{dc} \cdot \frac{Pal}{Pal + K_{dc}} + k_{ps} \\
 \frac{d Cer}{dt} &= k_{dc} \frac{Pal}{Pal + K_{dc}} - k_{sc} \cdot Cer + k_{cs} \\
 \frac{d pAMPK}{dt} &= -(k_{aa} + K_{1max} \frac{Cer}{Cer + K_1}) \cdot PP2A \frac{pAMPK}{pAMPK + K_M} - k_{a2c} \cdot PP2C \frac{pAMPK}{pAMPK + K_M} \\
 &\quad + (k_p + k_{spmax} \frac{SIRT1}{SIRT1 + K_{sp}}) \cdot (AMPK_t - pAMPK) \\
 \frac{d PP2C}{dt} &= k_{syn} + k_{synmax} \frac{TNFRa}{TNFRa + K_2} - k_{d1} \cdot PP2C \\
 \frac{d NF\kappa Ba}{dt} &= -a_1 \cdot k_v \cdot I\kappa Bn \cdot NF\kappa Ba + k_{ac} \cdot NF\kappa Bn - (k_{da} + k_{damax} \frac{SIRT1}{SIRT1 + K_{da}}) NF\kappa Ba \\
 \frac{d NF\kappa Bn}{dt} &= i_1 NF\kappa B - a_1 \cdot k_v \cdot I\kappa Bn - k_{ac} \cdot NF\kappa Bn + (k_{da} + k_{damax} \frac{SIRT1}{SIRT1 + K_{da}}) NF\kappa Ba \\
 \frac{d NF\kappa Bn - I\kappa Bn}{dt} &= a_1 \cdot k_v \cdot I\kappa Bn (NF\kappa Bn + NF\kappa Ba) - e_{2a} * NF\kappa Bn - I\kappa Ban
 \end{aligned}$$

TABLE II: Model parameters in the AMPK-SIRT1 regulation model

$k_{gn}$	constitutive $NAD^+$ generation rate constant	$k_{gne}$	rate constant of $NAD^+$ generation due to pAMPK
$K_{gne}$	Michaelis constant of pAMPK-induced $NAD^+$ generation	$k_{bo}$	palmitate oxidation rate constant
$K_{bo}$	Michaelis constant of palmitate oxidation	$k_{c1}$	rate constant for conversion from $NAD^+$ to NADH
$k_{c2}$	rate constant for conversion from NADH to $NAD^+$	$v_{dnad}$	maximum degradation rate for $NAD^+$
$K_{dnad}$	Michaelis constant for $NAD^+$ degradation	$k_{dn}$	nonspecific degradation rate constant
$k_{tran}$	rate constant of palmitate translocation to cytoplasm	$k_{dc}$	ceramide generation rate constant from palmitate
$K_{dc}$	Michaelis constant for ceramide degradation	$k_{ps}$	constitutive palmitate synthesis rate constant
$k_{sc}$	ceramide degradation rate constant	$k_{cs}$	constitutive ceramide synthesis rate constant
$k_{aa}$	PP2A-induced AMPK dephosphorylation rate constant	$K_{1max}$	maximum AMPK dephosphorylation rate due to ceramide
$K_1$	Michaelis constant for Cer-induced AMPK dephosphorylation	$k_{a2c}$	PP2C-induced AMPK dephosphorylation rate constant
$K_M$	Michaelis constant for PP2C-induced AMPK dephosphorylation	$k_p$	constitutive AMPK phosphorylation rate constant
$k_{spmax}$	SIRT1-induced AMPK phosphorylation rate constant	$K_{sp}$	Michaelis constant for SIRT1-induced AMPK phosphorylation
$k_{syn}$	constitutive PP2C synthesis rate constant	$k_{synmax}$	maximum PP2C synthesis rate enhanced by TNFRa
$K_2$	Michaelis constant for TNFRa-induced PP2C synthesis	$k_{d1}$	PP2C degradation rate constant
$k_{da}$	constitutive NF $\kappa$ B deacetylation rate	$k_{damax}$	maximum SIRT1-induced NF $\kappa$ B deacetylation rate
$K_{da}$	Michaelis constant for SIRT1-induced NF $\kappa$ B deacetylation	PP2A	the number of PP2A molecules