

OZONE'S POWER OVER PATHOGENS

Pathogen	<u>Dosage</u>
Aspergillus Niger (Black Mount)	Destroyed by 1.5 to 2 mg/l
Bacillus Bacteria	Destroyed by 0.2 m/l within 30 seconds
Bacillus Anthracis (causes anthrax in sheep, cattle and pigs. Also a human pathogen)	Ozone susceptible
Bacillus cereus	99% destruction after 5-min at 0.12 mg/l in water
B. cereus (spores)	99% destruction after 5-min at 2.3 mg/l in water
Bacillus subtilis	90% reduction at 0.10-PPM for 33 minutes
Bacteriophage f2	99.99% destruction at 0.41 mg/l for 10-seconds in water
Botrytis cinerea	3.8 mg/l for 2 minutes
c. difficile	99.999% destruction at 0.6 mg/l for 2 minutes (in water)
Candida Bacteria	Ozone susceptible
Clavibacter michiganense	99.99% destruction at 1.1 mg/l for 5 minutes
Cladosporium	90% reduction at 0.10-PPM for 12.1 minutes
Clostridium Bacteria	Ozone susceptible
Clostridium Botulinum Spores. Its toxin paralyses the central nerve system, being a poison multiplying in food and meals.	0.4 to 0.5 mg/l threshold value
Coxsackie Virus A9	95% destruction at 0.035 mg/l for 10-seconds in water
Coxsackie Virus B5	99.99% destruction at 0.4 mg/l for 2.5-minutes in sludge effluent
Diphtheria Pathogen	Destroyed by 1.5 to 2 mg/l
Eberth Bacillus (Typhus abdomanalis). Spreads typically by aqueous infection and causes typhoid.	Destroyed by 1.5 to 2 mg/l
Echo Virus 29: The virus most sensitive to ozone.	After a contact time of 1 minute at 1 mg/l of ozone, 99.999% killed.
Enteric virus	95% destruction at 4.1 mg/l for 29 minutes in raw wastewater
Escherichia Coli Bacteria (from feces)	Destroyed by 0.2 mg/l within 30 seconds in air
E-coli (in clean water)	99.99% destruction at 0.25 mg/l for 1.6 minutes
E-coli (in wastewater)	99.9% destruction at 2.2 mg/l for 19 minutes
Encephalomyocarditis Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l.
Endamoebic Cysts Bacteria	Ozone susceptible
Enterovirus Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l.
Fusarium oxysporum f.sp. lycopersici	1.1 mg/l for 10 minutes
Fusarium oxysporum f.sp. melonogea	99.99 % destruction at 1.1 mg/l for 20 minutes
GDVII Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l.
Hepatitis A virus	99.5% reduction at 0.25 mg/l for 2-seconds in a phosphate buffer
Herpes Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l.
Influenza Virus	0.4 to 0.5 mg/l threshold value
Klebs-Loffler Bacillus	Destroyed by 1.5 to 2 mg/l
Legionella pneumophila	99.99% destruction at 0.32 mg/l for 20 minutes in distilled water
Luminescent Basidiomycetes (species having no melanin pigment).	Destroyed in 10 minutes at 100-PPM
Mucor piriformis	3.8 mg/l for 2 minutes
Mycobacterium avium	99.9% with a CT value of 0.17 in water (scientifically reviewed document)
Mycobacterium foruitum	90% destruction at 0.25 mg/l for 1.6 minutes in water
Penicillium Bacteria	Ozone susceptible
Phytophthora parasitica	3.8 mg/l for 2 minutes
Poliomyelitis Virus	99.99% kill with 0.3 to 0.4 mg/l in 3-4 minutes
Poliovirus type 1	99.5% destruction at 0.25 mg/l for 1.6 minutes in water
Proteus Bacteria	Very susceptible
Pseudomonas Bacteria	Very susceptible
Rhabdovirus virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l
Salmonella Bacteria	Very susceptible
Salmonella typhimurium	99.99% destruction at 0.25 mg/l for 1.67 minutes in water
Schistosoma Bacteria	Very susceptible
Staph epidermidis	90% reduction at 0.1-ppm for 1.7 min
Staphylococci	Destroyed by 1.5 to 2.0 mg/l
Stomatitis Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l
Streptococcus Bacteria	Destroyed by 0.2 mg/l within 30 seconds
Verticillium dahlia	99.99 % destruction at 1.1 mg/l for 20 minutes
Vesicular Virus	Destroyed to zero level in less than 30 seconds with 0.1 to 0.8 mg/l
Virbrio Cholera Bacteria	Very susceptible
Vicia Faba progeny	Ozone causes chromosome aberration and its effect is twice that observed by the action of X-rays