

Linkage Tests in Upland Cotton, *Gossypium hirsutum* L. II.¹R. J. Kohel²

ABSTRACT

Results of linkage analyses of cotton performed at College Station, Texas are presented, and linkage information reported by other workers is summarized. The mutants withering bract, accessory involucre, abnormal palisades, Ragged Leaf, round leaf, Red Dwarf, male-sterile-1, male-sterile-2, Male-sterile-4, and Ligon Lintless were tested for linkage with several marker loci and in various combinations among themselves.

Ragged Leaf and round leaf were found to be linked (R.C. = 32%), and thus form a new linkage group, number X. Data are presented to document the simple dominant inheritance of Ligon Lintless; the symbol *Li* is assigned to the mutant allele.

Additional index words: Linkage Group X, Ligon Lintless inheritance.

THIS paper reviews the efforts by researchers at several locations to identify the 26 potential linkage groups of *Gossypium hirsutum* L., and it reports the progress made since the last published linkage report (34).

In the past, researchers at the Texas Station have relied heavily upon our multiple marker lines to expedite tests for linkage associations. These lines are still useful for testing new mutants, although most known mutants have been tested with the multiple marker stocks. Often, as in this report, the only remaining untested combinations in many of our linkage tests involve digenic combinations.

MATERIALS

Our most useful multiple marker stocks are T 582 (*cu*, *fg*, *cl*, *gl*, and *vi*), T 586 (*R*₂, *Lc*, *L*^o, *R*₁, *H*₂, *Y*, *N*, and *P*_a), and T 588 (*R*₂, *L*^o, *Lg*, *R*₁, *H*₂, *Y*, *N*, and *P*_a). Summary information concerning the mutant alleles in the multiple marker lines and the other lines involved in the linkage analyses is presented in Table 1. In this table and also in the text, the conventional method of numbering the chromosomes of *G. hirsutum* is followed in which numbers 1 to 13 are assigned to the A subgenome chromosomes and numbers 14 to 26 are assigned to the D subgenome chromosomes.

The mutants in the linkage tests with which I was primarily concerned were withering bract, accessory involucre, abnormal palisades, Ragged Leaf, round leaf, Red Dwarf, male-sterile-1, male-sterile-2, Male-sterile-4, and Ligon Lintless.

¹ Contribution from the Plant Science Research Division, Agricultural Research Service, U. S. Department of Agriculture, in cooperation with the Texas Agricultural Experiment Station, College Station, Texas. Received June 18, 1971.

² Research Geneticist, Plant Science Research Division, Agricultural Research Service, U. S. Department of Agriculture, College Station, Texas 77843.

Withering bract, initially called deciduous bract, was discovered in 'Stoneville 2B.' The inheritance of this mutant was determined by Knight (35) and it was found to be linked to *gl*₂ by Rhyne (46). Subsequent studies of withering bract revealed that it was controlled by duplicate loci (47). All *G. hirsutum* lines tested were recessive at the second locus (*bw*₂), which accounts for the apparent monogenic control of this mutant. The *bw*₁ locus is located on a chromosome of the A subgenome, and *bw*₂ is located on the homeologous D chromosome.

Accessory involucre is one of a series of mutants (Heritable Abnormality #7) found by H. C. McNamara. It is simply inherited and linked with frego-bract (34). Frego-bract is thought to reside on an A subgenome chromosome.

Abnormal palisades is controlled by duplicate recessive genes (31). The amphidiploid nature of *G. hirsutum* and the experiences of cotton geneticists with other duplicate genes suggest that the controlling loci reside on homeologous chromosomes. Convention places the lower numbered locus (*lp*₂) in the A subgenome and the other (*lp*₂) in the D subgenome.

Ragged Leaf is a simply inherited dominant in which the homozygote is usually lethal (33). Round leaf (3, 51) and crenate (58) are merely two names for the same condition.

Red Dwarf (*Rd*) has a phenotype similar to Red Plant (*R*₁) except that, as its name implies, plants with mutant alleles at the Red Dwarf locus have short stature associated with their red pigmentation (25, 40).

Male-sterile-1 and -2 are simply inherited recessives, whereas Male-sterile-4 is a simply inherited dominant. The chromosomal or subgenomic location of these genes is not known. All three male-steriles differ in their expression. Male-sterile-1 is conditionally male-sterile in that the mutant expression (non-dehiscent anthers) is influenced by environmental conditions. At some locations, it can be identified with a high degree of accuracy, but at College Station the degree of expression varies from year to year. Both male-sterile-2 and Male-sterile-4 impart

Table 1. Name, symbol, and chromosome or subgenome location of mutants involved in linkage tests (references in parentheses).

Name	Symbol	Chromosome or subgenome
abnormal palisades	<i>lp</i> ₁ , <i>lp</i> ₂ (31)	1, 15 D (10)
accessory involucre	<i>ia</i> (32)	
cluster fruiting	<i>cl</i> ₁ (24, 59)	16 (8, 12)
cup leaf	<i>cu</i> (38)	A (13)
frego bract	<i>fg</i> (14)	
glandless boll	<i>gl</i> ₁ (41)	D (12, 48)
male-sterile-1	<i>ms</i> ₁ (26)	
male-sterile-2	<i>ms</i> ₂ (52)	
round leaf	<i>rl</i> (3, 51)	
virescent-1	<i>v</i> ₁ (24, 30)	A (12, 13)
withering bract	<i>bw</i> ₁ , <i>bw</i> ₂ (35)	A, D (47)
Brown Lint	<i>Lc</i> ₁ (22, 24, 50, 61)	7 (11, 12, 27)
Green Lint	<i>Lg</i> (19, 24, 50)	15 (8)
Laetinate Leaf	<i>L</i> ^o	1 (62)
Ligon Lintless	<i>Li</i> (4, text)	D (42)
Male-sterile-4	<i>Ms</i> ₄ (1)	
Naked Seeds	<i>N</i> (5, 15, 19, 24, 29, 59, 60)	A (12)
Okra Leaf	<i>L</i> ^o (20, 24, 53)	15 (8, 12)
Petal Spot	<i>R</i> ₂ (17, 19, 20, 24)	7 (11, 12)
Pilose	<i>P</i> ₁ (55)	6 (6, 12)
Ragged Leaf	<i>Rg</i> (33)	
Red Dwarf	<i>Rd</i> (25, 40)	D (12, 42)
Red Plant	<i>R</i> ₁ (20, 24, 30, 39, 59)	16 (12)
Yellow Petals	<i>Y</i> ₁ (16, 24)	A (57)
Yellow Pollen	<i>P</i> _a (18, 24)	A (56)

complete male-sterility; however, male-sterile-2 results in normal-appearing, but indehiscent anthers, that contain aberrant pollen, whereas Male-sterile-4 varies in its expression from indehiscent anthers to the formation of only rudimentary filaments and anthers.

Ligon Lintless is a mutant that has been known since the 1930's (4) and it has been used extensively as a genetic marker at the Texas Station and elsewhere. It is known to be inherited as a simple complete dominant; however, its inheritance has never been documented in print. This paper will present data to document the inheritance and to assign gene symbols to Ligon Lintless in addition to reporting its performance in linkage tests.

RESULTS

Results of the linkage tests are presented in Table 2. The table contains the recombination values and the population sizes in the various test combinations. The dashes indicate previous tests with independent associations. The associations of *Rg* with *rl* and *fg* with *ia* were the only significant linkage associations detected among the test combinations.

The linkage between *rl* and *Rg* represents a new linkage group, number X, in *G. hirsutum*. Data from the initial test populations indicated that *Rd*, a D genome marker, was linked to *Rg*. However, information from a larger population failed to indicate linkage.

Kohel, Lewis and Richmond (34) identified linkage group VI (*ia-fg*). My data from additional populations of this combination provided a slightly lower estimate of the recombination value than the original one.

Although geneticists agree that Ligon Lintless is a simply inherited dominant mutant, this fact has not been documented, and a gene symbol has not been assigned. Table 3 summarizes the segregation data from those linkage populations reported in this paper and a previous one (34), and the symbol *Li* is assigned to the mutant allele.

Results verified that *Li* is inherited as a simple dominant and also showed that Ligon Lintless plants are less vigorous than normal reflected by the deficient number of plants in the mutant class. The variation in goodness-of-fit to the single gene model was related in part to whether the plants were classified in the greenhouse soon after germination or in the field after they had been transplanted.

Ligon Lintless mutants are characterized by contorted leaf laminae and a twisted appearance of the branches and stems. This expression is first evident in the cotyledonary seedling stage. Another feature characteristic of this mutant is its short lint (approximately 0.2 cm long compared with lint approximately 2.5 cm long in normal cotton). In addition, the homozygous mutant plants appear slightly smaller than the heterozygous ones.

DISCUSSION AND REVIEW

The results of linkage analyses in *G. hirsutum* were summarized in a recent report (2). However, subsequent papers have provided new information; and this information has been incorporated in the current linkage map of *G. hirsutum* presented in Table 4.

Endrizzi and Taylor (11) and Poisson (43) isolated the monosome for chromosome 7, and found that it is associated with linkage group I. As a result of these

Table 2. Summary of linkage tests (the upper number is the recombination percent, and the lower number is the number of plants classified in the test combination).

Testers	Mutants tested									
	bw ₁	ia	lp ₁ lp ₂	itg	rl	Rd	ms ₁	ms ₂	Ms ₄	Li
R ₂	53 100	-	50 176	-	-	-	54 93	-	67 98	
Le ₁				-	-		48 94	-	42 99	
L [*]	47 102	-	56 176	-	-	-	53 95	-	50 98	
Lg	43 99	-	57 176	-	-	-	55 95	-	56 99	
cl ₁	45 192	53 178		-	45 193	56 112		-	44 64	
R ₁	51 102	-	49 176	-	-		58 95	-	55 98	
H ₂	53 102	-	47 176	-	-	-	50 95	-	58 98	
fg	58 192	29.89 572		-	55 196	48 230		-	44 64	
L ^L				43 106	44 98	52 103		47 97		
rl			32.45 607			55 433	45 88			58 112
Rg						54 433	54 93			56 77
Rd								51 95		48 92
Y ₁	44 100	-	48 176	-	-	-		-	46 98	
N	50 99		53 176						60 99	
v ₁	58 193	52 192	50 186	-	43 197	52 112		-	58 64	
cu	57 195	43 192	63 186	-	60 197	55 112		-	53 64	
gl ₁	52 194	58 192	57 186	-	53 197	52 112		-	50 64	
Pa	54 100	-		-	-		55 92	-	54 50	
Li				56 77	58 112	48 92		52 98		

Table 3. Segregation of Ligon Lintless in backcross and F₂ populations.

Population	Number of plants		Chi-square value	P
	Ligon Lintless (Li-)	Normal (lil)		
Backcross (1:1)				
III (III- LiLi)-1	31	45	2.19	.2-.1
-2	45	47	0.10	.5-.7
-3	45	53	0.65	.5-.3
(LiLi- lil)(III-1	75	112	7.32	<.01
-2	48	64	2.28	.2-.1
(III- LiLi)(lil)	88	105	1.50	.3-.2
Pooled	332	426	11.66	<.01
Heterogeneity			2.38	.8-.7
F ₂ (3:1)				
(III- LiLi)⊙-1	133	60	3.82	.1-.05
-2	127	44	0.05	.9-.8
-3	219	86	1.66	.2-.1
Pooled	479	150	.44	.7-.5
Heterogeneity			5.09	.05-.02

Table 4. Linkage map of *Gossypium hirsutum*.

Linkage group	Mutant loci and recombination values	Chromosome or genome location	Reference
I	R ₂ -16-cl ₂ -4-yg ₂ -32-Le ₁	7	11, 21, 27, 43, 44, 45, 54, 58
II	lp ₂ -?-L [*] -32-c*-44-Lg-7-cr	15	9, 10, 58, 63
III	Dw-29-yg ₁ -14-R ₁ -15-cl ₁ -25-c ms ₃ -20-rl-ms ₃ -2-yg ₁	16	9, 21, 44, 45, 28
IV	H ₂ -4-c-18-Le ₂	6	9, 58
V	gl ₂ -33-ne ₁ gl ₂ -15-bw ₁	A	23, 49, 46, 47
VI	ia-30-fg	A	34
VII	L ^L -?-lp ₁	I	10, 62
VIII	ml-41-st ₁	4	7, 62
IX	gl ₃ -39-ne ₂ gl ₃ -20-bw ₂	D	23, 49, 37, 47
X	rl-32-Rg		

* c = centromere of chromosomes 6, 15 and 16.

tests, Endrizzi and Taylor (ibid.) and Kammacher (27) found that the Naked Seed locus was not on chromosome 7 as had been previously reported (58). However, the evidence remains that Naked Seed is an A subgenome mutant (12).

Wilson and Kohel (63) used a translocation stock to provide additional information about the location of L° on chromosome 15 (linkage group II). Results suggested an increase in the overall length of the linkage group. The tentative assignment of lp_2 to this linkage group increases its length even more (see discussion below).

Kammacher and Schwendiman (28) found ms_3 linked to R_1 and yg_1 . This finding adds another identified locus to linkage group III. The homozygous mutant alleles $ms_3 ms_3$ were reported to produce partial male sterility, but Kammacher was able to increase the degree of male sterility through selection (personal communication).

The homeologous linkage groups V (gl_2-bw_1) and IX (gl_3-bw_2) were identified by Lee (36) and by Rhyne (47). Subsequently, Holder, Jenkins and Maxwell (23) found that the duplicate factors for nectariless (ne_1 and ne_2) were also members of these homeologous groups.

Monosomic tests by Endrizzi and Stith (10) have placed lp_1 on chromosome I along with L^L (linkage group VII). Chromosomes 1 and 15 are homeologues; thus, lp_2 should reside on chromosome 15. However, the results of the present study indicate no linkage of lp_2 with L° or Lg . L° and Lg are in opposite arms of chromosome 15 (9), and if lp_2 were in the proximal region of 15, the tests should have detected linkages with one or the other of these marker loci. Therefore, lp_2 must be distal to L° or Lg .

ACKNOWLEDGEMENT

Drs. J. E. Endrizzi and J. A. Lee are acknowledged for their permission to include their unpublished data and their suggestions in the preparation of this paper.

REFERENCES

- Allison, D. C. and W. D. Fisher. 1964. A dominant gene for male-sterility in Upland cotton. *Crop Sci.* 4:548-549.
- Anonymous. 1968. Genetics and cytology of cotton 1956-1967. Southern Coop. Series Bul. 139.
- Brown, H. B. and J. R. Cotton. 1937. "Round leaf" cotton. Notes on the appearance and behavior of a peculiar new strain. *J. Hered.* 28:45-48.
- and J. O. Ware. 1958. Cotton (p. 131). McGraw and Hill Book Co., New York.
- Carver, W. A. 1929. The inheritance of certain seed, leaf, and flower characters in *Gossypium hirsutum* and their genetic interrelations. *J. Amer. Soc. Agron.* 21:467-480.
- Endrizzi, J. E. 1963. Genetic analysis of six primary monosomes and one tertiary monosome in *Gossypium hirsutum*. *Genetics* 48:1625-1633.
- . 1970. Personal communication.
- and M. S. Brown. 1964. Identification of monosomes for six chromosomes in *Gossypium hirsutum*. *Amer. J. Bot.* 51:117-120.
- and R. J. Kohel. 1966. Use of telosomes in mapping three chromosomes in cotton. *Genetics* 54:535-550.
- and L. S. Stith. 1970. Association of two marker genes with chromosome I of cotton. *Agron. Abstr.* p. 9.
- and T. Taylor. 1968. Cytogenetic studies of $N L_1$, yg_2 , R_2 marker genes and chromosome deficiencies in cotton. *Genet. Res.* 12:295-304.
- Gerstel, D. U. and L. L. Phillips. 1958. Segregation of synthetic amphiploids in *Gossypium* and *Nicotiana*. Cold Spring Harbor Symp. Quant. Biol. 23:225-237.
- Giles, J. A. 1962. The comparative genetics of *Gossypium hirsutum* L. and the synthetic amphidiploid, *Gossypium arboreum* L. \times *Gossypium thurberi* Tod. *Genetics* 47:45-49.
- Green, J. M. 1955. Frege bract, a genetic marker in Upland cotton. *J. Hered.* 46:232.
- Griffee, F. and L. L. Ligon. 1929. Occurrence of "lintless" cotton plants and the inheritance of the character "lintless." *J. Amer. Soc. Agron.* 21:711-717.
- Harland, S. C. 1920. Studies of inheritance in cotton. I. The inheritance of corolla colour. *W. Ind. Bull.* 18:13-19.
- . 1929a. The genetics of cotton. I. The inheritance of petal spot in New World cottons. *J. Genet.* 20:365-385.
- . 1929b. The genetics of cotton. II. The inheritance of pollen colour in New World cottons. *J. Genet.* 20:387-399.
- . 1929c. The work of the Genetics Department of the Cotton Research Station, Trinidad. *Emp. Cot. Gr. Rev.* 6:304-314.
- . 1932. The genetics of *Gossypium*. *Bib. Genet.* 9:107-182.
- . 1935. The genetics of cotton. XII. Homologous genes for anthocyanin pigmentation in New and Old World Cottons. *J. Genet.* 30:465-476.
- . 1935. The genetics of cotton. XIV. The inheritance of brown lint in New World cottons. *J. Genet.* 31:27-37.
- Holder, David S., Johnie N. Jenkins, and Fowden G. Maxwell. 1968. Duplicate linkage of glandless and nectariless genes in Upland cotton, *Gossypium hirsutum* L. *Crop Sci.* 8:577-580.
- Hutchinson, J. B. and R. A. Silow. 1939. Gene symbols for use in cotton genetics. *J. Hered.* 30:461-464.
- Johnson, B. L. 1949. Complementary factors for dark-red plant color in Upland cotton. *J. Agr. Res.* 78:535-543.
- Justus, N. and C. L. Leinweber. 1960. A heritable partial male sterile character in cotton. *J. Hered.* 51:191-192.
- Kammacher, P. 1968. New investigations of linkage group I of *Gossypium hirsutum*. *Coton Fib. Trop.* (English Ed.) 23:179-183.
- and J. Schwendiman. 1967. Etude de la localisation chromosomique du gène ms_3 de stérilité pollinique du cotonnier. *Coton Fib. Trop.* 22:417-420.
- Kearney, T. H. and G. J. Harrison. 1927. Inheritance of smooth seeds in cotton. *J. Agr. Res.* 35:193-217.
- Killough, D. T. and W. R. Horlacher. 1933. The inheritance of virescent yellow and red plant characters in cotton. *Genetics* 18:329-334.
- Kohel, R. J. 1964. Inheritance of abnormal palisade mutant in American Upland cotton, *Gossypium hirsutum* L. *Crop Sci.* 4:112-113.
- . 1965. Inheritance of accessory involucre mutant in American Upland cotton, *Gossypium hirsutum* L. *Crop Sci.* 5:119-120.
- and C. F. Lewis. 1962. Inheritance of ragged leaf mutant in American Upland cotton, *Gossypium hirsutum* L. *Crop Sci.* 2:61-62.
- , ———, and T. R. Richmond. 1965. Linkage tests in Upland cotton, *Gossypium hirsutum* L. *Crop Sci.* 5:582-585.
- Knight, R. L. 1962. The genetics of withering or deciduous bracteoles in cotton. *J. Genet.* 50:392-395.
- Lee, J. A. 1965. The genomic allocation of the principal foliar-gland loci in *Gossypium hirsutum* and *Gossypium barbadense*. *Evolution* 19:182-188.
- . 1968. Personal communication.
- Lewis, C. F. 1954. The inheritance of cup leaf in cotton. *J. Hered.* 45:127-128.
- McLendon, C. A. 1912. Mendelian inheritance in cotton hybrids. *Ga. Exp. Sta. Bul.* 89:141-228.
- McMichael, S. C. 1942. Occurrence of the dwarf-red character in Upland cotton. *J. Agr. Res.* 64:477-481.
- . 1954. Glandless boll in Upland cotton and its use in the study of natural crossing. *Agron. J.* 46:527-528.
- Phillips, L. L. 1962. Segregation in new allopolyploids of *Gossypium*. IV. Segregation in New World \times Asiatic and New World \times Wild American hexaploids. *Amer. J. Bot.* 49:51-57.
- Poisson, C. 1968. Preliminary report on a monosome of *Gossypium hirsutum* corresponding to linkage group I. *Coton Fib. Trop.* (English Ed.) 23:183-186.
- Rhyne, C. L. 1955. The inheritance of yellow-green, a possible mutation in cotton. *Genetics* 40:235-245.
- . 1957. Duplicate linkage groups in cotton. *J. Hered.* 48:59-62.
- . 1962. Inheritance of the glandless-leaf phenotype of Upland cotton. *J. Hered.* 53:115-123.

47. ———. 1965. Duplicate linkage blocks in glandless-leaf cotton. *J. Hered.* 56:247-252.
48. ———. 1965. Diploidization in *Gossypium hirsutum* as indicated by glandless-stem and boll inheritance. *Amer. Nat.* 96:265-276.
49. ———. 1965. Inheritance of extra-floral nectaries in cotton. *Adv. Front. Plt. Sci.* 13:121-137.
50. Richmond, T. R. 1943. Inheritance of green and brown lint in Upland cotton. *J. Amer. Soc. Agron.* 35:967-975.
51. ——— and R. E. Harper. 1937. Inheritance of okra-leaf and round-leaf in Upland cotton. A note on Brown's and Cotton's data. *J. Hered.* 28:215-216.
52. ——— and R. J. Kohel. 1961. Analysis of a completely male-sterile character in American Upland cotton. *Crop Sci.* 1:397-401.
53. Shoemaker, D. N. 1908. A study of leaf characters in cotton hybrids. *Rep. Amer. Breed. Assoc.* 5:116-119.
54. Silow, R. A. 1946. Evidence on chromosome homology and gene homology in the amphidiploid New World cottons. *J. Genet.* 47:219-221.
55. Simpson, D. M. 1947. Fuzzy leaf in cotton and its association with short lint. *J. Hered.* 48:153-156.
56. Stephens, S. C. 1954. Interspecific homologies between gene loci in *Gossypium*. I. Pollen color. *Genetics* 39:701-711.
57. ———. 1954. Interspecific homologies between gene loci in *Gossypium*. II. Corolla color. *Genetics* 39:712-723.
58. ———. 1955. Linkage in Upland cotton. *Genetics* 40:903-917.
59. Thadani, K. I. 1923. Linkage relations in the cotton plant. *Agr. J. India* 18:572-579.
60. Ware, J. O. 1929. Inheritance of lint percentage in cotton. *J. Amer. Soc. Agron.* 21:876-894.
61. ———. 1932. Inheritance of lint colors in Upland cotton. *J. Amer. Soc. Agron.* 24:550-562.
62. White, T. G. and J. E. Endrizzi. 1965. Tests for the association of marker loci with chromosomes in *Gossypium hirsutum* by the use of aneuploids. *Genetics* 51:605-612.
63. Wilson, F. D. and R. J. Kohel. 1970. Linkage of Green-Lint and Okra-Leaf genes in a reciprocal translocation stock of Upland cotton. *Can. J. Genet. Cytol.* 12:100-104.