

A Partially Male-Sterile Character in Upland Cotton¹

Norman Justus, James R. Meyer, and Jean B. Roux²

THE first heritable male-sterile line of upland cotton was reported by Justus and Leinweber (1). It was a partial male-sterile, controlled by a single recessive gene, *ms*-1. Richmond and Kohel (5) reported the development of a completely male-sterile line, *ms*₂, also controlled by a single recessive gene. Meyer and Meyer (4) have reported cytoplasmic-genetic sterility developed from interspecific crosses in cotton. The purpose of this paper is to report results of a study of the inheritance of a second partially male-sterile character in upland cotton.

MATERIALS AND METHODS

The original partially male-sterile plant was found by the junior author in a D₂ smooth progeny that had M8 (M8948) as a recurrent parent. Meyer (2) described the development of D₂ smoothness. The partially male-sterile plant was morphologically normal except for flowers that lacked pollen or had a reduced amount.

¹Contribution from the Delta Branch of the Mississippi Agricultural Experiment Station, Stoneville, Miss., in cooperation with the Crops Research Division, ARS, USDA. Journal Paper 1102, of the Mississippi Agr. Exp. Sta. Supported in part by funds provided by the Regional Cotton Genetics Project S-1. Received Apr. 4, 1963.

²Research Agronomist and Geneticist, Crops Research Division, ARS, USDA, and Delta Branch of the Mississippi Agr. Exp. Sta., and Geneticist with I.R.C.T., Paris, France.

The partial male-sterile plant or plants from its self-pollinated (S₁) progeny were crossed to 5 normal lines: M8, Dixie Triumph, Acala W 29-6, Delfos 9169, and Cook 307-6. M8 is a doubled haploid from Deltapine 14 (3) and the other four lines were from the Regional Collection of Upland Cotton. The resulting F₁ progenies were self-pollinated and backcrossed to the partially male-sterile line. Each plant in the segregating generations was classified male-sterile or male-fertile based on the scale described by Justus and Leinweber (1) where 0 = no anthers dehiscid, 1 = about 25% of anthers dehiscid, 2 = about 50% of anthers dehiscid, 3 = about 75% of anthers dehiscid, and 4 = all anthers dehiscid. Flowers from each plant were scored several times in 1958, a few times in 1959, and 5 times in 1960. A plant consistently scoring 4 was considered male-fertile, and plants scoring less than 4 were classed as partially male-sterile. Selected plants from the 1959 planting were moved to the greenhouse and self-pollinated seeds were obtained from them. The parent plants were again moved to the field and they and their progenies were scored for male-fertility and male-sterility.

RESULTS AND DISCUSSION

Data presented in Tables 1 and 2 indicate that the partial male-sterility in these populations is controlled by a single recessive gene. The symbol *ms*-3 is proposed for this gene. The 1958 backcross (BC₁) and F₂ progenies involving partial male-sterile and M8 indicated a good fit to the expected ratios. The segregating progenies of the partial male-sterile crossed with the remaining 4 male-fertile lines

Population	Number of plants		χ^2	P
	Male fertile	Male sterile		
BC₁:				
(ms-3 ms-3 × M8)-1 × (ms-3 ms-3)-1	4	7	0.8182	.50 - .25*
(ms-3 ms-3 × M8)-1 × (ms-3 ms-3)-3	8	3	2.2737	.25 - .10
(ms-3 ms-3 × M8)-1 × (ms-3 ms-3)-4	6	4	0.4000	.050 - .025
(ms-3 ms-3 × M8)-3 × (ms-3 ms-3)-3	3	2	0.2000	.25 - .10
(ms-3 ms-3 × M8)-7 × (ms-3 ms-3)-3	3	3	0.0000	1
(ms-3 ms-3 × M8)-7 × (ms-3 ms-3)-1	8	2	3.6000	.10 - .05
Total	32	21	7.2919	
Deviation			2.2830	.25 - .10
Heterogeneity			5.0079	.50 - .25
F₂:				
(ms-3 ms-3 × M8)-1-1	68	18	0.760	.50 - .25†
(ms-3 ms-3 × M8)-2-1	28	6	0.980	.50 - .25
(ms-3 ms-3 × M8)-4-1	19	7	0.051	.90 - .75
Total	115	31	1.791	
Deviation			1.105	.50 - .25
Heterogeneity			0.686	.80 - .70

	Population	Number of plants		χ^2	P
		Male- fertile	Male- sterile		
BC₁:					
ms-3 ms-3 \times (ms-3 ms-3 \times Dixie Triumph)	4	2	0.667	.50-.25*	
(ms-3 ms-3 \times Dixie Triumph) \times ms-3 ms-3	3	10	3.769	.10-.05	
(Dixie Triumph \times ms-3 ms-3) \times ms-3 ms-3	6	5	0.091	.90-.75	
(Acala W 29-6 \times ms-3 ms-3) \times ms-3 ms-3	9	16	1.960	.25-.10	
(ms-3 ms-3 \times Delfos 9169) \times ms-3 ms-3	4	1	1.800	.25-.10	
(ms-3 ms-3 \times Cook 307-6) \times ms-3 ms-3	4	8	1.333	.25-.10	
Total	30	42	9.619		
Deviation			2.000	.25-.10	
Heterogeneity			7.619	.20-.10	
F₂:					
ms-3 ms-3 \times Dixie Triumph	4	7	8.758	.01†	
Acala W 29-6 \times ms-3 ms-3	7	6	3.103	.10-.05	
ms-3 ms-3 \times Delfos 9169	10	1	1.485	.30-.20	
ms-3 ms-3 \times Cook 307-6	5	2	0.048	.90-.80	
Total	26	16	13.392		
Deviation			3.841	.05	
Heterogeneity			9.551	.05-.32	

No.	Parent plant*				Progeny test		
	1959 mean†	1960 array	Assumed genotype	Genera- tion	Observed‡		P
					F	S	
1	0	0-1-1-2-3	ms-3 ms-3	F ₂	0	11	-
2	1	1-1-0-0-1	ms-3 ms-3	F ₂	0	10	-
3	1	2-0-2-1-2	ms-3 ms-3	F ₂	0	9	-
4	2	4-4-4-4-4	Ms-3 ms-3	F ₂	17	8	.653
5	2	0-1-1-1-4	ms-3 ms-3	F ₂	0	11	.50-30
6	2	0-1-1-2-0	ms-3 ms-3	F ₂	0	14	-
7	3	1-1-3-0-0	ms-3 ms-3	F ₂	0	13	-
8	3	4-4-4-4-4	Ms-3 ms-3	F ₂	9	3	.000
9	4	4-4-4-4-4	Ms-3 ms-3	F ₂	16	0	-
10	4	4-4-4-4-4	Ms-3 ms-3	F ₂	12	2	.857
11	1	3-1-1-2-4	ms-3 ms-3	F ₂	0	13	.50-25
12	3	2-0-1-1-2	ms-3 ms-3	F ₂	0	14	-
13	4	3-1-0-2-1	ms-3 ms-3	F ₂	0	13	-

The flowers of the advanced-generation steriles resembled the original male-steriles in that they lack pollen or have



Seed of the *ms-3 ms-3* genotype has been placed in the Regional Collection of Upland Cotton.

A second recessive gene for partial male-sterility of upland cotton is reported. The percent of sterile anthers for homozygous plants varies in response to environment. The symbol *ms-3* is proposed for this gene.

1. JUSTUS, NORMAN, and LEINWEBER, C. L. A heritable partially male-sterile character in cotton. *J. of Hered.* 51:191-192. 1960.
2. MEYER, JAMES R. Origin and inheritance of D₂ smoothness in upland cotton. *J. Hered.* 48:249-250. 1957.
3. ———, and JUSTUS, NORMAN. Properties of doubled haploids of cotton. *Crop Sci.* 1:462-464. 1961.
4. ———, and MEYER, VESTA G. Cytoplasmic male-sterility in cotton. *Genetics* 46(8):883. 1961.
5. RICHMOND, T. R., and KOHEL, R. J. Analysis of a completely male-sterile character in American Upland cotton. *Crop Sci.* 1:397-401. 1961.
6. SMITH, LUTHER. Possible practical method for producing hybrid seed of self-pollinated crops through the use of male-sterility. *J. Am. Soc. Agron.* 39:260-261. 1947.