

HN-02 · VITAMINS & MINERALS · HUMAN NUTRITION

HS-VitaA

Retinol / Beta-Carotene

INCI	CAS	EC	FAMILY	SUPPLIED AS
Retinol / Beta-Carotene	68-26-8 +1	—	HN-02	Per vitamin

01 POSITIONING

Fortification is arithmetic — it only works when the assay on the drum matches the number on the COA.

02 HOW IT WORKS

- 01 Defined nutrient function
Each vitamin covers an established metabolic role — immune, vision, bone, antioxidant.
- 02 Assay-driven dosing
Fortification math depends on a tight, reproducible assay per lot.
- 03 Stabilized formats
Coated and CWS grades protect fragile vitamins through processing and shelf.

03 APPLICATIONS

- Multivitamin tablets and capsules
- Effervescent and powder drinks
- Food and beverage fortification
- Gummies (grade-dependent)
- Infant and clinical nutrition

04 FORMULATION GUIDE

USE LEVEL	Per RDA and matrix
PH & CONDITIONS	Vitamin-dependent; protect C and A from oxidation and light

- Overage the fragile vitamins (A, C) per stability data to hit label claim at end of shelf-life.
- Choose coated/CWS grades for effervescent and moisture-exposed matrices.

05 TYPICAL SPECIFICATION

Appearance	per vitamin: powder / oil / beadlet
Assay	per pharmacopoeia (USP/EP)
Grade	USP / FCC / feed per TDS

Typical parameters. The specification on the current TDS and the per-lot Certificate of Analysis govern.

06 PERFORMANCE — DIRECTIONAL

Assay consistency lot-to-lot
Pharmacopoeia grade

Stability of coated vs plain
Coated / CWS grade

Values are representative of the ingredient class, based on published technical literature, and shown directionally for comparison. Final performance must be validated in the customer's formulation.