

PC-02 · SILICONES · PERSONAL CARE

HS 949

Amodimethicone & Cetrimonium Chloride & Trideceth-12

INCI	CAS	EC	FAMILY	SUPPLIED AS
Amodimethicone & Cetrimonium Chloride & Trideceth-12	—	—	PC-02	Emulsion system

01 POSITIONING

Damage-targeted chemistry — the claim 'repairs where it's needed' with a mechanism behind it.

02 HOW IT WORKS

- 01

Charge-directed deposition

Protonated amine groups bind the negatively charged damaged cuticle first.
- 02

Durable film

Resists several wash cycles — conditioning with memory.
- 03

Thermal protection support

The film evens heat distribution from dryers and irons.

03 APPLICATIONS

- Masks and deep conditioners
- Post-chemical treatment lines
- Leave-in repair
- Textile softening (premium)

04 FORMULATION GUIDE


USE LEVEL	0.5–4%
PH & CONDITIONS	Formulate pH 3.5–6 to keep amine protonated
— Post-add below 40 °C; excess dosage can build up on virgin hair — respect the range.	
— Combine with cationic guar for a two-layer conditioning system.	

05 TYPICAL SPECIFICATION

Appearance	Milky emulsion / clear microemulsion
Solids	per TDS
Amine value	per grade

Typical parameters. The specification on the current TDS and the per-lot Certificate of Analysis govern.

06 PERFORMANCE — DIRECTIONAL



Values are representative of the ingredient class, based on published technical literature, and shown directionally for comparison. Final performance must be validated in the customer's formulation.