

Supplementary Material for Bayesian Network Problem in Project 1

Jun Chu

September 22, 2016

1 Linear Gaussian Model

Consider a Bayesian network where continuous random variable Y has k parents $\mathbf{X} = \{X_1, \dots, X_k\}$. Y is said to obey a linear Gaussian model with parameters β_1, \dots, β_k and σ^2 if $P(Y|X_1, \dots, X_k) \sim \mathcal{N}(\beta_0 + \beta_1 X_1 + \dots + \beta_k X_k; \sigma^2)$. Let's define $\boldsymbol{\theta} = (\beta_1, \dots, \beta_k, \sigma^2)$. The log-likelihood is Linear Gaussian Model

$$\begin{aligned} L(\boldsymbol{\theta}) &= \log(P(Y|X_1, \dots, X_k)) \\ &= \sum_{n=1}^N \left[-\frac{1}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} (\beta_0 x_0[n] + \beta_1 x_1[n] + \dots + \beta_k x_k[n] - y[n])^2 \right]. \end{aligned}$$

Here, we assume $x_0[n] \equiv 1$. It is possible to get a closed-form solution. It involves solving a set of equations which are obtained by taking partial derivatives of the log-likelihood function, as follows. Gradient of log-likelihood with respect to β_i is

$$\begin{aligned} \frac{\partial L(\boldsymbol{\theta})}{\partial \beta_i} &= -\frac{1}{\sigma^2} \sum_{n=1}^N \{(\beta_0 x_0[n] + \beta_1 x_1[n] + \dots + \beta_k x_k[n] - y[n]) x_i[n]\} \\ &= -\frac{1}{\sigma^2} \left(\beta_0 \sum_{n=1}^N x_0[n] x_i[n] + \beta_1 \sum_{n=1}^N x_1[n] x_i[n] + \dots + \beta_k \sum_{n=1}^N x_k[n] x_i[n] - \sum_{n=1}^N y[n] x_i[n] \right). \end{aligned}$$

Equating to zero and rearranging we get

$$\beta_0 \sum_{n=1}^N x_0[n] x_i[n] + \beta_1 \sum_{n=1}^N x_1[n] x_i[n] + \dots + \beta_k \sum_{n=1}^N x_k[n] x_i[n] = \sum_{n=1}^N y[n] x_i[n], \quad i = 0, 1, \dots, k.$$

If we define:

$$A = \begin{pmatrix} \sum_{n=1}^N x_0[n] x_0[n] & \sum_{n=1}^N x_1[n] x_0[n] & \dots & \sum_{n=1}^N x_k[n] x_0[n] \\ \sum_{n=1}^N x_0[n] x_1[n] & \sum_{n=1}^N x_1[n] x_1[n] & \dots & \sum_{n=1}^N x_k[n] x_1[n] \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{n=1}^N x_0[n] x_k[n] & \sum_{n=1}^N x_1[n] x_k[n] & \dots & \sum_{n=1}^N x_k[n] x_k[n] \end{pmatrix},$$

$$\boldsymbol{\beta} = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_k \end{pmatrix}$$

and

$$\mathbf{y} = \begin{pmatrix} \sum_{n=1}^N y[n]x_0[n] \\ \sum_{n=1}^N y[n]x_1[n] \\ \vdots \\ \sum_{n=1}^N y[n]x_k[n] \end{pmatrix}.$$

Therefore $A\boldsymbol{\beta} = \mathbf{y}$. $\boldsymbol{\beta}$ could be solved using $\boldsymbol{\beta} = A^{-1}\mathbf{y}$.

Gradient of log-likelihood with respect to σ is

$$\frac{\partial L(\boldsymbol{\theta})}{\partial \sigma} = -\frac{N}{\sigma} + \frac{1}{\sigma^3} \sum_{n=1}^N (\beta_0 x_0[n] + \beta_1 x_1[n] + \dots + \beta_k x_k[n] - y[n])^2.$$

Set it to zero, we get

$$\sigma^2 = \frac{1}{N} \sum_{n=1}^N (\beta_0 x_0[n] + \beta_1 x_1[n] + \dots + \beta_k x_k[n] - y[n])^2.$$

x

2 Bayesian Network Factorization

Given a Bayesian network G of d variables $\mathbf{X} = \{X_1, X_2, \dots, X_d\}$, the joint probability distribution is given by

$$p(\mathbf{X}) = \prod_{i=1}^d p(X_i | \text{pa}(X_i)),$$

where $\text{pa}(x_i)$ are the parent variables of x_i . The log-likelihood is

$$\log p(\mathbf{X}) = \sum_{i=1}^d \log p(X_i | \text{pa}(X_i)).$$

Each term could be maximized separately using the aforementioned linear gaussian model closed form solution.