

Chapter 1

Exercise 1.1

1. (i) coefficient of $x^2 = 3$
(ii) coefficient of $x = -9$
(iii) independent term = 5.
2. (i) degree 2
(ii) degree 3
(iii) degree 4.
3. $\frac{-4}{x} = -4x^{-1}$, -1 is not a positive power;
 $x^{\frac{3}{2}}$, $\frac{3}{2}$ is not an integer.
4. (i) $3x^2 - 6x + 7 + 5x^2 + 2x - 9 = 8x^2 - 4x - 2$
(ii) $x^3 - 4x^2 - 5x + 3x^3 + 6x^2 - x = 4x^3 + 2x^2 - 6x$
(iii) $x(x + 4) + 3x(2x - 3) = x^2 + 4x + 6x^2 - 9x$
 $= 7x^2 - 5x$
(iv) $3(x^2 - 7) + 2x(3x - 1) - 7x + 2 = 3x^2 - 21 + 6x^2 - 2x - 7x + 2$
 $= 9x^2 - 9x - 19$
5. (i) $3x^2(4x + 2) + 5x^2(2x - 5) = 12x^3 + 6x^2 + 10x^3 - 25x^2$
 $= 22x^3 - 19x^2$
(ii) $x^3(x - 2) + 4x^3(2x - 6) = x^4 - 2x^3 + 8x^4 - 24x^3$
 $= 9x^4 - 26x^3$
(iii) $x(x^3 + 4x^2 - 7x) + 3x^2(2x^2 - 3x + 4) = x^4 + 4x^3 - 7x^2 + 6x^4 - 9x^3 + 12x^2$
 $= 7x^4 - 5x^3 + 5x^2$
(iv) $3x(x^2 - 7x + 1) + 2x^2(6x - 5) = 3x^3 - 21x^2 + 3x + 12x^3 - 10x^2$
 $= 15x^3 - 31x^2 + 3x$
6. (i) $(x + 4)(2x + 5) = 2x^2 + 5x + 8x + 20 = 2x^2 + 13x + 20$
(ii) $(2x + 3)(x - 2) = 2x^2 - 4x + 3x - 6 = 2x^2 - x - 6$
(iii) $(3x - 2)(x + 3) = 3x^2 + 9x - 2x - 6 = 3x^2 + 7x - 6$
(iv) $(3x - 2)(4x - 1) = 12x^2 - 3x - 8x + 2 = 12x^2 - 11x + 2$
(v) $(3x - 1)(2x + 5) = 6x^2 + 15x - 2x - 5 = 6x^2 + 13x - 5$
(vi) $(4x + 1)(2x - 6) = 8x^2 - 24x + 2x - 6 = 8x^2 - 22x - 6$
(vii) $(x - 2)(x + 2) = x^2 + 2x - 2x - 4 = x^2 - 4$
(viii) $(2x + 5)(2x - 5) = 4x^2 - 10x + 10x - 25 = 4x^2 - 25$
(ix) $(ax - by)(ax + by) = a^2x^2 + abxy - abxy - b^2y^2 = a^2x^2 - b^2y^2$
7. (i) $(x + 2)^2 = x^2 + 4x + 4$
(ii) $(x - 3)^2 = x^2 - 6x + 9$
(iii) $(x + 5)^2 = x^2 + 10x + 25$
(iv) $(a + b)^2 = a^2 + 2ab + b^2$
(v) $(x - y)^2 = x^2 - 2xy + y^2$
(vi) $(a + 2b)^2 = a^2 + 4ab + 4b^2$
(vii) $(3x - y)^2 = 9x^2 - 6xy + y^2$
(viii) $(x - 5y)^2 = x^2 - 10xy + 25y^2$
(ix) $(2x + 3y)^2 = 4x^2 + 12xy + 9y^2$

$$8. \quad (i) \left(x + \frac{1}{2}\right)^2 = x^2 + 2(x)\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 = x^2 + x + \frac{1}{4}$$

$$(ii) \quad 8\left(x - \frac{1}{4}\right)^2 = 8\left(x^2 - 2(x)\left(\frac{1}{4}\right) + \left(\frac{-1}{4}\right)^2\right) = 8\left(x^2 - \frac{x}{2} + \frac{1}{16}\right) \\ = 8x^2 - 4x + \frac{1}{2}$$

$$(iii) \quad -(1-x)^2 = -(1-2x+x^2) = -1+2x-x^2$$

$$9. \quad (i) \quad x^2 + 5x + 25; \text{ No, cannot be written in the form } (x+k)^2$$

$$(ii) \quad 9x^2 - 6x - 1; \text{ No, cannot be written in the form } (x+k)^2$$

$$(iii) \quad 4 + 12x + 9x^2 = (2 + 3x)^2; \text{ YES.}$$

$$10. \quad px^2 + 4x + 1 = (ax + 1)^2$$

$$px^2 + 4x + 1 \text{ can be written in the form } (ax + 1)^2 = a^2x^2 + 2ax + 1$$

$$\therefore 2a = 4 \Rightarrow a = 2$$

$$\therefore (ax + 1)^2 = (2x + 1)^2 = 4x^2 + 4x + 1$$

$$\therefore p = 4.$$

$$11. \quad 25x^2 + tx + 4 = (5x + 2)^2 \quad \text{or} \quad 25x^2 + tx + 4 = (5x - 2)^2 \\ = 25x^2 + 20x + 4 \quad \quad \quad = 25x^2 - 20x + 4$$

$$\Rightarrow \quad t = 20 \quad \quad \quad \Rightarrow \quad t = -20$$

$$12. \quad 9x^2 + 24x + s = (3x + a)^2 = 9x^2 + 6ax + a^2$$

$$\Rightarrow 6a = 24$$

$$a = 4.$$

$$\therefore 9x^2 + 24x + s = (3x + 4)^2 \\ = 9x^2 + 24x + 16$$

$$\Rightarrow s = 16.$$

$$13. \quad (i) \quad (x + 2)(x^2 + 2x + 6) = x^3 + 2x^2 + 6x + 2x^2 + 4x + 12 \\ = x^3 + 4x^2 + 10x + 12$$

$$(ii) \quad (x - 4)(2x^2 + 3x - 1) = 2x^3 + 3x^2 - x - 8x^2 - 12x + 4 \\ = 2x^3 - 5x^2 - 13x + 4$$

$$(iii) \quad (2x + 3)(x^2 - 3x + 2) = 2x^3 - 6x^2 + 4x + 3x^2 - 9x + 6 \\ = 2x^3 - 3x^2 - 5x + 6$$

$$(iv) \quad (3x - 2)(2x^2 - 4x + 2) = 6x^3 - 12x^2 + 6x - 4x^2 + 8x - 4 \\ = 6x^3 - 16x^2 + 14x - 4.$$

$$14. \quad (x + y)(x^2 - xy + y^2) = x^3 - \cancel{x^2y} + \cancel{xy^2} + \cancel{x^2y} - \cancel{xy^2} + y^3 \\ = x^3 + y^3$$

$$15. \quad (x - y)(x^2 + xy + y^2) = x^3 + \cancel{x^2y} + \cancel{xy^2} - \cancel{x^2y} - \cancel{xy^2} - y^3 \\ = x^3 - y^3$$

$$16. \quad (2x - 3)(3x^2 - 2x + 4) = 6x^3 - 4x^2 + 8x - 9x^2 + 6x - 12 \\ = 6x^3 - 13x^2 + 14x - 12 \\ \Rightarrow \text{coefficient of } x = 14.$$

$$17. \quad (x + 3)(x - 4)(2x + 1) = (x + 3)(2x^2 + x - 8x - 4) \\ = (x + 3)(2x^2 - 7x - 4) \\ = 2x^3 - 7x^2 - 4x + 6x^2 - 21x - 12 \\ = 2x^3 - x^2 - 25x - 12.$$

$$18. \quad (x^2 - 3x - 2)(2x^2 - 4x + 1) = 2x^4 - 4x^3 + x^2 - 6x^3 + 12x^2 - 3x \\ - 4x^2 + 8x - 2 \\ = 2x^4 - 10x^3 + 9x^2 + 5x - 2.$$

19. $(3x^2 + 5x - 1)(2x^2 - 6x - 5)$; x^2 coefficients include $-15x^2 - 30x^2 - 2x^2$
 $= -47x^2$
 coefficient of $x^2 = -47$.

20. (i) $\frac{3x+6}{3} = x+2$

(ii) $\frac{x^2+2x}{x} = x+2$

(iii) $\frac{3x^3-6x^2}{3x} = \frac{3x^2(x-2)}{3x} = x(x-2) = x^2-2x$

(iv) $\frac{15x^2y-10xy^2}{5xy} = 3x-2y$

21. (i) $\frac{6x^2y+9xy^2-3xy}{3xy} = x+3y-1$

(ii) $\frac{6x^4-9x^3+12x^2}{3x^2} = \frac{3x^2(x^2-3x+4)}{3x^2}$
 $= x^2-3x+4$

22. (i) $\frac{\cancel{1}^4\cancel{2}^4a^2b}{\cancel{3}^4\cancel{a}^4b} = 4a$

(ii) $\frac{\cancel{1}^4\cancel{2}^4a^2b\cancel{c}^4}{\cancel{3}^4\cancel{a}^4\cancel{c}^4} = 4ab$

(iii) $\frac{\cancel{4}^2xy^2z}{\cancel{2}^2xy} = 2yz$

(iv) $\frac{3xy}{2} \cdot \frac{4}{6x^2} = \frac{\cancel{1}^2\cancel{2}^2xy}{\cancel{1}^2\cancel{2}^2x^2} = \frac{y}{x}$

23. (i) $\frac{2x^2+5x-3}{2x-1} = \frac{\cancel{(2x-1)}^{\cancel{2x-1}}(x+3)}{\cancel{2x-1}} = x+3$

(ii) $\frac{2x^2-2x-12}{x-3} = \frac{2(x^2-x-6)}{x-3} = \frac{2\cancel{(x-3)}^{\cancel{x-3}}(x+2)}{\cancel{x-3}} = 2(x+2) = 2x+4$

(iii) $\frac{8x^2+8x-6}{4x-2} = \frac{2(4x^2+4x-3)}{2(2x-1)} = \frac{\cancel{2}^{\cancel{2}}\cancel{(2x-1)}^{\cancel{2x-1}}(2x+3)}{\cancel{2}^{\cancel{2}}\cancel{(2x-1)}} = 2x+3$

24. (i)
$$\begin{array}{r} x^2 - 7x + 12 \\ x-1 \overline{) x^3 - 8x^2 + 19x - 12} \\ \underline{x^3 - x^2} \\ -7x^2 + 19x - 12 \\ \underline{-7x^2 + 7x} \\ 12x - 12 \\ \underline{12x - 12} \\ 0 \end{array}$$

(ii)
$$\begin{array}{r} x^2 - 1 \\ 2x-1 \overline{) 2x^3 - x^2 - 2x + 1} \\ \underline{2x^3 - x^2} \\ -2x + 1 \\ \underline{-2x + 1} \\ 0 \end{array}$$

(iii)
$$\begin{array}{r} x^2 - 1 \\ 3x-4 \overline{) 3x^3 - 4x^2 - 3x + 4} \\ \underline{3x^3 - 4x^2} \\ -3x + 4 \\ \underline{-3x + 4} \\ 0 \end{array}$$

$$\begin{array}{r}
 \text{(iv)} \quad 4x^2 + 5x - 6 \\
 x - 3 \overline{) 4x^3 - 7x^2 - 21x + 18} \\
 \underline{4x^3 - 12x^2} \\
 + 5x^2 - 21x + 18 \\
 \underline{+ 5x^2 - 15x} \\
 - 6x + 18 \\
 \underline{- 6x + 18} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(v)} \quad x^2 - 5x + 3 \\
 x + 5 \overline{) x^3 - 22x + 15} \\
 \underline{x^3 + 5x^2} \\
 - 5x^2 - 22x + 15 \\
 \underline{- 5x^2 - 25x} \\
 + 3x + 15 \\
 \underline{+ 3x + 15} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(vi)} \quad 2x^2 + 3x + 6 \\
 x - 2 \overline{) 2x^3 - x^2 - 12} \\
 \underline{2x^3 - 4x^2} - 12 \\
 3x^2 - 12 \\
 \underline{3x^2 - 6x} \\
 6x - 12 \\
 \underline{6x - 12} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{25. (i)} \quad x - 2 \\
 x^2 + 2 \overline{) x^3 - 2x^2 + 2x - 4} \\
 \underline{x^3 + 2x} \\
 - 2x^2 - 4 \\
 \underline{- 2x^2 - 4} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(ii)} \quad x - 3 \\
 x^2 - 6x + 9 \overline{) x^3 - 9x^2 + 27x - 27} \\
 \underline{x^3 - 6x^2 + 9x} \\
 - 3x^2 + 18x - 27 \\
 \underline{- 3x^2 + 18x - 27} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(iii)} \quad 3x - 1 \\
 x^2 + x - 2 \overline{) 3x^3 + 2x^2 - 7x + 2} \\
 \underline{3x^3 + 3x^2 - 6x} \\
 - x^2 - x + 2 \\
 \underline{- x^2 - x + 2} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(iv)} \quad x + 2 \\
 5x^2 + 4x - 1 \overline{) 5x^3 + 14x^2 + 7x - 2} \\
 \underline{5x^3 + 4x^2 - x} \\
 + 10x^2 + 8x - 2 \\
 \underline{+ 10x^2 + 8x - 2} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{26. (i)} \quad x^2 + 2x + 4 \\
 x - 2 \overline{) x^3 - 8} \\
 \underline{x^3 - 2x^2} \\
 2x^2 - 8 \\
 \underline{2x^2 - 4x} \\
 4x - 8 \\
 \underline{4x - 8} \\
 0
 \end{array}$$

$$\begin{array}{r}
 \text{(ii)} \quad \begin{array}{r} 4x^2 + 6xy + 9y^2 \\ 2x - 3y \overline{) 8x^3 - 27y^3} \\ \underline{8x^3 - 12x^2y} \\ 12x^2y \\ \underline{12x^2y - 18xy^2} \\ 18xy^2 - 27y^3 \\ \underline{18xy^2 - 27y^3} \end{array}
 \end{array}$$

Exercise 1.2

1. x cm length of smaller side,

$(x + 4)$ cm length of longer side.

(i) $A(x) = x(x + 4) = (x^2 + 4x) \text{ cm}^2$

(ii) $P(x) = 2[x + (x + 4)] = (4x + 8) \text{ cm}$

2. (i) Area = length \times width

$$\begin{aligned}
 \Rightarrow \text{width} &= \frac{\text{Area}}{\text{length}} = \frac{6x^2 + 4x - 2}{3x - 1} \\
 &= \frac{(3x - 1)(2x + 2)}{3x - 1} = 2x + 2
 \end{aligned}$$

(ii) Perimeter = $2(\text{length} + \text{width})$

$$= 2((3x - 1) + (2x + 2))$$

$$P(x) = 10x + 2$$

3. (a) $V(x) = (2x + 3)(x)(x + 1)$

$$= (2x + 3)(x^2 + x)$$

$$= 2x^3 + 2x^2 + 3x^2 + 3x$$

$$= 2x^3 + 5x^2 + 3x$$

(b) $S(x) = (x)(2x + 3) + 2(x)(x + 1) + 2(2x + 3)(x + 1)$

$$= 2x^2 + 3x + 2x^2 + 2x + 4x^2 + 4x + 6x + 6$$

$$= 8x^2 + 15x + 6$$

(c) (i) $V(5) = 2(5)^3 + 5(5)^2 + 3(5) = 390 \text{ cm}^3$

(ii) $S(5) = 8(5)^2 + 15(5) + 6 = 281 \text{ cm}^2$

4. $f(x) = 2x^3 - x^2 - 5x - 4$

(a) $f(0) = 2(0)^3 - (0)^2 - 5(0) - 4 = -4$

(b) $f(1) = 2(1)^3 - (1)^2 - 5(1) - 4 = -8$

(c) $f(-2) = 2(-2)^3 - (-2)^2 - 5(-2) - 4 = -14$

(d) $f(3a) = 2(3a)^3 - (3a)^2 - 5(3a) - 4 = 54a^3 - 9a^2 - 15a - 4$

5. $f(x) = x^2 - 3x + 6$

(a) $f(0) = (0)^2 - 3(0) + 6 = 6$

(b) $f(-5) = (-5)^2 - 3(-5) + 6 = 46$

(c) $f\left(-\frac{1}{2}\right) = \left(-\frac{1}{2}\right)^2 - 3\left(-\frac{1}{2}\right) + 6 = \frac{31}{4} = 7.75$

(d) $f\left(\frac{a}{4}\right) = \left(\frac{a}{4}\right)^2 - 3\left(\frac{a}{4}\right) + 6 = \frac{a^2}{16} - \frac{3a}{4} + 6$

6. Length = $(x - y)$.

Width = $(2x + 3y)$.

(a) Area = $(x - y)(2x + 3y) = 2x^2 + 3xy - 2xy - 3y^2$
 $= 2x^2 + xy - 3y^2$

(b) Perimeter = $2[(x - y) + (2x + 3y)] = 2[3x + 2y] = 6x + 4y$

7. Length = x cm

Width = $(x - 5)$ cm

Height = $2x$ cm.

$$\begin{aligned} \text{(a) Volume} &= \text{Length} \times \text{Width} \times \text{Height} \\ &= (x)(x - 5)(2x) = 2x^3 - 10x^2 \end{aligned}$$

$$\begin{aligned} \text{(b) Surface area} &= 2(x)(x - 5) + 4(2x)(x - 5) + 4(2x)(x) \\ &= 2x^2 - 10x + 8x^2 - 40x + 8x^2 \\ &= 18x^2 - 50x. \end{aligned}$$

8. (i) $d(4)$ = the number of diagonals in a 4-sided polygon


(ii) $d(5)$ = the number of diagonals in a 5-sided polygon.

$$d(4) = \frac{(4)^2}{2} - \frac{3(4)}{2} = 8 - 6 = 2.$$

$$d(5) = \frac{(5)^2}{2} - \frac{3(5)}{2} = \frac{25}{2} - \frac{15}{2} = \frac{10}{2} = 5.$$

$$d(6) = \frac{(6)^2}{2} - \frac{3(6)}{2} = \frac{36}{2} - \frac{18}{2} = \frac{18}{2} = 9.$$

$d(3) = 0$ because a triangle has no diagonal.



9. If $f(x) = x + 5$,

$$\begin{aligned} f(a^2) - 3f(a) + 2 &= a^2 + 5 - 3(a + 5) + 2 \\ &= a^2 + 5 - 3a - 15 + 2 \\ &= a^2 - 3a - 8. \end{aligned}$$

10. $f(x) = x^2 - 3x + 6$.

$$\text{(i) } f(-2t) = (-2t)^2 - 3(-2t) + 6 = 4t^2 + 6t + 6.$$

$$\text{(ii) } f(t^2) = (t^2)^2 - 3(t^2) + 6 = t^4 - 3t^2 + 6.$$

$$\begin{aligned} \text{(iii) } f(t - 2) &= (t - 2)^2 - 3(t - 2) + 6 = t^2 - 4t + 4 - 3t + 6 + 6 \\ &= t^2 - 7t + 16. \end{aligned}$$

(i) $4t^2 + 6t + 6$ is of degree 2.

(ii) $t^4 - 3t^2 + 6$ is of degree 4.

(iii) $t^2 - 7t + 16$ is of degree 2.

11. $V(r, h) = \frac{1}{3}\pi r^2 h$.

$$\text{(i) } V(r = 14, h = 21) = \frac{1}{3}\pi \cdot 14^2 \cdot 21 = 1372\pi \text{ cm}^3$$

$$\text{(ii) } V(r, h = r) = \frac{1}{3}\pi \cdot r^2 \cdot r = \frac{1}{3}\pi r^3$$

$$\text{(iii) } V(r = 2h, h) = \frac{1}{3}\pi(2h)^2 \cdot h = \frac{4}{3}\pi h^3$$

12. The dimensions of the base of the box are $24 - 2x$ and $18 - 2x$, in cm.

The height of the box is x , in cm.

Thus the volume, V , of the box, in cm^3 , is

$$V = x(24 - 2x)(18 - 2x)$$

$$V = x(432 - 48x - 36x + 4x^2)$$

$$V = x(4x^2 - 84x + 432)$$

$$V = 4x^3 - 84x^2 + 432x$$

The smallest possible value of x is 0. The largest possible value of x is given by

$$18 - 2x = 0$$

$$9 = x$$

Hence $0 < x < 9$, and so $a = 0$ and $b = 9$.

13. $T = 2\pi\sqrt{\frac{l}{g}}$

$$\therefore T^2 = 4\pi^2 \frac{l}{g}$$

$$\therefore \frac{gT^2}{4\pi^2} = l$$

$$\therefore l = \frac{gT^2}{4\pi^2}$$

$$\Rightarrow \text{when } T = 4 \text{ and } g = 10, l = \frac{10 \cdot 4^2}{4\pi^2} = \frac{40}{\pi^2} \text{ m.}$$

14. $V = \frac{4}{3}\pi r^3$

$$\Rightarrow \frac{4}{3}\pi r^3 = V$$

$$\Rightarrow r^3 = \frac{3V}{4\pi}$$

$$\Rightarrow r = \sqrt[3]{\frac{3V}{4\pi}}$$

$$\text{when } V = \frac{792}{7} \text{ and } \pi = \frac{22}{7}, r = \sqrt[3]{\frac{3 \times \frac{792}{7} \times 7}{4 \times \frac{22}{7}}} = \sqrt[3]{\frac{198}{4}} = \sqrt[3]{27} = 3 \text{ m}$$

15. $H(x) = \frac{x}{2}(x - 1)$, x = number of students.

(i) $x = 5 \Rightarrow H(5) = \frac{5}{2}(5 - 1) = 10$

(ii) $x = 6 \Rightarrow H(6) = \frac{6}{2}(6 - 1) = 15$

(iii) $x = 10 \Rightarrow H(10) = \frac{10}{2}(10 - 1) = 45$

(iv) $H(x) = 136 = \frac{x}{2}(x - 1)$

$$272 = x(x - 1)$$

\Rightarrow The product of two consecutive numbers = 272.

$$\Rightarrow \text{if } x = 16, x - 1 = 15 \quad \therefore 16 \times 15 = 240.$$

$$\text{if } x = 17, x - 1 = 16 \quad \therefore 17 \times 16 = 272.$$

$$\therefore x = 17.$$

or

$$272 = x^2 - x$$

$$\Rightarrow x^2 - x - 272 = 0$$

$$(x - 17)(x + 16) = 0$$

$$\therefore x - 17 = 0 \Rightarrow x = 17$$

or $x + 16 = 0 \Rightarrow x = -16$ which is invalid
since x stands for the number of students

$$\therefore x = 17.$$

Exercise 1.3

1. $5x^2 - 10x = 5x(x - 2)$

2. $6ab - 12bc = 6b(a - 2c)$

3. $3x^2 - 6xy = 3x(x - 2y)$

4. $2x^2y - 6x^2z = 2x^2(y - 3z)$
5. $2a^3 - 4a^2 + 8a = 2a(a^2 - 2a + 4)$
6. $5xy^2 - 20x^2y = 5xy(y - 4x)$
7. $2a^2b - 4ab^2 + 12abc = 2ab(a - 2b + 6c)$
8. $3x^2y - 9xy^2 + 15xyz = 3xy(x - 3y + 5z)$
9. $4\pi r^2 + 6\pi rh = 2\pi r(2r + 3h)$
10. $3a(2b - c) - 4(2b - c) = (2b - c)(3a - 4)$
11. $x^2 - ax + 3x - 3a = x(x - a) + 3(x - a)$
 $= (x - a)(x + 3)$
12. $2c^2 - 4cd + c - 2d = 2c(c - 2d) + c - 2d$
 $= (c - 2d)(2c + 1)$
13. $8ax + 4ay - 6bx - 3by = 4a(2x + y) - 3b(2x + y)$
 $= (2x + y)(4a - 3b)$
14. $7y^2 - 21by + 2ay - 6ab = 7y(y - 3b) + 2a(y - 3b)$
 $= (y - 3b)(7y + 2a)$
15. $6xy + 12yz - 8xz - 9y^2 = 6xy - 9y^2 + 12yz - 8xz$
 $= 3y(2x - 3y) + 4z(3y - 2x)$
 $= 3y(2x - 3y) - 4z(2x - 3y)$
 $= (2x - 3y)(3y - 4z)$
16. $6x^2 - 3y(3x - 2a) - 4ax = 6x^2 - 4ax - 3y(3x - 2a)$
 $= 2x(3x - 2a) - 3y(3x - 2a)$
 $= (3x - 2a)(2x - 3y)$
17. $3ax^2 - 3ay^2 - 4bx^2 + 4by^2 = 3a(x^2 - y^2) - 4b(x^2 - y^2)$
 $= (x^2 - y^2)(3a - 4b)$
 $= (x - y)(x + y)(3a - 4b)$
18. $a^2 - b^2 = (a - b)(a + b)$
19. $x^2 - 4y^2 = (x - 2y)(x + 2y)$
20. $9x^2 - y^2 = (3x - y)(3x + y)$
21. $16x^2 - 25y^2 = (4x)^2 - (5y)^2 = (4x - 5y)(4x + 5y)$
22. $36x^2 - 25 = (6x - 5)(6x + 5)$
23. $1 - 36x^2 = (1 - 6x)(1 + 6x)$
24. $49a^2 - 4b^2 = (7a)^2 - (2b)^2 = (7a - 2b)(7a + 2b)$
25. $x^2y^2 - 1 = (xy - 1)(xy + 1)$
26. $4a^2b^2 - 16c^2 = (2ab)^2 - (4c)^2 = (2ab - 4c)(2ab + 4c)$
27. $3x^2 - 27y^2 = 3(x^2 - 9y^2)$
 $= 3(x - 3y)(x + 3y)$
28. $45 - 5x^2 = 5(9 - x^2)$
 $= 5(3 - x)(3 + x)$
29. $45a^2 - 20 = 5(9a^2 - 4)$
 $= 5(3a - 2)(3a + 2)$

$$30. (2x + y)^2 - 4 = (2x + y - 2)(2x + y + 2)$$

$$31. (3a - 2b)^2 - 9 = (3a - 2b - 3)(3a - 2b + 3)$$

$$32. a^4 - b^4 = (a^2)^2 - (b^2)^2 = (a^2 - b^2)(a^2 + b^2) \\ = (a - b)(a + b)(a^2 + b^2)$$

$$33. x^2 + 9x + 14 = (x + 2)(x + 7)$$

$$34. 2x^2 + 7x + 3 = (2x + 1)(x + 3)$$

$$35. 2x^2 + 11x + 14 = (2x + 7)(x + 2)$$

$$36. x^2 - 9x + 14 = (x - 2)(x - 7)$$

$$37. x^2 - 11x + 28 = (x - 7)(x - 4)$$

$$38. 2x^2 - 7x + 3 = (2x - 1)(x - 3)$$

$$39. 3x^2 - 17x + 20 = (3x - 5)(x - 4)$$

$$40. 7x^2 - 18x + 8 = (7x - 4)(x - 2)$$

$$41. 2x^2 - 7x - 15 = (2x + 3)(x - 5)$$

$$42. 3x^2 + 11x - 20 = (3x - 4)(x + 5)$$

$$43. 12x^2 - 11x - 5 = (4x - 5)(3x + 1)$$

$$44. 6x^2 + x - 15 = (3x + 5)(2x - 3)$$

$$45. 3x^2 + 13x - 10 = (3x - 2)(x + 5)$$

$$46. 6x^2 - 11x + 3 = (3x - 1)(2x - 3)$$

$$47. 36x^2 - 7x - 4 = (9x - 4)(4x + 1)$$

$$48. 15x^2 - 14x - 8 = (5x + 2)(3x - 4)$$

$$49. 6y^2 + 11y - 35 = (3y - 5)(2y + 7)$$

$$50. 12x^2 + 17xy - 5y^2 = (4x - y)(3x + 5y)$$

$$51. (i) x^2 + 3\sqrt{3}x + 6.$$

$$\Rightarrow a = 1, b = 3\sqrt{3}, c = 6.$$

$$\therefore x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3\sqrt{3} \pm \sqrt{27 - 4(1)(6)}}{2(1)} \\ = \frac{-3\sqrt{3} \pm \sqrt{3}}{2}$$

$$\therefore x = \frac{-3\sqrt{3} + \sqrt{3}}{2} \quad \text{or} \quad x = \frac{-3\sqrt{3} - \sqrt{3}}{2}$$

$$\therefore x = -\sqrt{3} \quad \text{or} \quad -2\sqrt{3}$$

$$\therefore \text{Factors are } (x + \sqrt{3}) \text{ and } (x + 2\sqrt{3})$$

$$(ii) x^2 + 2\sqrt{5}x - 15.$$

$$\Rightarrow a = 1, b = 2\sqrt{5}, c = -15.$$

$$\therefore x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-2\sqrt{5} \pm \sqrt{20 - 4(1)(-15)}}{2(1)} \\ = \frac{-2\sqrt{5} \pm \sqrt{80}}{2} \\ = \frac{-2\sqrt{5} \pm 4\sqrt{5}}{2}$$

$$\therefore x = \frac{-2\sqrt{5} + 4\sqrt{5}}{2} \quad \text{or} \quad x = \frac{-2\sqrt{5} - 4\sqrt{5}}{2}$$

$$\therefore x = \sqrt{5} \quad \text{or} \quad -3\sqrt{5}$$

$$\therefore \text{Factors are } (x - \sqrt{5}) \text{ and } (x + 3\sqrt{5})$$

(iii) $2x^2 - 5\sqrt{2}x - 6$.

$\Rightarrow a = 2, b = -5\sqrt{2}, c = -6$.

$$\therefore x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{5\sqrt{2} \pm \sqrt{50 - 4(2)(-6)}}{2(2)}$$

$$= \frac{5\sqrt{2} \pm \sqrt{98}}{4}$$

$$= \frac{5\sqrt{2} \pm 7\sqrt{2}}{4}$$

$$\therefore x = \frac{5\sqrt{2} + 7\sqrt{2}}{4} \quad \text{or} \quad x = \frac{5\sqrt{2} - 7\sqrt{2}}{4}$$

$$\therefore x = 3\sqrt{2} \quad \text{or} \quad \frac{-\sqrt{2}}{2}$$

\therefore Factors are $(x - 3\sqrt{2})$ and $\left(x + \frac{\sqrt{2}}{2}\right)$

But since coefficient of x^2 is 2, one of the factors must

contain $2x$. $\therefore x + \frac{\sqrt{2}}{2} = 0$

$$\Rightarrow 2x + \sqrt{2} = 0$$

\therefore Factors are $(x - 3\sqrt{2})$ and $(2x + \sqrt{2})$

52. (i) $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

(ii) $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

(iii) $8x^3 + y^3 = ((2x)^3 + y^3) = (2x + y)(4x^2 - 2xy + y^2)$

53. (i) $27x^3 - y^3 = (3x)^3 - y^3 = (3x - y)(9x^2 + 3xy + y^2)$

(ii) $x^3 - 64 = x^3 - 4^3 = (x - 4)(x^2 + 4x + 16)$

(iii) $8x^3 - 27y^3 = (2x)^3 - (3y)^3 = (2x - 3y)(4x^2 + 6xy + 9y^2)$

54. (i) $8 + 27k^3 = (2)^3 + (3k)^3 = (2 + 3k)(4 - 6k + 9k^2)$

(ii) $64 - 125a^3 = (4)^3 - (5a)^3 = (4 - 5a)(16 + 20a + 25a^2)$

(iii) $27a^3 + 64b^3 = (3a)^3 + (4b)^3 = (3a + 4b)(9a^2 - 12ab + 16b^2)$

55. (i) $a^3 - 8b^3c^3 = a^3 - (2bc)^3 = (a - 2bc)(a^2 + 2abc + 4b^2c^2)$

(ii) $5x^3 + 40y^3 = 5(x^3 + 8y^3)$

$$= 5(x^3 + (2y)^3)$$

$$= 5(x + 2y)(x^2 - 2xy + 4y^2)$$

(iii) $(x + y)^3 - z^3 = (x + y - z)[(x + y)^2 + (x + y)z + z^2]$

Exercise 1.4

1. (i) $\frac{\cancel{8}^4 \cancel{y}^2}{\cancel{2}^1 \cancel{y}^2} = \frac{4}{y^2}$ (ii) $\frac{\cancel{7}^1 \cancel{a}^6 \cancel{b}^3}{\cancel{14}^2 \cancel{a}^3 \cancel{b}^4} = \frac{a}{2b}$

(iii) $\frac{(2x)^2}{4x} = \frac{\cancel{4}^2 \cancel{x}^2}{\cancel{4}^1 \cancel{x}^1} = x$

(iv) $\frac{7y + 2y^2}{7y} = \frac{y(7 + 2y)}{\cancel{y}^1} = \frac{7 + 2y}{7}$

(v) $\frac{5ax}{15a + 10a^2} = \frac{\cancel{5}^1 \cancel{a}^1 x}{\cancel{5}^1 \cancel{a}^1 (3 + 2a)} = \frac{x}{3 + 2a}$

2. (a) $\frac{2x}{5} + \frac{4x}{3} = \frac{6x}{15} + \frac{20x}{15} = \frac{26x}{15}$

(b) $\frac{3x}{5} - \frac{x}{2} = \frac{6x}{10} - \frac{5x}{10} = \frac{x}{10}$

(c) $\frac{2x + 3}{4} + \frac{x}{3} = \frac{6x + 9}{12} + \frac{4x}{12} = \frac{10x + 9}{12}$

$$(d) \frac{x+1}{4} + \frac{2x-1}{5} = \frac{5x+5}{20} + \frac{8x-4}{20} = \frac{13x+1}{20}$$

$$(e) \frac{3x-4}{6} - \frac{2x+1}{3} = \frac{3x-4}{6} - \frac{4x+2}{6} = \frac{-x-6}{6}$$

$$(f) \frac{3x-2}{6} - \frac{x-3}{4} = \frac{6x-4}{12} - \frac{3x-9}{12} = \frac{3x+5}{12}$$

$$(g) \frac{5x-1}{4} - \frac{2x-4}{5} = \frac{25x-5}{20} - \frac{8x-16}{20} = \frac{17x+11}{20}$$

$$(h) \frac{3x+5}{6} - \frac{2x+3}{4} - \frac{1}{12} = \frac{6x+10}{12} - \frac{6x+9}{12} - \frac{1}{12} \\ = \frac{0}{12} = 0$$

$$(i) \frac{3x-2}{4} + \frac{3}{5} - \frac{2x-1}{10} = \frac{15x-10}{20} + \frac{12}{20} - \frac{4x-2}{20} \\ = \frac{11x+4}{20}$$

$$(j) \frac{1}{3x} + \frac{1}{5x} = \frac{5}{15x} + \frac{3}{15x} = \frac{8}{15x}$$

$$(k) \frac{3}{4x} - \frac{5}{8x} = \frac{6}{8x} - \frac{5}{8x} = \frac{1}{8x}$$

$$(l) \frac{1}{x} + \frac{1}{x+3} = \frac{x+3+x}{x(x+3)} = \frac{2x+3}{x(x+3)}$$

$$(m) \frac{2}{x+2} + \frac{3}{x+4} = \frac{2(x+4)+3(x+2)}{(x+2)(x+4)} = \frac{5x+14}{(x+2)(x+4)}$$

$$(n) \frac{2}{x-2} + \frac{3}{2x-1} = \frac{2(2x-1)+3(x-2)}{(x-2)(2x-1)} = \frac{7x-8}{(x-2)(2x-1)}$$

$$(o) \frac{5}{3x-1} - \frac{2}{x+3} = \frac{5(x+3)-2(3x-1)}{(3x-1)(x+3)} = \frac{-x+17}{(3x-1)(x+3)}$$

$$(p) \frac{3}{2x-7} - \frac{1}{5x+2} = \frac{3(5x+2)-(2x-7)}{(2x-7)(5x+2)} = \frac{13x+13}{(2x-7)(5x+2)}$$

$$(q) \frac{2}{3x-5} - \frac{1}{4} = \frac{8-(3x-5)}{4(3x-5)} = \frac{13-3x}{4(3x-5)}$$

$$(r) \frac{5}{2x-1} - \frac{3}{x-2} = \frac{5(x-2)-3(2x-1)}{(2x-1)(x-2)} = \frac{-x-7}{(2x-1)(x-2)}$$

$$(s) \frac{x}{x-y} - \frac{y}{x+y} = \frac{x(x+y)-y(x-y)}{(x-y)(x+y)} = \frac{x^2 + \cancel{xy} + \cancel{xy} + y^2}{(x-y)(x+y)} \\ = \frac{x^2+y^2}{x^2-y^2}$$

$$(t) \frac{3}{x} + \frac{4}{3y} - \frac{2}{3xy} = \frac{3(3y)+4(x)-2}{3xy} = \frac{9y+4x-2}{3xy}$$

$$(u) \frac{3}{x} - \frac{2}{x-1} - \frac{4}{x(x-1)} = \frac{3(x-1)-2(x)-4}{x(x-1)} = \frac{x-7}{x(x-1)}$$

$$3. (i) \frac{2z^2-4z}{2z^2-10z} = \frac{\cancel{2z}(z-2)}{\cancel{2z}(z-5)} = \frac{z-2}{z-5}$$

$$(ii) \frac{y^2+7y+10}{y^2-25} = \frac{(y+5)(y+2)}{(y+5)(y-5)} = \frac{y+2}{y-5}$$

$$(iii) \frac{t^2+3t-4}{t^2-3t+2} = \frac{(t+4)(\cancel{t-1})}{(t-2)(\cancel{t-1})} = \frac{t+4}{t-2}$$

$$(iv) \frac{x}{x^2-4} - \frac{1}{x+2} = \frac{x}{(x+2)(x-2)} - \frac{1}{x+2} \\ = \frac{x-(x-2)}{(x+2)(x-2)} = \frac{2}{(x+2)(x-2)}$$

$$\begin{aligned} \text{(v)} \quad \frac{2}{a+3} - \frac{a+2}{a^2-9} &= \frac{2}{a+3} - \frac{a+2}{(a+3)(a-3)} \\ &= \frac{2(a-3) - (a+2)}{(a+3)(a-3)} = \frac{a-8}{(a+3)(a-3)} \end{aligned}$$

$$\begin{aligned} \text{(vi)} \quad \frac{x-1}{x^2-4} + \frac{1}{x-2} &= \frac{x-1}{(x-2)(x+2)} + \frac{1}{x-2} \\ &= \frac{x-1+x+2}{(x-2)(x+2)} = \frac{2x+1}{(x-2)(x+2)} \end{aligned}$$

$$\begin{aligned} \text{4. (i)} \quad \frac{10}{2x^2-3x-2} - \frac{2}{x-2} &= \frac{10}{(2x+1)(x-2)} - \frac{2}{(x-2)} \\ &= \frac{10-2(2x+1)}{(2x+1)(x-2)} \\ &= \frac{8-4x}{(2x+1)(x-2)} = \frac{4(2-x)}{(2x+1)(x-2)} \\ &= \frac{-4(\cancel{x-2})}{(2x+1)(\cancel{x-2})} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \frac{x+2}{2x^2-x-1} - \frac{1}{x-1} &= \frac{x+2}{(2x+1)(x-1)} - \frac{1}{x-1} \\ &= \frac{x+2-(2x+1)}{(2x+1)(x-1)} \\ &= \frac{-x+1}{(2x+1)(x-1)} = \frac{\cancel{-(x-1)}}{(2x+1)(\cancel{x-1})} = \frac{-1}{2x+1} \end{aligned}$$

$$\begin{aligned} \text{5. (i)} \quad \frac{1}{x^2-9} - \frac{2}{x^2-x-6} &= \frac{1}{(x-3)(x+3)} - \frac{2}{(x-3)(x+2)} \\ &= \frac{x+2-2(x+3)}{(x-3)(x+3)(x+2)} \\ &= \frac{-x-4}{(x-3)(x+3)(x+2)} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \frac{3}{x^2+x-2} - \frac{2}{x^2+3x+2} &= \frac{3}{(x+2)(x-1)} - \frac{2}{(x+2)(x+1)} \\ &= \frac{3(x+1)-2(x-1)}{(x+2)(x-1)(x+1)} \\ &= \frac{x+5}{(x+2)(x-1)(x+1)} \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad \frac{2}{6x^2-5x-4} - \frac{3}{9x^2-16} &= \frac{2}{(3x-4)(2x+1)} - \frac{3}{(3x-4)(3x+4)} \\ &= \frac{2(3x+4)-3(2x+1)}{(3x-4)(2x+1)(3x+4)} \\ &= \frac{\cancel{6x}+8-\cancel{6x}-3}{(3x-4)(2x+1)(3x+4)} = \frac{5}{(3x-4)(2x+1)(3x+4)} \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad \frac{1}{xy-x^2} - \frac{1}{y^2-xy} &= \frac{1}{x(y-x)} - \frac{1}{y(y-x)} \\ &= \frac{y-x}{x(y-x)y} \\ &= \frac{\cancel{(y-x)}}{\cancel{(y-x)}xy} = \frac{1}{xy} \end{aligned}$$

$$6. \quad (i) \quad \frac{\frac{1}{2} + \frac{3}{4}}{\frac{1}{4}} = \frac{\frac{2}{4} + \frac{3}{4}}{\frac{1}{4}} = \frac{\frac{5}{4}}{\frac{1}{4}} = \frac{5}{4} \times \frac{4}{1} = 5$$

$$(ii) \quad \frac{\frac{2}{3} + \frac{5}{6}}{\frac{3}{8}} = \frac{\frac{4}{6} + \frac{5}{6}}{\frac{3}{8}} = \frac{\frac{9}{6}}{\frac{3}{8}} = \frac{\cancel{9}^3}{\cancel{3}_1} \times \frac{\cancel{8}^4}{\cancel{3}_1} = 4$$

$$(iii) \quad \frac{x - \frac{1}{x}}{1 + \frac{1}{x}} = \frac{\frac{x^2 - 1}{x}}{\frac{x+1}{x}} = \frac{x^2 - 1}{x+1} = \frac{(x-1)\cancel{(x+1)}}{\cancel{(x+1)}} = x - 1$$

$$7. \quad (i) \quad \frac{\frac{1}{x} + 1}{\frac{1}{x} - 1} = \frac{\frac{1+x}{x}}{\frac{1-x}{x}} = \frac{(1+x)}{\cancel{x}} \cdot \frac{\cancel{x}}{(1-x)} = \frac{1+x}{1-x}$$

$$(ii) \quad \frac{\frac{1}{x^2} - 4}{\frac{1}{x} - 2} = \frac{\frac{1-4x^2}{x^2}}{\frac{1-2x}{x}} = \frac{(1-4x^2)}{x^2} \cdot \frac{x^1}{(1-2x)} = \frac{(1-2x)(1+2x)}{x(1-2x)} = \frac{1+2x}{x}$$

$$(iii) \quad \frac{x+y}{\frac{1}{x} + \frac{1}{y}} = \frac{x+y}{\frac{y+x}{xy}} = \frac{\cancel{(x+y)}}{1} \cdot \frac{xy}{\cancel{(x+y)}} = xy$$

$$8. \quad (i) \quad \frac{4y - \frac{3}{2}}{2} = \frac{\frac{8y-3}{2}}{2} = \frac{8y-3}{2} \times \frac{1}{2} = \frac{8y-3}{4}$$

$$(ii) \quad \frac{2 - \frac{1}{x}}{2} = \frac{\frac{2x-1}{x}}{2} = \frac{2x-1}{x} \cdot \frac{1}{2} = \frac{2x-1}{2x}$$

$$(iii) \quad \frac{3x + \frac{1}{x}}{2} = \frac{\frac{3x^2+1}{x}}{2} = \frac{3x^2+1}{x} \cdot \frac{1}{2} = \frac{3x^2+1}{2x}$$

$$(iv) \quad \frac{y + \frac{1}{4}}{\frac{1}{2}} = \frac{\frac{4y+1}{4}}{\frac{1}{2}} = \frac{4y+1}{4} \cdot \frac{\cancel{2}^1}{\cancel{1}_2} = \frac{4y+1}{2}$$

$$9. \quad (i) \quad \frac{z - \frac{1}{3}}{z - \frac{1}{2}} = \frac{\frac{3z-1}{3}}{\frac{2z-1}{2}} = \frac{3z-1}{3} \cdot \frac{2}{2z-1} = \frac{6z-2}{6z-3}$$

$$(ii) \quad \frac{2x + \frac{1}{2}}{x + \frac{1}{4}} = \frac{\frac{4x+1}{2}}{\frac{4x+1}{4}} = \frac{\cancel{(4x+1)}}{\cancel{2}_1} \cdot \frac{\cancel{4}^2}{\cancel{(4x+1)}} = 2$$

$$(iii) \quad \frac{z - \frac{1}{2z}}{z - \frac{1}{3z}} = \frac{\frac{2z^2-1}{2z}}{\frac{3z^2-1}{3z}} = \frac{(2z^2-1)}{\cancel{2z}} \cdot \frac{\cancel{3z}}{(3z^2-1)} = \frac{6z^2-3}{6z^2-2}$$

$$(iv) \quad \frac{x - \frac{1}{x+1}}{x-1} = \frac{\frac{x(x+1)-1}{x+1}}{x-1} = \frac{x^2+x-1}{x+1} \cdot \frac{1}{x-1} = \frac{x^2+x-1}{x^2-1}$$

$$10. \quad (i) \quad \frac{1 + \frac{2}{x}}{\frac{x+2}{x-2}} = \frac{\frac{x+2}{x}}{\frac{x+2}{x-2}} = \frac{\cancel{(x+2)}}{x} \cdot \frac{x-2}{\cancel{(x+2)}} = \frac{x-2}{x}$$

$$(ii) \quad \frac{2 + \frac{1}{x}}{2x^2 + x} = \frac{\frac{2x+1}{x}}{x(2x+1)} = \frac{\cancel{(2x+1)}}{x} \cdot \frac{1}{\cancel{x(2x+1)}} = \frac{1}{x^2}$$

$$\begin{aligned}
 \text{(iii)} \quad \frac{x + \frac{2x}{x-2}}{1 + \frac{4}{(x+2)(x-2)}} &= \frac{\frac{x(x-2) + 2x}{(x-2)}}{\frac{(x+2)(x-2) + 4}{(x+2)(x-2)}} \\
 &= \frac{x^2 - \cancel{2x} + \cancel{2x}}{(x-2)} \cdot \frac{(x+2)(x-2)}{x^2 - \cancel{4} + \cancel{4}} \\
 &= \frac{\cancel{x^2}}{\cancel{(x-2)}} \cdot \frac{(x+2)\cancel{(x-2)}}{\cancel{x^2}} = x + 2
 \end{aligned}$$

$$\begin{aligned}
 \text{11. (i)} \quad \frac{\left(\frac{a+b}{a-b}\right) - \left(\frac{a-b}{a+b}\right)}{1 + \left(\frac{a-b}{a+b}\right)} &= \frac{\frac{(a+b)(a+b) - (a-b)(a-b)}{(a-b)(a+b)}}{\frac{\cancel{(a+b)} + \cancel{(a-b)}}{(a+b)}} \\
 &= \frac{a^2 + 2ab + b^2 - (a^2 - 2ab + b^2)}{(a-b)(a+b)} \cdot \frac{(a+b)}{2a} \\
 &= \frac{\cancel{a^2} + 2ab + \cancel{b^2} - \cancel{a^2} + 2ab - \cancel{b^2}}{(a-b)(a+b)} \cdot \frac{(a+b)}{2a} \\
 &= \frac{\cancel{2ab} \cancel{(a+b)}}{(a-b)\cancel{(a+b)} \cancel{2a}} = \frac{2b}{a-b}
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad \frac{x + \frac{3}{x}}{x - \frac{9}{x^3}} &= \frac{\frac{x^2+3}{x}}{\frac{x^4-9}{x^3}} = \frac{(x^2+3)}{\cancel{x}} \cdot \frac{x^{\cancel{3}^2}}{(x^4-9)} \\
 &= \frac{\cancel{(x^2+3)}(x^2)}{(x^2-3)\cancel{(x^2+3)}} = \frac{x^2}{x^2-3}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii)} \quad \frac{9 - \frac{1}{y^2}}{9 + \frac{6}{y} + \frac{1}{y^2}} &= \frac{\frac{9y^2-1}{y^2}}{\frac{9y^2+6y+1}{y^2}} \\
 &= \frac{9y^2-1}{\cancel{y^2}} \cdot \frac{\cancel{y^2}}{9y^2+6y+1} \\
 &= \frac{(3y-1)\cancel{(3y+1)}}{(3y+1)\cancel{(3y+1)}} = \frac{3y-1}{3y+1}
 \end{aligned}$$

$$\begin{aligned}
 \text{12.} \quad \frac{3x-5}{x-2} + \frac{1}{2-x} &= \frac{(3x-5)(2-x) + (x-2)}{(x-2)(2-x)} \\
 &= \frac{6x - 3x^2 - 10 + 5x + x - 2}{(x-2)(2-x)} \\
 &= \frac{-3x^2 + 12x - 12}{(x-2)(2-x)} \\
 &= \frac{-3(x^2 - 4x + 4)}{(x-2)(2-x)} \\
 &= \frac{-3\cancel{(x-2)}\cancel{(x-2)}}{-\cancel{(x-2)}\cancel{(x-2)}} = 3.
 \end{aligned}$$

Exercises 1.5

1. By calculator:

$$\text{(i)} \quad \binom{7}{4} = 35 \quad \text{(ii)} \quad \binom{6}{2} = 15 \quad \text{(iii)} \quad \binom{6}{4} = 15$$

$$\text{(iv)} \quad \binom{15}{4} = 1365 \quad \text{(v)} \quad \binom{10}{9} = 10$$

2. (i) $\binom{9}{0} = 1$ (ii) $\binom{10}{1} = 10$ (iii) $\binom{13}{13} = 1$

(iv) $\binom{30}{0} = 1$ (v) $\binom{18}{17} = \binom{18}{1} = 18$

3. $\binom{12}{3} = \binom{12}{12-3} = \binom{12}{9}$. Hence $k = 9$.

4. $\binom{16}{12} = \binom{16}{16-12} = \binom{16}{4}$. Hence $k = 4$.

5. $(a + 2b)^4 = \binom{4}{0}a^4 + \binom{4}{1}a^3(2b) + \binom{4}{2}a^2(2b)^2 + \binom{4}{3}a(2b)^3 + \binom{4}{4}(2b)^4$
 $= (1)a^4 + (4)a^3(2b) + (6)a^2(4b^2) + (4)a(8b^3) + (1)(16b^4)$
 $= a^4 + 8a^3b + 24a^2b^2 + 32ab^3 + 16b^4$

6. The coefficients in the 6th row of Pascal's triangle are 1 6 15 20 15 6 1.

Thus

$$(x + y)^6 = x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + y^6.$$

7. The fifth row of Pascal's triangle is 1 5 10 10 ...

The coefficient of the 4th term is thus 10. The 4th term is then

$$10(1)^2(2x)^3 = 10(8x^3) = 80x^3$$

Thus the coefficient is 80.

8. (i) $(a - 2b)^4 = \binom{4}{0}a^4 + \binom{4}{1}a^3(-2b) + \binom{4}{2}a^2(-2b)^2 + \binom{4}{3}a(-2b)^3 + \binom{4}{4}(-2b)^4$
 $= (1)a^4 + (4)a^3(-2b) + (6)a^2(4b^2) + (4)a(-8b^3) + (1)(16b^4)$
 $= a^4 - 8a^3b + 24a^2b^2 - 32ab^3 + 16b^4$

(ii) $(2x - y)^3 = \binom{3}{0}(2x)^3 + \binom{3}{1}(2x)^2(-y) + \binom{3}{2}(2x)(-y)^2 + \binom{3}{3}(-y)^3$
 $= (1)(8x^3) + (3)(4x^2)(-y) + (3)(2x)(y^2) + (1)(-y^3)$
 $= 8x^3 - 12x^2y + 6xy^2 - y^3$

(iii) $(p + 3q)^4 = \binom{4}{0}p^4 + \binom{4}{1}p^3(3q) + \binom{4}{2}p^2(3q)^2 + \binom{4}{3}p(3q)^3 + \binom{4}{4}(3q)^4$
 $= (1)p^4 + 4p^3(3q) + (6)p^2(9q^2) + (4)p(27q^3) + (1)(81q^4)$
 $= p^4 + 12p^3q + 54p^2q^2 + 108pq^3 + 81q^4$

(iv) $(1 + 2y)^5 = \binom{5}{0}(1)^5 + \binom{5}{1}(1)^4(2y) + \binom{5}{2}(1)^3(2y)^2 + \binom{5}{3}(1)^2(2y)^3$
 $+ \binom{5}{4}(1)(2y)^4 + \binom{5}{5}(2y)^5$
 $= (1)(1) + (5)(1)(2y) + (10)(1)(4y^2) + (10)(1)(8y^3) + (5)(1)(16y^4) + (1)(32y^5)$
 $= 1 + 10y + 40y^2 + 80y^3 + 80y^4 + 32y^5$

9. (i) $(2 + 3p)^6 = \binom{6}{0}2^6 + \binom{6}{1}2^5(3p) + \binom{6}{2}2^4(3p)^2 + \binom{6}{3}2^3(3p)^3 + \binom{6}{4}2^2(3p)^4$
 $+ \binom{6}{5}2(3p)^5 + \binom{6}{6}(3p)^6$
 $= (1)(64) + (6)(32)(3p) + (15)(16)(9p^2) + (20)(8)(27p^3) + (15)(4)(81p^4)$
 $+ (6)(2)(243p^5) + (1)(729p^6)$
 $= 64 + 576p + 2160p^2 + 4320p^3 + 4860p^4 + 2916p^5 + 729p^6$

(ii) $(1 - b)^7 = \binom{7}{0}1^7 + \binom{7}{1}1^6(-b) + \binom{7}{2}1^5(-b)^2 + \binom{7}{3}1^4(-b)^3 + \binom{7}{4}1^3(-b)^4$
 $+ \binom{7}{5}1^2(-b)^5 + \binom{7}{6}1(-b)^6 + \binom{7}{7}(-b)^7$
 $= (1)(1) + (7)(1)(-b) + (21)(1)(b^2) + (35)(1)(-b^3) + (35)(1)(b^4)$
 $+ (21)(1)(-b^5) + (7)(1)(b^6) + (1)(-b^7)$
 $= 1 - 7b + 21b^2 - 35b^3 + 35b^4 - 21b^5 + 7b^6 - b^7$

$$\begin{aligned}
 \text{(iii)} \quad (p - 4q)^5 &= \binom{5}{0}p^5 + \binom{5}{1}p^4(-4q) + \binom{5}{2}p^3(-4q)^2 + \binom{5}{3}p^2(-4q)^3 \\
 &\quad + \binom{5}{4}p(-4q)^4 + \binom{5}{5}(-4q)^5 \\
 &= (1)p^5 + (5)p^4(-4q) + (10)p^3(16q^2) + (10)p^2(-64q^3) + (5)p(256q^4) \\
 &\quad + (1)(-1024q^5) \\
 &= p^5 - 20p^4q + 160p^3q^2 - 640p^2q^3 + 1280pq^4 - 1024q^5
 \end{aligned}$$

10. The coefficients in the 7th row are then

$$\begin{array}{cccccccc}
 1 & 1+6 & 6+15 & 15+20 & 20+15 & 15+6 & 6+1 & 1 \\
 \text{or } 1 & 7 & 21 & 35 & 35 & 21 & 7 & 1.
 \end{array}$$

11. General term $= \binom{8}{r}x^{8-r}y^r$. For the 5th term, $r = 4$. Thus the 5th term is

$$\binom{8}{4}x^{8-4}y^4 = 70x^4y^4.$$

12. General term $= \binom{9}{r}x^{9-r}(-y)^r$. For the 4th term, $r = 3$. Thus the 4th term is

$$\binom{9}{3}x^{9-3}(-y)^3 = (84)x^6(-y^3) = -84x^6y^3$$

13. General term $= \binom{10}{r}(2x)^{10-r}y^r$. For the 6th term, $r = 5$. Thus the 6th term is

$$\binom{10}{5}(2x)^5y^5 = (252)(32x^5)y^5 = 8064x^5y^5$$

14. The number of terms in $(p + 2q)^6$ is $6 + 1 = 7$.

The middle term is then the 4th term (3 before and 3 after)

The general term is $\binom{6}{r}p^{6-r}(2q)^r$. For the 4th term, $r = 3$. Thus the middle term is

$$\binom{6}{3}p^{6-3}(2q)^3 = (20)p^3(8q^3) = 160p^3q^3.$$

The coefficient of this term is 160.

15. (i) $(2x - y)^8 = \binom{8}{0}(2x)^8 + \binom{8}{1}(2x)^7(-y) + \binom{8}{2}(2x)^6(-y)^2 + \binom{8}{3}(2x)^5(-y)^3$

$$\begin{aligned}
 &+ \binom{8}{4}(2x)^4(-y)^4 + \binom{8}{5}(2x)^3(-y)^5 + \binom{8}{6}(2x)^2(-y)^6 \\
 &+ \binom{8}{7}(2x)(-y)^7 + \binom{8}{8}(-y)^8 \\
 &= (1)(256x^8) + (8)(128x^7)(-y) + (28)(64x^6)(y^2) + (56)(32x^5)(-y^3) \\
 &\quad + (70)(16x^4)(y^4) + (56)(8x^3)(-y^5) + (28)(4x^2)(y^6) \\
 &\quad + (8)(2x)(-y^7) + (1)(y^8) \\
 &= 256x^8 - 1024x^7y + 1792x^6y^2 - 1792x^5y^3 + 1120x^4y^4 - 448x^3y^5 \\
 &\quad + 112x^2y^6 - 16xy^7 + y^8
 \end{aligned}$$

(ii) $(a + 2b)^9 = \binom{9}{0}a^9 + \binom{9}{1}a^8(2b) + \binom{9}{2}a^7(2b)^2 + \binom{9}{3}a^6(2b)^3 + \binom{9}{4}a^5(2b)^4$

$$\begin{aligned}
 &+ \binom{9}{5}a^4(2b)^5 + \binom{9}{6}a^3(2b)^6 + \binom{9}{7}a^2(2b)^7 + \binom{9}{8}a(2b)^8 + \binom{9}{9}(2b)^9 \\
 &= (1)a^9 + (9)a^8(2b) + (36)a^7(4b^2) + (84)a^6(8b^3) + (126)a^5(16b^4) \\
 &\quad + (126)a^4(32b^5) + (84)a^3(64b^6) + (36)a^2(128b^7) \\
 &\quad + (9)a(256b^8) + (1)(512b^9) \\
 &= a^9 + 18a^8b + 144a^7b^2 + 672a^6b^3 + 2016a^5b^4 + 4032a^4b^5 + 5376a^3b^6 \\
 &\quad + 4608a^2b^7 + 2304ab^8 + 512b^9
 \end{aligned}$$

16. The general term in the expansion of $(5x + 1)^{10}$ is

$$\binom{10}{r}(5x)^{10-r}(1)^r.$$

For the 3rd term, $r = 2$. Thus the 3rd term is

$$\binom{10}{2}(5x)^8(1)^2 = (45)(390625x^8) = 17578125x^8$$

17. The general term in the expansion of $\left(x - \frac{3y}{2}\right)^9$ is

$$\binom{9}{r}x^{9-r}\left(\frac{-3y}{2}\right)^r$$

For the 4th term, $r = 3$. Thus the 4th term is

$$\binom{9}{3}x^6\left(\frac{-3y}{2}\right)^3 = (84)x^6\left(\frac{-27y^3}{8}\right) = -\frac{567}{2}x^6y^3$$

The coefficient is $-\frac{567}{2} = -283.5$.

Exercise 1.6

1. $ax^2 + bx + c = (2x - 3)(3x + 4)$ for all x
 $= 6x^2 + 8x - 9x - 12$
 $= 6x^2 - x - 12$
 $\therefore a = 6, b = -1, c = -12$

2. $(3x - 2)(x + 5) = 3x^2 + px + q$ for all x
 $3x^2 + 15x - 2x - 10 = 3x^2 + px + q$
 $3x^2 + 13x - 10 = 3x^2 + px + q$
 $\therefore p = 13, q = -10$

3. $x^2 + 6x + 16 = (x + a)^2 + b$ for all x
 $x^2 + 6x + 16 = x^2 + 2ax + a^2 + b$
 $\therefore 2a = 6 \Rightarrow a = 3$
and $a^2 + b = 16$
 $\therefore 9 + b = 16 \Rightarrow b = 7$

4. $x^2 + 4x - 6 = (x + a)^2 + b$ for all x
 $x^2 + 4x - 6 = x^2 + 2ax + a^2 + b$
 $\therefore 2a = 4 \Rightarrow a = 2$
and $a^2 + b = -6$
 $\therefore 4 + b = -6 \Rightarrow b = -10$

5. $2x^2 + 5x + 6 = p(x + q)^2 + r$ for all x
 $= p(x^2 + 2xq + q^2) + r$
 $= px^2 + 2pqx + pq^2 + r$
 $\therefore p = 2$
and $2pq = 5$
 $\therefore 2(2)q = 5 \Rightarrow q = \frac{5}{4}$
and $pq^2 + r = 6$
 $\therefore 2\left(\frac{5}{4}\right)^2 + r = 6$
 $\frac{25}{8} + r = 6 \Rightarrow r = 6 - \frac{25}{8}$
 $= \frac{48 - 25}{8} = \frac{23}{8}$

6. $(2x + a)^2 = 4x^2 + 12x + b$ for all values of x

$$4x^2 + 4ax + a^2 = 4x^2 + 12x + b$$

$$\therefore 4a = 12 \Rightarrow a = 3$$

$$\text{and } a^2 = b \Rightarrow 9 = b$$

7. $x^2 - 4x - 5 = (x - n)^2 - m$ for all values of x

$$x^2 - 4x - 5 = x^2 - 2nx + n^2 - m$$

$$\therefore -4 = -2n \Rightarrow n = 2$$

$$\text{and } -5 = n^2 - m$$

$$\therefore -5 = 4 - m \Rightarrow m = 9$$

8. (i) $V(x) = ax^3 + bx^2 + cx + d = (x + 5)(x + 3)(x + 2)$ for all x

$$= (x^2 + 8x + 15)(x + 2)$$

$$= x^3 + 2x^2 + 8x^2 + 16x + 15x + 30$$

$$= x^3 + 10x^2 + 31x + 30$$

$$\therefore a = 1, b = 10, c = 31, d = 30.$$

(ii) $S(x) = px^2 + qx + r = 2(x + 3)(x + 2) + 2(x + 5)(x + 3) + (x + 5)(x + 2)$

$$= 2(x^2 + 5x + 6) + 2(x^2 + 8x + 15) + (x^2 + 7x + 10)$$

$$= 5x^2 + 33x + 52$$

$$\therefore p = 5, q = 33, r = 52$$

9. $3(x - p)^2 + q = 3x^2 - 12x + 7$ for all x

$$\therefore 3(x^2 - 2px + p^2) + q = 3x^2 - 12x + 7$$

$$\therefore 3x^2 - 6px + 3p^2 + q = 3x^2 - 12x + 7$$

$$\therefore -6p = -12 \Rightarrow p = 2$$

$$\text{and } 3p^2 + q = 7$$

$$\therefore 3(2)^2 + q = 7 \Rightarrow q = -5$$

10. $V(x) = x^3 + 12x^2 + bx + 30 = (x^2 + cx + 4)(x + a)$

$$= x^3 + ax^2 + cx^2 + acx + 4x + 4a$$

$$= x^3 + x^2(a + c) + x(ac + 4) + 4a$$

$$\therefore a + c = 12$$

$$\text{and } b = ac + 4$$

$$\text{and } 4a = 30 \Rightarrow a = \frac{30}{4} = \frac{15}{2}.$$

$$\therefore a + c = 12 \Rightarrow c = 12 - a$$

$$c = 12 - \frac{15}{2} = \frac{9}{2}$$

$$\therefore b = ac + 4 \Rightarrow b = \frac{15}{2} \cdot \frac{9}{2} + 4 = \frac{135 + 16}{4} = 37\frac{3}{4}$$

11. $(x - 4)^3 = x^3 + px^2 + qx - 64$ for all x

$$= (x - 4)(x - 4)(x - 4)$$

$$= (x^2 - 8x + 16)(x - 4)$$

$$= x^3 - 4x^2 - 8x^2 + 32x + 16x - 64$$

$$= x^3 - 12x^2 + 48x - 64$$

$$\therefore p = -12, q = 48$$

12. $(x + a)(x^2 + bx + 2) = x^3 - 2x^2 - x - 6$ for all x

$$= x^3 + bx^2 + 2x + ax^2 + abx + 2a$$

$$= x^3 + x^2(b + a) + x(2 + ab) + 2a$$

$$\therefore b + a = -2$$

$$\text{and } 2 + ab = -1$$

$$\text{and } 2a = -6 \Rightarrow a = -3$$

$$\therefore b + a = -2 \Rightarrow b - 3 = -2 \Rightarrow b = 1$$

$$\begin{aligned}
 13. \quad (x-2)(x^2+bx+c) &= x^3+2x^2-5x-6 \quad \text{for all } x \\
 &= x^3+bx^2+cx-2x^2-2bx-2c \\
 &= x^3+x^2(b-2)+x(c-2b)-2c \\
 \therefore b-2 &= 2 \Rightarrow b=4 \\
 \text{and } c-2b &= -5 \Rightarrow c-2(4) = -5 \Rightarrow c=3
 \end{aligned}$$

$$\begin{aligned}
 14. \quad (5a-b)x+b+2c &= 0 \quad \text{for all } x \\
 \therefore (5a-b)x+b+2c &= 0.x+0 \\
 \therefore 5a-b &= 0 \Rightarrow b=5a \\
 \text{and } b+2c &= 0 \Rightarrow b=-2c \\
 \therefore 5a &= -2c \\
 a &= \frac{-2c}{5}
 \end{aligned}$$

$$\begin{aligned}
 15. \quad (4x+r)(x^2+s) &= 4x^3+px^2+qx+2 \quad \text{for all } x \\
 &= 4x^3+4xs+rx^2+rs \\
 &= 4x^3+rx^2+4xs+rs \\
 \therefore r &= p \\
 \text{and } 4s &= q \\
 \text{and } rs &= 2 \quad \left. \vphantom{\begin{matrix} r=p \\ 4s=q \end{matrix}} \right\} \Rightarrow pq = r.4s = 4rs \\
 \therefore pq &= 4rs = 4(2) = 8
 \end{aligned}$$

$$\begin{aligned}
 16. \quad (x+s)(x-s)(ax+t) &= ax^3+bx^2+cx+d \quad \text{for all } x \\
 &= (x^2-s^2)(ax+t) \\
 &= ax^3+tx^2-as^2x-ts^2 \\
 \therefore t &= b \\
 \text{and } -as^2 &= c \\
 \text{and } -ts^2 &= d \quad \left. \vphantom{\begin{matrix} t=b \\ -as^2=c \end{matrix}} \right\} \therefore \frac{-as^2}{-ts^2} = \frac{c}{d} \\
 &\Rightarrow -ad = -ct \\
 \text{but } t &= b \quad \therefore -ad = -cb \\
 &\Rightarrow ad = cb
 \end{aligned}$$

$$\begin{aligned}
 17. \quad \frac{1}{(x+1)(x-1)} &= \frac{A}{(x+1)} + \frac{B}{(x-1)} \quad \text{for all } x \\
 \Rightarrow \frac{1}{(x+1)(x-1)} &= \frac{A(x-1)+B(x+1)}{(x+1)(x-1)} \\
 \Rightarrow 1 &= A(x-1)+B(x+1) \\
 \Rightarrow 1 &= Ax-A+Bx+B \\
 \Rightarrow 1 &= x(A+B)-A+B \\
 \therefore 0.x+1 &= (A+B)x-A+B \\
 \therefore A+B &= 0 \\
 \text{and } -A+B &= 1 \\
 \text{adding: } 2B &= 1 \\
 B &= \frac{1}{2} \\
 \text{since } A+B &= 0 \Rightarrow A = -B = -\frac{1}{2}.
 \end{aligned}$$

$$\begin{aligned}
 18. \quad \frac{1}{(x+2)(x-3)} &= \frac{C}{(x+2)} + \frac{D}{(x-3)} \\
 \frac{1}{(x+2)(x-3)} &= \frac{C(x-3)+D(x+2)}{(x+2)(x-3)} \\
 \Rightarrow 1 &= C(x-3)+D(x+2) \\
 1 &= Cx-3C+Dx+2D \\
 0.x+1 &= x(C+D)-3C+2D
 \end{aligned}$$

$$\begin{array}{l} \Rightarrow C + D = 0 \\ \times 3 \text{ and } -3C + 2D = 1 \\ \hline \Rightarrow 3C + 3D = 0 \\ \therefore \text{adding: } 5D = 1 \Rightarrow \end{array}$$

$$D = \frac{1}{5}$$

since $C + D = 0 \Rightarrow C = -D = -\frac{1}{5}$.

$$19. \frac{1}{(x+1)(x+4)} = \frac{A}{(x+1)} + \frac{B}{(x+4)}$$

$$\Rightarrow \frac{1}{(x+1)(x+4)} = \frac{A(x+4) + B(x+1)}{(x+1)(x+4)}$$

$$\Rightarrow 1 = Ax + 4A + Bx + B$$

$$0.x + 1 = x(A + B) + 4A + B$$

$$\therefore A + B = 0$$

$$\text{and } 4A + B = 1$$

$$\text{subtracting: } -3A = -1 \Rightarrow A = \frac{1}{3}$$

$$\text{since } A + B = 0 \Rightarrow B = -A = -\frac{1}{3}.$$

$$20. (x-3)^2 \text{ is a factor of } x^3 + ax + b$$

$$\Rightarrow (x-3)^2(x+k) = x^3 + ax + b$$

$$\therefore (x^2 - 6x + 9)(x+k) = x^3 + ax + b$$

$$\therefore x^3 + kx^2 - 6x^2 - 6kx + 9x + 9k = x^3 + ax + b$$

$$\therefore x^3 + x^2(k-6) + x(-6k+9) + 9k = x^3 + ax + b = x^3 + 0.x^2 + ax + b$$

$$\therefore k-6=0 \Rightarrow k=6$$

$$\text{also } -6k+9=a \quad \therefore -6(6)+9=a \Rightarrow a=-27$$

$$\text{also } 9k=b \quad \therefore 9(6)=b \Rightarrow b=54.$$

$$21. (x-2)^2 \text{ is a factor of } x^3 + px + q$$

$$\Rightarrow (x-2)^2(x-k) = x^3 + px + q$$

$$\therefore (x^2 - 4x + 4)(x-k) = x^3 + 0.x^2 + px + q$$

$$\therefore x^3 - kx^2 - 4x^2 + 4kx + 4x - 4k = x^3 + 0.x^2 + px + q$$

$$\therefore x^3 + x^2(-k-4) + x(4k+4) - 4k =$$

$$\therefore -k-4=0$$

$$\Rightarrow k=-4$$

$$\text{also } p=4k+4=4(-4)+4=-12$$

$$q=-4k=-4(-4)=+16.$$

$$22. (x^2-4) \text{ is a factor of } x^3 + cx^2 + dx - 12$$

$$\therefore (x^2-4)(x+k) = x^3 + cx^2 + dx - 12$$

$$\therefore x^3 + kx^2 - 4x - 4k = x^3 + cx^2 + dx - 12$$

$$\therefore k=c$$

$$\text{also } d=-4$$

$$\text{and } -4k=-12 \Rightarrow k=3=c$$

$$\therefore (x^2-4)(x+3) = x^3 + 3x^2 - 4x - 12$$

$$\Rightarrow (x-2)(x+2)(x+3) = x^3 + 3x^2 - 4x - 12.$$

$$23. (x^2+b) \text{ is a factor of } x^3 - 3x^2 + bx - 15$$

$$\Rightarrow (x^2+b)(x+k) = x^3 - 3x^2 + bx - 15$$

$$\therefore x^3 + kx^2 + bx + bk = x^3 - 3x^2 + bx - 15$$

$$\Rightarrow k=-3$$

$$\text{also } bk=-15 \Rightarrow b = \frac{-15}{k} = \frac{-15}{-3} = 5.$$

24. $x^2 - px + 9$ is a factor of $x^3 + ax + b$

$$\begin{aligned}\therefore (x^2 - px + 9)(x + k) &= x^3 + 0x^2 + ax + b \\ \therefore x^3 + kx^2 - px^2 - pkx + 9x + 9k &= x^3 + 0x^2 + ax + b \\ \therefore x^3 + x^2(k - p) + x(-pk + 9) + 9k &= x^3 + 0x^2 + ax + b \\ \therefore k - p = 0 &\Rightarrow k = p \\ -pk + 9 = a &\Rightarrow -p(p) + 9 = a \\ \therefore a &= 9 - p^2 \\ \text{also } b = 9k &\Rightarrow b = 9p \\ a + b &= 17 \\ \Rightarrow 9 - p^2 + 9p &= 17 \\ -p^2 + 9p - 8 &= 0 \\ p^2 - 9p + 8 &= 0 \\ (p - 8)(p - 1) &= 0 \\ \therefore p &= 8, 1\end{aligned}$$

25. $x^2 - kx + 1$ is a factor of $ax^3 + bx + c$

$$\begin{array}{r} ax + ak \\ x^2 - kx + 1 \overline{) ax^3 + + bx + c} \\ \underline{ax^3 - akx^2 + ax} \\ akx^2 + bx - ax + c \\ \underline{akx^2 + (b - a)x + c} \\ \underline{akx^2 - ak^2x + ak} \\ (b - a)x + ak^2x + c - ak \\ (b - a + ak^2)x + c - ak \text{ (remainder)} \end{array}$$

Since $x^2 - kx + 1$ is a factor, there can be no remainder.

$$\therefore (b - a + ak^2)x + c - ak = 0 \quad \text{for all } x$$

$$\Rightarrow b - a + ak^2 = 0$$

$$\text{and } c - ak = 0 \Rightarrow k = \frac{c}{a}$$

$$\therefore b - a + a\left(\frac{c}{a}\right)^2 = 0$$

$$\therefore b - a + \frac{c^2}{a} = 0 \Rightarrow ab - a^2 + c^2 = 0$$

$$\Rightarrow c^2 = a^2 - ab = a(a - b)$$

26. $(x - a)^2$ is a factor of $x^3 + 3px + c$

i.e. $x^2 - 2ax + a^2$ is a factor of $x^3 + 3px + c$.

$$\begin{array}{r} x + 2a \\ x^2 - 2ax + a^2 \overline{) x^3 + + 3px + c} \\ \underline{x^3 - 2ax^2 + a^2x} \\ 2ax^2 + 3px - a^2x + c \\ \underline{2ax^2 + (3p - a^2)x + c} \\ \underline{2ax^2 - 4a^2x + 2a^3} \\ (3p - a^2)x + 4a^2x + c - 2a^3 \text{ (remainder)} \end{array}$$

Since $x^2 - 2ax + a^2$ is a factor, there can be no remainder.

$$\therefore (3p - a^2 + 4a^2)x + c - 2a^3 = 0 \quad \text{for all } x$$

$$\therefore 3p - a^2 + 4a^2 = 0 \Rightarrow 3p = -3a^2 \Rightarrow p = -a^2$$

$$\text{also, } c - 2a^3 = 0 \Rightarrow c = 2a^3.$$

27. $x^2 + ax + b$ is a factor of $x^3 - k$.

$$\begin{array}{r}
 \overline{x - a} \\
 \therefore x^2 + ax + b \overline{\begin{array}{r}
 x^3 - k \\
 \underline{x^3 + ax^2 + bx} \\
 -ax^2 - bx - k \\
 \underline{-ax^2 - a^2x - b} \\
 -bx + a^2x - k + b \\
 x(-b + a^2) - k + ba \text{ (remainder)}
 \end{array}}
 \end{array}$$

Since $x^2 + ax + b$ is a factor, there can be no remainder.

$$\therefore (-b + a^2) = 0 \Rightarrow b = a^2$$

$$(i) \text{ also, } -k + ba = 0 \Rightarrow k = ab = a \cdot a^2 = a^3$$

$$(ii) \text{ Since } b = a^2 \Rightarrow b^3 = (a^2)^3 = a^6 = k^2.$$

28.

$$\begin{array}{r}
 \phantom{2x - \sqrt{3}} \overline{2x - 1} \\
 2x - \sqrt{3} \overline{\begin{array}{r}
 4x^2 - 2(1 + \sqrt{3})x + \sqrt{3} \\
 \underline{4x^2 - 2\sqrt{3}x} \\
 -2(1 + \sqrt{3})x + 2\sqrt{3}x + \sqrt{3} \\
 -2x - 2\sqrt{3}x + 2\sqrt{3}x + \sqrt{3} \\
 -2x + \sqrt{3} \\
 \underline{-2x + \sqrt{3}} \\
 0
 \end{array}}
 \end{array}$$

$\therefore 2x - \sqrt{3}$ is a factor

and the second factor is $2x - 1$.

29. $5x + 3 = Ax(x + 3) + Bx(x - 1) + C(x - 1)(x + 3)$ for all x

$$\begin{aligned}
 5x + 3 &= Ax^2 + 3Ax + Bx^2 - Bx + C(x^2 + 3x - x - 3) \\
 &= Ax^2 + 3Ax + Bx^2 - Bx + Cx^2 + 3Cx - Cx - 3C \\
 &= Ax^2 + 3Ax + Bx^2 - Bx + Cx^2 + 2Cx - 3C \\
 &= (A + B + C)x^2 + (3A - B + 2C)x - 3C
 \end{aligned}$$

$$\therefore A + B + C = 0$$

$$\text{also } 3A - B + 2C = 5$$

$$\text{and } -3C = 3 \Rightarrow C = -1$$

$$\therefore A + B = 1$$

$$3A - B = 7$$

$$\text{adding: } 4A = 8 \Rightarrow A = 2$$

$$\text{since } A + B = 1 \Rightarrow B = 1 - A = 1 - 2 = -1.$$

Exercise 1.7

1. (i) $3x - 2y = 4$

$$3x = 4 + 2y$$

$$x = \frac{4 + 2y}{3}$$

(ii) $2x - b = 4c$

$$2x = 4c + b$$

$$x = \frac{4c + b}{2}$$

(iii) $5x - 4 = \frac{y}{2}$

$$5x = \frac{y}{2} + 4$$

$$x = \frac{\frac{y}{2} + 4}{5} = \frac{y + 8}{10}$$

$$(iv) \quad 5(x - 3) = 2y$$

$$x - 3 = \frac{2y}{5}$$

$$x = \frac{2y}{5} + 3 = \frac{2y + 15}{5}$$

$$(v) \quad 3y = \frac{x}{3} - 2$$

$$\frac{-x}{3} = -3y - 2$$

$$\frac{x}{3} = 3y + 2$$

$$x = 9y + 6$$

$$(vi) \quad xy = xz + yz$$

$$xy - xz = yz$$

$$x(y - z) = yz$$

$$x = \frac{yz}{y - z}$$

$$2. (i) \quad 2x - \frac{y}{3} = \frac{1}{3}$$

$$2x = \frac{1}{3} + \frac{y}{3} = \frac{y + 1}{3}$$

$$x = \frac{y + 1}{6}$$

$$(ii) \quad z = \frac{y - 2x}{3}$$

$$3z = y - 2x$$

$$2x = y - 3z$$

$$x = \frac{y - 3z}{2}$$

$$(iii) \quad \frac{a}{x} - b = c$$

$$a - bx = cx$$

$$-cx - bx = -a$$

$$cx + bx = a$$

$$x(c + b) = a$$

$$x = \frac{a}{b + c}$$

$$3. (a) \quad V = \pi r^2 h$$

$$\Rightarrow \pi r^2 h = V$$

$$\Rightarrow r^2 = \frac{V}{\pi h}$$

$$r = \sqrt{\frac{V}{\pi h}}$$

$$(b) \quad A = 2\pi r h$$

$$\Rightarrow 2\pi r h = A$$

$$r = \frac{A}{2\pi h}$$

$$(c) \quad r = \sqrt{\frac{V}{\pi h}} \text{ and } r = \frac{A}{2\pi h}$$

$$\Rightarrow \sqrt{\frac{V}{\pi h}} = \frac{A}{2\pi h}$$

$$\begin{aligned}\therefore \frac{V}{\pi h} &= \frac{A^2}{4\pi^2 h^2} \\ \Rightarrow A^2 &= \frac{4\pi^2 h^2 V}{\pi h} = 4\pi h V\end{aligned}$$

4. (a) $A_{\text{circle}} = \pi r^2$

(b) The side of the square = $2r$

$$\Rightarrow A_{\text{square}} = l^2 = (2r)^2 = 4r^2$$

(c) $A_{\text{corners}} = 4r^2 - \pi r^2 = (4 - \pi)r^2$

(d) area of new square = $(4r)^2 = 16r^2$


$$\text{area of new circle} = \pi \left(\frac{r}{2}\right)^2 = \frac{\pi r^2}{4}$$

$$\begin{aligned}\Rightarrow \text{area of new shaded section} &= 16r^2 - \frac{\pi r^2}{4} \\ &= r^2 \left(16 - \frac{\pi}{4}\right) \\ &= \frac{r^2}{2} (64 - \pi)\end{aligned}$$

(e) To find the radius of the outer circle we need to find the distance from the centre of the circle to a corner (vertex) of the square.

$$d = \sqrt{r^2 + r^2} = \sqrt{2r^2} = \sqrt{2}r.$$

$$\begin{aligned}\Rightarrow \text{Area of outer circle} &= \pi d^2 \\ &= \pi (\sqrt{2}r)^2 \\ &= 2\pi r^2 \\ &= \text{twice the area of the inner circle.}\end{aligned}$$



5. (i) $f' = \frac{fc}{c-u}$

$$\Rightarrow f'(c-u) = fc$$

$$c-u = \frac{fc}{f'}$$

$$-u = \frac{fc}{f'} - c$$

$$u = c - \frac{fc}{f'} = \frac{f'c - fc}{f'} = \frac{c(f' - f)}{f'}$$

(ii) $f' = \frac{fc}{c-u}$

$$f'(c-u) = fc$$

$$f'c - f'u = fc$$

$$f'c - fc = f'u$$

$$c(f' - f) = f'u$$

$$c = \frac{f'u}{f' - f}$$

6. (i) $T = 2\pi \sqrt{\frac{l}{g}}$

$$T^2 = 4\pi^2 \frac{l}{g}$$

$$l = \frac{gT^2}{4\pi^2}$$

(ii) $T = 3, g = 10. \Rightarrow l = \frac{10.3^2}{4\pi^2} = \frac{90}{39.48} \simeq 2.3 \text{ m}$

7. (i) $\frac{x}{y} = \frac{a+b}{a-b}$

$$x(a-b) = y(a+b)$$

$$ax - bx = ay + by$$

$$ax - ay = bx + by$$

$$a(x-y) = b(x+y)$$

$$a = \frac{b(x+y)}{(x-y)}$$

(ii) $bc - ac = ac$

$$-ac - ac = -bc$$

$$-2ac = -bc$$

$$a = \frac{-bc}{-2c} = \frac{b}{2}$$

$$\begin{aligned}
 8. \quad (i) \quad y &= \frac{3(u-v)}{4} \\
 4y &= 3u - 3v \\
 3v &= 3u - 4y \\
 v &= \frac{3u - 4y}{3}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad s &= \frac{t}{2}(u+v) \\
 2s &= tu + tv \\
 -tv &= tu - 2s \\
 tv &= 2s - tu \\
 v &= \frac{2s - tu}{t}
 \end{aligned}$$


$$\begin{aligned}
 9. \quad A &= P\left(1 + \frac{i}{100}\right)^3 \\
 P\left(1 + \frac{i}{100}\right)^3 &= A \\
 \left(1 + \frac{i}{100}\right) &= \sqrt[3]{\frac{A}{P}} \\
 1 + \frac{i}{100} &= \sqrt[3]{\frac{A}{P}} \\
 100 + i &= 100\sqrt[3]{\frac{A}{P}} \\
 i &= 100\sqrt[3]{\frac{A}{P}} - 100 \\
 P &= 2500, \quad A = 2650 \\
 \therefore i &= 100\sqrt[3]{\frac{2650}{2500}} - 100 \\
 &= 100(1.0196) - 100 \\
 &= 1.961 \\
 \therefore i &= 2\%
 \end{aligned}$$

$$\begin{aligned}
 10. \quad (i) \quad d &= \sqrt{\frac{a-b}{ac}} \\
 d^2 &= \frac{a-b}{ac} \\
 acd^2 &= a-b \\
 c &= \frac{a-b}{ad^2}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad b &= \frac{2c-1}{c-1} \\
 \Rightarrow b(c-1) &= 2c-1 \\
 bc - b &= 2c-1 \\
 bc - 2c &= b-1 \\
 c(b-2) &= b-1 \\
 c &= \frac{b-1}{b-2}
 \end{aligned}$$

$$\begin{aligned}
 11. \quad (i) \quad \text{From Pythagoras: } h^2 + r^2 &= 15^2 \\
 h^2 &= 15^2 - r^2 \\
 h &= \sqrt{15^2 - r^2} \\
 (ii) \quad \text{at } r = 5 \text{ cm: } h &= \sqrt{15^2 - 5^2} \\
 &= \sqrt{225 - 25} = \sqrt{200} \\
 &= 10\sqrt{2} \text{ cm} \\
 (iii) \quad r = \frac{15}{2}; h &= \sqrt{15^2 - \left(\frac{15}{2}\right)^2} \\
 &= \sqrt{168.75} = 12.99 \text{ cm} \\
 h &= 13 \text{ cm}
 \end{aligned}$$

$$\begin{aligned}
 12. \quad (i) \quad 2W + L &= 300 \\
 L &= 300 - 2W \\
 (ii) \quad A &= L \times W \\
 &= (300 - 2W) \cdot W = 300W - 2W^2 \\
 (iii) \quad 10000 &= 300W - 2W^2 \\
 2W^2 - 300W + 10000 &= 0 \\
 W^2 - 150W + 5000 &= 0 \\
 (W - 50)(W - 100) &= 0
 \end{aligned}$$



$\Rightarrow W = 50$ or $W = 100$
 hence, $L = 300 - 2(50)$ or $L = 300 - 2(100)$
 $= 200$ $= 100$
 answer: $(50, 200)$ or $(100, 100)$

Exercise 1.8

1. (a) 4, 7, 10, 13, 16 ... constant 1st difference \Rightarrow linear
- (b) -2, 2, 6, 10, 14 ... constant 1st difference \Rightarrow linear
- (c) -4, -3, 0, 5, 12
 $+1, +3, +5, +7$
 $+2, +2, +2$ constant 2nd difference \Rightarrow quadratic
- (d) 2, 1, -2, -7, -14, -23...
 $-1, -3, -5, -7,$
 $-2, -2, -2$... constant 2nd difference \Rightarrow quadratic
- (e) 2, 7, 22, 47
 $+5, +15, +25$
 $+10, +10$ constant 2nd difference \Rightarrow quadratic
- (f) 3, 1, -5, -15, -29,
 $-2, -6, -10, -14$
 $-4, -4, -4, \dots$ constant 2nd difference \Rightarrow quadratic
- (g) 1, -4, -19, -44, -79
 $-5, -15, -25, -35$
 $-10, -10, -10$... constant 2nd difference \Rightarrow quadratic
- (h) 3, -2, -7, -12, -17...
 $-5, -5, -5, -5$... constant 1st difference \Rightarrow linear
- (i) 0, 3, 12, 27, 48
 $+3, +9, +15, +21$
 $+6, +6, +6$ constant 2nd difference \Rightarrow quadratic
- (j) 5, 17, 37, 65, 101
 $+12, +20, +28, +36$
 $+8, +8, +8$... constant 2nd difference \Rightarrow quadratic.

2. (a) -1, 3, 15, 35, 63
 first differences = +4, +12, +20, +28
 second differences = +8, +8, +8.
 \Rightarrow quadratic pattern of the form $ax^2 + bx + c$
 also, $2a = 8 \Rightarrow a = 4$.
 $\therefore 4x^2 + bx + c$
 let $x = 1 \Rightarrow 4(1)^2 + b(1) + c = -1$
 $\Rightarrow b + c = -1 - 4 = -5$
 let $x = 2 \Rightarrow 4(2)^2 + b(2) + c = 3$
 $\Rightarrow 2b + c = 3 - 16 = -13$
 using simultaneous equations: $2b + c = -13$
 $\begin{array}{r} b + c = -5 \\ \underline{b} = -8 \\ \Rightarrow -8 + c = -5 \\ \Rightarrow c = -5 + 8 = 3 \end{array}$
 $\therefore ax^2 + bx + c = 4x^2 - 8x + 3$ for $x = 1, 2, 3, \dots$
 Note also we could let $x = 0 \Rightarrow 4(0)^2 + b(0) + c = -1$
 $\Rightarrow c = -1$

$$\begin{aligned}\text{let } x = 1 &\Rightarrow 4(1)^2 + b(1) + c = 3 \\ &4 + b + c = 3 \\ &4 + b - 1 = 3 \\ &b = 0.\end{aligned}$$

$$\therefore ax^2 + bx + c = 4x^2 - 1 \quad \text{for } x = 0, 1, 2, \dots$$

(b) 4, 3, 0, -5, -12, -21, -32

first difference = -1, -3, -5, -7, -9, -11

second difference = -2, -2, -2, -2

\Rightarrow quadratic pattern of the form $ax^2 + bx + c$

also, $2a = -2 \Rightarrow a = -1$.

$\therefore -x^2 + bx + c$

$$\begin{aligned}\text{let } x = 0 &\Rightarrow -(0)^2 + b(0) + c = 4 \\ &c = 4\end{aligned}$$

$$\begin{aligned}\text{let } x = 1 &\Rightarrow -(1)^2 + b(1) + c = 3 \\ -1 + b + 4 &= 3 \\ b &= 0\end{aligned}$$

$\therefore ax^2 + bx + c = -x^2 + 4$ is the pattern for $x = 0, 1, 2, \dots$

3. (i) 2, 7, 12, 17, 22, ...

first difference = 5, a constant \Rightarrow a linear pattern.

$\therefore f(x) = ax + b = 5x + b$.

Let $x = 0$ be the 1st term of the pattern.

$\Rightarrow f(0) = 5(0) + b = 2$

$\therefore b = 2$

$\therefore f(x) = 5x + 2$ for $x = 0, 1, 2, \dots$

(ii) -6, -2, 2, 6, 10...

first difference = 4, a constant \Rightarrow a linear pattern.

$\therefore f(x) = ax + b = 4x + b$

Let $x = 0$ be the 1st term of the pattern.

$\Rightarrow f(0) = 4(0) + b = -6$

$\therefore b = -6$.

$\therefore f(x) = 4x - 6$ for $x = 0, 1, 2, \dots$

(iii) 3, 2, 1, 0, -1, -2, ...

first difference = -1, a constant \Rightarrow a linear relationship

$\therefore f(x) = ax + b = -x + b$

Let $x = 0$ be the 1st term of the pattern.

$\Rightarrow f(0) = -(0) + b = 3$

$\therefore b = 3$

$$\begin{aligned}\therefore f(x) &= -x + 3 \\ &= 3 - x.\end{aligned}$$

(iv) -2, 7, -12, -17, -22, ...

first difference = -5, a constant \Rightarrow a linear relationship

$\therefore f(x) = ax + b = -5x + b$

Let $x = 0$ be the first term of the pattern.

$\Rightarrow f(0) = -5(0) + b = -2$

$\therefore b = -2$

$\therefore f(x) = -5x - 2$

(v) 3, 3.5, 4, 4.5, 5, ...

first difference = 0.5, a constant \Rightarrow a linear relationship.

$\therefore f(x) = ax + b = 0.5x + b$

Let $x = 0$ be the first term of the pattern.

$$\therefore f(0) = 0.5(0) + b = 3$$

$$\Rightarrow b = 3$$

$$\therefore f(x) = 0.5(x) + 3$$

$$= \frac{x}{2} + 3 \text{ for } x = 0, 1, 2, \dots$$

(vi) $-1, -0.8, -0.6, -0.4, -0.2, \dots$

first difference = 0.2, a constant \Rightarrow a linear relationship.

$$\therefore f(x) = ax + b = 0.2x + b$$

Let $x = 0$ be the first term of the pattern.

$$\therefore f(0) = 0.2(0) + b = -1$$

$$\therefore b = -1$$

$$\therefore f(x) = 0.2x - 1 \text{ for } x = 0, 1, 2, \dots$$

4. 11, 13, 15, 17, 19, ...

first difference = 2, a constant \Rightarrow a linear relationship.

$$\therefore f(x) = ax + b = 2x + b$$

Let $x = 3$ be the first term of the pattern.

$$\Rightarrow f(3) = 2(3) + b = 11$$

$$\Rightarrow b = 5$$

$$\therefore f(x) = 2x + 5 \text{ for } x = 3, 4, 5, \dots$$

5. 1, 3, 5, 7, 9, ...

first difference = 2, a constant \Rightarrow a linear relationship.

$$\therefore f(x) = ax + b = 2x + b$$

Let $x = -2$ be the first term of the pattern.

$$\therefore f(-2) = 2(-2) + b = 1$$

$$b = 1 + 4 = 5.$$

$$\therefore f(x) = 2x + 5 \text{ for } x = -2, -1, 0, \dots$$

6. (a) 3, 6, 9, ...

a first difference = 3 (a constant) \Rightarrow a linear pattern.

$$\Rightarrow f(x) = ax + b = 3x + b$$

Let $x = 1$ be the first element of the pattern.

$$\therefore f(1) = 3(1) + b = 3$$

$$\Rightarrow b = 0.$$

$$\therefore f(x) = 3x \text{ for } x = 1, 2, 3, \dots$$

\Rightarrow for the 15th element, $x = 15$.

$$\therefore f(15) = 3(15) = 45 \text{ matchsticks are needed.}$$

(b) 4, 8, 12, ...

a first difference = 4 (a constant) \Rightarrow a linear pattern.

$$\Rightarrow f(x) = ax + b = 4x + b$$

Let $x = 1$ be the first element of the pattern.

$$\therefore f(1) = 4(1) + b = 4$$

$$\Rightarrow b = 0.$$

$$\Rightarrow \therefore f(x) = 4x \text{ for } x = 1, 2, 3, \dots$$

\Rightarrow for the 15th element, $x = 15$.

$$\therefore f(15) = 4(15) = 60 \text{ matchsticks are needed.}$$

(c) 3, 5, 7, ...

a first difference = 2 (constant) \Rightarrow a linear pattern.

$$\Rightarrow f(x) = ax + b = 2x + b$$

Let $x = 1$ be the first element of the pattern.

$$\therefore f(1) = 2(1) + b = 3$$

$$\Rightarrow b = 1$$

$$\therefore f(x) = 2x + 1 \text{ for } x = 1, 2, 3, \dots$$

\Rightarrow For the 15th element, $x = 15$.

$$\therefore f(15) = 2(15) + 1 = 31 \text{ matchsticks are needed.}$$

7. Plan A = $35x + 70$

Plan B = $24x + 125$

Both plans repay the same amount if

$$35x + 70 = 24x + 125$$

$$\Rightarrow 11x = 55$$

$$x = 5 \text{ months.}$$

8. 4, 7, 14, 25, 40

first difference: 3, 7, 11, 15

second difference : 4, 4, 4 \Rightarrow a quadratic pattern, $f(t) = at^2 + bt + c$

$$\therefore 2a = 4 \Rightarrow a = 2.$$

$$\therefore f(t) = 2t^2 + bt + c$$

At $t = 0$, $f(0) = 2(0)^2 + b(0) + c = 4$ (i.e. at the start there were 4 bacteria)

$$\Rightarrow c = 4$$

At $t = 1$, $f(1) = 2(1)^2 + b(1) + c = 7$

$$\Rightarrow 2 + b + 4 = 7$$

$$\Rightarrow b = 1$$

$$\therefore f(t) = 2t^2 + t + 4 \text{ for } t = 0, 1, 2, 3, \dots$$

when is $f(t) = 529$, assuming $f(t) = 2t^2 + t + 4$?

if $t = 10 \Rightarrow 2(10)^2 + 10 + 4 = 214$ too small

$t = 15 \Rightarrow 2(15)^2 + 15 + 4 = 469$ too small

$t = 16 \Rightarrow 2(16)^2 + 16 + 4 = 532$ too large

\therefore In the 16th hour, the number of bacteria was 529.

Exercise 1.9

1. (i) $y = 2x^2 + 2x - 1$ is not linear because the highest power is not 1.

(ii) $y = 2(x - 1)^{-1}$ is not linear because the highest power of x is not 1.

(iii) $y^2 = 3x + 4$

$$\Rightarrow y = \sqrt{3x + 4} \text{ is not linear because the highest power of } x \text{ is not } 1.$$

$$= (3x + 4)^{\frac{1}{2}}$$

2. (i) Solve $5x - 3 = 32$

$$\Rightarrow 5x = 35$$

$$x = \frac{35}{5} = 7$$

(ii) Solve $3x + 2 = x + 8$

$$\Rightarrow 3x - x = 8 - 2$$

$$2x = 6$$

$$x = 3$$

(iii) Solve $2 - 5x = 8 - 3x$

$$\Rightarrow 3x - 5x = 8 - 2$$

$$-2x = 6$$

$$x = \frac{6}{-2} = -3$$

$$\begin{aligned}
 \text{3. (i) Solve } 2(x - 3) + 5(x - 1) &= 3 \\
 \Rightarrow 2x - 6 + 5x - 5 &= 3 \\
 \Rightarrow 7x = 3 + 11 &= 14 \\
 x &= 2
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) Solve } 2(4x - 1) - 3(x - 2) &= 14 \\
 \Rightarrow 8x - 2 - 3x + 6 &= 14 \\
 5x = 14 - 4 &= 10 \\
 x &= 2
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) Solve } 3(x - 1) - 4(x - 2) &= 6(2x + 3) \\
 \Rightarrow 3x - 3 - 4x + 8 &= 12x + 18 \\
 -x - 12x &= 18 - 5 \\
 -13x &= 13 \\
 x &= -1
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) Solve } 3(x + 5) + 2(x + 1) - 3x &= 22 \\
 3x + 15 + 2x + 2 - 3x &= 22 \\
 2x &= 22 - 17 \\
 2x &= 5 \\
 x &= \frac{5}{2} = 2.5
 \end{aligned}$$

$$\begin{aligned}
 \text{4. (i) } \frac{2x + 1}{5} &= 1 \\
 \Rightarrow 2x + 1 &= 5 \\
 2x &= 4 \\
 x &= 2
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \frac{3x - 1}{4} &= 8 \\
 \Rightarrow 3x - 1 &= 32 \\
 3x &= 33 \\
 x &= 11
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } \frac{x - 3}{4} &= \frac{x - 2}{5} \\
 \Rightarrow 5(x - 3) &= 4(x - 2) \\
 5x - 15 &= 4x - 8 \\
 5x - 4x &= 15 - 8 \\
 x &= 7
 \end{aligned}$$

$$\begin{aligned}
 \text{5. (i) } \frac{2a}{3} - \frac{a}{4} &= \frac{5}{6} \\
 \Rightarrow \text{multiplying each term by } 12 : 4(2a) - 3(a) &= 2(5) \\
 \Rightarrow 8a - 3a &= 10 \\
 5a &= 10 \\
 a &= 2
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \frac{b + 2}{4} - \frac{b - 3}{3} &= \frac{1}{2} \\
 \Rightarrow \text{multiplying each term by } 12 : 3(b + 2) - 4(b - 3) &= 6 \\
 \Rightarrow 3b + 6 - 4b + 12 &= 6 \\
 -b &= 6 - 18 \\
 -b &= -12 \\
 b &= 12
 \end{aligned}$$

$$(iii) \frac{3c-1}{6} - \frac{c-3}{4} = \frac{4}{3}$$

\Rightarrow multiplying each term by 12 : $2(3c-1) - 3(c-3) = 4(4)$

$$\Rightarrow 6c - 2 - 3c + 9 = 16$$

$$3c = 16 - 7$$

$$3c = 9$$

$$c = 3$$

$$6. (i) \frac{x-2}{5} + \frac{2x-3}{10} = \frac{1}{2}$$

multiplying each term by 10 we get : $2(x-2) + (2x-3) = 5(1)$

$$\Rightarrow 2x - 4 + 2x - 3 = 5$$

$$4x = 5 + 7$$

$$4x = 12$$

$$x = 3$$

$$(ii) \frac{3y-12}{5} + 3 = \frac{3(y-5)}{2}$$

multiplying each term by 10 we get : $2(3y-12) + 10(3) = 5.3(y-5)$

$$\Rightarrow 6y - 24 + 30 = 15y - 75$$

$$\therefore 6y - 15y = 24 - 30 - 75$$

$$-9y = -81$$

$$y = \frac{-81}{-9} = +9$$

$$(iii) \frac{3p-2}{6} - \frac{3p+1}{4} = \frac{2}{3}$$

multiplying each term by 12 we get : $2(3p-2) - 3(3p+1) = 4(2)$

$$\Rightarrow 6p - 4 - 9p - 3 = 8$$

$$-3p = 8 + 7$$

$$-3p = 15$$

$$p = -5$$

$$(iv) \frac{3r-2}{5} - \frac{2r-3}{4} = \frac{1}{2}$$

multiplying each term by 20 we get : $4(3r-2) - 5(2r-3) = 10(1)$

$$\Rightarrow 12r - 8 - 10r + 15 = 10$$

$$2r = 10 - 7$$

$$2r = 3$$

$$r = \frac{3}{2} = 1.5$$

$$7. (i) \frac{3}{4}(2x-1) - \frac{2}{3}(4-x) = 2$$

multiplying each term by 12 we get : $3.3(2x-1) - 4.2(4-x) = 12.2$

$$\Rightarrow 18x - 9 - 32 + 8x = 24$$

$$26x = 24 + 41$$

$$26x = 65$$

$$x = \frac{65}{26} = 2.5$$

$$(ii) \frac{2}{3}(x-1) - \frac{1}{5}(x-3) = x+1$$

multiplying each term by 15 we get : $5.2(x-1) - 3.1(x-3) = 15x + 15$

$$\Rightarrow 10x - 10 - 3x + 9 = 15x + 15$$

$$7x - 15x = 15 + 1$$

$$-8x = 16$$

$$x = -2.$$

Exercise 1.10

$$\begin{array}{rcl}
 \text{1. (i)} & 3x - 2y = 8 & \Rightarrow & 3x - 2y = 8 \\
 & x + y = 6 & & \underline{2x + 2y = 12} \\
 & & \text{(adding)} & 5x = 20 \\
 & & & x = 4
 \end{array}$$

$$\text{since } x + y = 6 \Rightarrow 4 + y = 6$$

$$\therefore y = 2.$$

$$\therefore \text{ solution } (x, y) = (4, 2)$$

$$\begin{array}{rcl}
 \text{(ii)} & 3x - y = 1 & \Rightarrow & 6x - 2y = 2 \\
 & x - 2y = -8 & & \underline{x - 2y = -8} \\
 & & \text{(subtracting)} & 5x = 10 \\
 & & & x = 2
 \end{array}$$

$$\text{since } x - 2y = -8 \Rightarrow 2 - 2y = -8$$

$$-2y = -10$$

$$y = 5$$

$$\therefore \text{ solution } (x, y) = (2, 5)$$

$$\begin{array}{rcl}
 \text{(iii)} & 2x - 5y = 1 & \Rightarrow & 4x - 10y = 2 \\
 & 4x - 3y - 9 = 0 & & \underline{4x - 3y = 9} \\
 & & \text{(subtracting)} & -7y = -7 \\
 & & & y = 1
 \end{array}$$

$$\text{since } 2x - 5y = 1 \Rightarrow 2x - 5(1) = 1$$

$$\Rightarrow 2x = 6$$

$$x = 3$$

$$\therefore \text{ solution } (x, y) = (3, 1)$$

$$\begin{array}{rcl}
 \text{2. (i)} & 4x - 5y = 22 & \Rightarrow & 12x - 15y = 66 \\
 & 7x + 3y - 15 = 0 & & \underline{35x + 15y = 75} \\
 & & \text{(adding)} & 47x = 141
 \end{array}$$

$$x = \frac{141}{47} = 3$$

$$\text{since } 4x - 5y = 22 \Rightarrow 4(3) - 5y = 22$$

$$-5y = 22 - 12$$

$$-5y = 10$$

$$y = -2$$

$$\therefore \text{ solution } (x, y) = (3, -2)$$

$$\begin{array}{rcl}
 \text{(ii)} & \frac{x}{2} - \frac{y}{6} = \frac{1}{6} & \Rightarrow & 3x - y = 1 \\
 & x - 2y = -8 & & \underline{x - 2y = -8} \\
 & & \Rightarrow & \underline{6x - 2y = 2} \\
 & & & \underline{x - 2y = -8}
 \end{array}$$

$$\text{(subtracting)} \quad 5x = 10$$

$$x = 2$$

$$\text{since } 3x - y = 1 \Rightarrow 3(2) - y = 1$$

$$6 - y = 1$$

$$-y = -5$$

$$y = 5$$

$$\therefore \text{ solution } (x, y) = (2, 5)$$

$$\begin{aligned}
 \text{(iii)} \quad \frac{4x-2}{5} &= \frac{8y}{10} & \Rightarrow & \quad 8x-4=8y \\
 18x-20y &= 4 & \Rightarrow & \quad 18x-20y=4 \\
 & & \Rightarrow & \quad 8x-8y=4 \\
 & & & \quad 18x-20y=4 \\
 & & \Rightarrow & \quad 40x-40y=20 \\
 & & & \quad 36x-40y=8 \\
 & & \text{(subtracting)} & \quad 4x=12 \\
 & & & \quad x=3
 \end{aligned}$$

$$\begin{aligned}
 \text{since } 18x-20y &= 4 \\
 \Rightarrow 18(3)-20y &= 4 \\
 -20y &= 4-54 = -50 \\
 y &= 2\frac{1}{2} \\
 \therefore \text{solution } (x, y) &= (3, 2\frac{1}{2})
 \end{aligned}$$

$$\begin{aligned}
 \text{3. } \frac{2x-5}{3} + \frac{y}{5} &= 6 \Rightarrow 5(2x-5) + 3y = 15.6 \\
 &\Rightarrow 10x-25+3y=90 \\
 10x+3y &= 115 \\
 \frac{3x}{10} + 2 &= \frac{3y-5}{2} \Rightarrow 3x+10.2=5(3y-5) \\
 &\Rightarrow 3x+20=15y-25 \\
 &\Rightarrow 3x-15y=-45 \\
 \therefore 10x+3y &= 115 & \Rightarrow & \quad 50x+15y=575 \\
 3x-15y &= -45 & \Rightarrow & \quad 3x-15y=-45 \\
 & & \text{(adding)} & \quad 53x=530 \\
 & & & \quad x=10
 \end{aligned}$$

$$\begin{aligned}
 \text{since } 10x+3y &= 115 \\
 10(10)+3y &= 115 \\
 3y &= 15 \\
 y &= 5
 \end{aligned}$$

$$\therefore \text{solution } (x, y) = (10, 5)$$

$$\begin{aligned}
 \text{4. } y &= 3x-23 \Rightarrow y=3x-23 \\
 y &= \frac{x}{2} + 2 \Rightarrow 2y=x+4 \\
 &\Rightarrow 2y=6x-46 \\
 &\quad 2y=x+4 \\
 &\quad \underline{\hspace{1cm}} \\
 \text{(subtracting)} \quad 0 &= 5x-50 \\
 \Rightarrow 5x &= 50 \\
 x &= 10
 \end{aligned}$$

$$\begin{aligned}
 \text{since } y &= 3x-23 \\
 \Rightarrow y &= 3(10)-23=7 \\
 \therefore \text{solution } (x, y) &= (10, 7)
 \end{aligned}$$

5. (i) $A: 2x + y + z = 8$ $3A: 6x + \cancel{3y} + 3z = 24$
 $B: 5x - 3y + 2z = 3$ $B: \underline{5x - \cancel{3y} + 2z = 3}$
 $C: 7x + y + 3z = 20$ $D: (\text{adding}) \quad 11x \quad + 5z = 27$
also $B: 5x - \cancel{3y} + 2z = 3$
 $3C: \underline{21x + \cancel{3y} + 9z = 60}$
 $E: (\text{adding}) \quad 26x \quad + 11z = 63$
 $\Rightarrow 11D: 121x + \cancel{55z} = 297$
 $5E: \underline{130x + \cancel{55z} = 315}$
 $(\text{subtracting}) \quad -9x = -18$
 $x = 2$
since $11x + 5z = 27$
 $11(2) + 5z = 27$
 $5z = 27 - 22 = 5$
 $z = 1$
since $2x + y + z = 8$
 $2(2) + y + 1 = 8$
 $y = 3$
 \therefore solution $(x, y, z) = (2, 3, 1)$
- (ii) $A: 2x - y - z = 6$ $2A: 4x - \cancel{2y} - 2z = 12$
 $B: 3x + 2y + 3z = 3 \Rightarrow B: \underline{3x + \cancel{2y} + 3z = 3}$
 $C: 4x + y - 2z = 3$ $D: (\text{adding}) \quad 7x \quad + z = 15$
also $B: 3x + \cancel{2y} + 3z = 3$
 $2C: \underline{8x + \cancel{2y} - 4z = 6}$
 $E: (\text{subtracting}) \quad -5x \quad + 7z = -3$
since $D: 7x + z = 15 \Rightarrow 7D = 49x + \cancel{7z} = 105$
also $E = \underline{-5x + \cancel{7z} = -3}$
subtracting: $54x = 108$
 $x = 2.$
since $D: 7x + z = 15$
 $\Rightarrow 7(2) + z = 15$
 $z = 1$
also, since $A: 2x - y - z = 6$
 $\Rightarrow 2(2) - y - 1 = 6$
 $-y = 3$
 $y = -3.$
 \therefore solution $(x, y, z) = (2, -3, 1)$
- (iii) $A: 2x + y - z = 9$ $A: 2x + y - \cancel{z} = 9$
 $B: x + 2y + z = 6 \Rightarrow B: \underline{x + 2y + \cancel{z} = 6}$
 $C: 3x - y + 2z = 17$ $D (\text{adding}): 3x + 3y = 15$
also $2B: 2x + 4y + \cancel{2z} = 12$
 $C: \underline{3x - y + \cancel{2z} = 17}$
 $E (\text{subtracting}): -x + 5y = -5$

$$\text{since } D: 3x + 3y = 15$$

$$\underline{3E: -3x + 15y = -15}$$

$$\text{adding: } 18y = 0$$

$$y = 0.$$

$$\text{since } E: -x + 5y = -5$$

$$-x + 5(0) = -5$$

$$x = 5.$$

$$\text{also, } A: 2x + y - z = 9$$

$$2(5) + 0 - z = 9$$

$$-z = 9 - 10 = -1$$

$$+z = 1$$

$$\therefore \text{ solution } (x, y, z) = (5, 0, 1)$$

$$6. \quad (i) \quad A: 2a + b + c = 8 \quad \Rightarrow \quad 3A: 6a + \cancel{3b} + 3c = 24$$

$$B: 5a - 3b + 2c = 3 \quad B: \underline{5a - \cancel{3b} + 2c = -3}$$

$$C: 7a - 3b + 3c = 1 \quad D: \text{adding: } 11a + 5c = 21$$

$$\text{also } B: 5a - \cancel{3b} + 2c = -3$$

$$C: \underline{7a - \cancel{3b} + 3c = 1}$$

$$E: \text{subtracting: } -2a - c = -4.$$

$$\text{since } D: 11a + \cancel{5c} = 21$$

$$\text{and } 5E: \underline{-10a - \cancel{5c} = -20}$$

$$\text{adding: } a = 1$$

$$\text{since } D: 11a + 5c = 21$$

$$\Rightarrow 11(1) + 5c = 21$$

$$5c = 10$$

$$c = 2.$$

$$\text{also, } A: 2a + b + c = 8$$

$$2(1) + b + 2 = 8$$

$$b = 4$$

$$\therefore \text{ solution } (a, b, c) = (1, 4, 2)$$

$$(ii) \quad A: x + y + 2z = 3 \quad \Rightarrow \quad 2A: 2x + \cancel{2y} + 4z = 6$$

$$B: 4x + 2y + z = 13 \quad B: \underline{4x + \cancel{2y} + z = 13}$$

$$C: 2x + y - 2z = 9 \quad D: (\text{subtracting}): -2x + 3z = -7$$

$$\text{also } B: \cancel{4x} + \cancel{2y} + z = 13$$

$$2C: \underline{\cancel{4x} + \cancel{2y} - 4z = 18}$$

$$E: (\text{subtracting}): 5z = -5$$

$$\Rightarrow z = -1$$

$$\text{since } D: -2x + 3z = -7$$

$$\Rightarrow -2x + 3(-1) = -7$$

$$-2x - 3 = -7$$

$$-2x = -7 + 3$$

$$-2x = -4$$

$$x = 2.$$

$$\text{also, A: } x + y + 2z = 3$$

$$2 + y + 2(-1) = 3$$

$$y = 3$$

$$\therefore \text{ solution } (x, y, z) = (2, 3, -1).$$

$$(iii) \text{ A: } x + y + z = 2 \qquad \text{A: } x + y + z = 2$$

$$\text{B: } 2x + 3y + z = 7$$

$$\text{C: } \frac{x}{2} + \frac{y}{2} + \frac{z}{3} = \frac{2}{3} \Rightarrow \text{C: } \underline{3x - y + 2z = 4}$$

$$\text{D: (adding) } 4x \quad + 3z = 6$$

$$\text{also B: } 2x + 3y + z = 7$$

$$\underline{3\text{C: } 9x - 3y + 6z = 12}$$

$$\text{E: (adding) } 11x \quad + 7z = 19$$

$$\therefore 11\text{D: } 44x + 33z = 66$$

$$\underline{4\text{E: } 44x + 28z = 76}$$

$$\text{subtracting: } 5z = -10$$

$$z = -2.$$

$$\text{since D: } 4x + 3z = 6$$

$$4x + 3(-2) = 6$$

$$4x = 12$$

$$x = 3$$

$$\text{also, A: } x + y + z = 2$$

$$\Rightarrow 3 + y - 2 = 2$$

$$y = 2 - 1$$

$$y = 1$$

$$\therefore \text{ solution } (x, y, z) = (3, 1, -2)$$

$$7. \text{ A: } 6x + 4y - 2z - 5 = 0 \Rightarrow \text{A: } 6x + \cancel{4y} - 2z = 5$$

$$\text{B: } 3x - 2y + 4z + 10 = 0 \qquad \underline{2\text{B: } 6x - \cancel{4y} + 8z = -20}$$

$$\text{C: } 5x - 2y + 6z + 13 = 0 \quad \text{D: (adding): } 12x \quad + 6z = -15$$

$$\text{also B: } 3x - \cancel{2y} + 4z = -10$$

$$\underline{\text{C: } 5x - \cancel{2y} + 6z = -13}$$

$$\text{E (subtracting): } -2x \quad - 2z = 3$$

$$\text{also, D: } 12x + \cancel{6z} = -15$$

$$\underline{3\text{E: } -6x - \cancel{6z} = 9}$$

$$\text{adding: } 6x \quad = -6$$

$$x = -1$$

$$\text{since D: } 12x + 6z = -15$$

$$12(-1) + 6z = -15$$

$$6z = -3$$

$$z = -\frac{1}{2}$$

$$\text{also, A: } 6x + 4y - 2z = 5$$

$$\Rightarrow 6(-1) + 4y - 2\left(-\frac{1}{2}\right) = 5$$

$$-6 + 4y + 1 = 5$$

$$4y = 10$$

$$y = 2\frac{1}{2}$$

$$\therefore \text{solution } (x, y, z) = (-1, 2\frac{1}{2}, -\frac{1}{2}).$$

8. Curve $f(x) = ax^2 + bx + c$

$$(1, 2) \text{ on curve} \Rightarrow \text{when } x = 1, f(x) = 2$$

$$\Rightarrow 2 = a(1)^2 + b(1) + c$$

$$\Rightarrow 2 = a + b + c \quad : A$$

$$(2, 4) \text{ on curve} \Rightarrow \text{when } x = 2, f(x) = 4$$

$$\Rightarrow 4 = a(2)^2 + b(2) + c$$

$$\Rightarrow 4 = 4a + 2b + c \quad : B$$

$$(3, 8) \text{ on curve} \Rightarrow \text{when } x = 3, f(x) = 8$$

$$\Rightarrow 8 = a(3)^2 + b(3) + c$$

$$\Rightarrow 8 = 9a + 3b + c \quad : C$$

$$\text{since } A: \quad a + b + c = 2$$

$$\text{and } B: \quad \underline{4a + 2b + c = 4}$$

$$D(\text{subtracting}): \quad -3a - b = -2$$

$$\text{since } B: 4a + 2b + c = 4$$

$$\text{and } C: \quad \underline{a + 3b + c = 8}$$

$$E(\text{subtracting}): \quad -5a - b = -4$$

$$\text{also } D: \quad -3a - b = -2$$

$$E: \quad \underline{-5a - b = -4}$$

$$\text{subtracting:} \quad 2a = 2$$

$$\Rightarrow \quad a = 1$$

$$\text{since } D: \quad -3a - b = -2$$

$$\Rightarrow \quad -3(1) - b = -2$$

$$-b = +1$$

$$b = -1$$

$$\text{also } A: a + b + c = 2$$

$$\Rightarrow \quad 1 - 1 + c = 2$$

$$c = 2$$

$$\therefore \text{solution } (a, b, c) = (1, -1, 2)$$

9. Point 1, (1, 1)

$$\text{point 2, } (0, -6)$$

$$\text{point 3, } (-2, -8)$$

$$\text{Curve} = f(x) = ax^2 + bx + c$$

$$(1, 1) \text{ on curve} \Rightarrow \text{when } x = 1, f(x) = 1$$

$$\Rightarrow 1 = a(1)^2 + b(1) + c$$

$$\Rightarrow 1 = a + b + c \quad : A$$

$$(0, -6) \text{ on curve} \Rightarrow \text{when } x = 0, f(x) = -6$$

$$\Rightarrow -6 = a(0) + b(0) + c$$

$$\Rightarrow -6 = c \quad : B$$

$$(-2, -8) \text{ on curve} \Rightarrow \text{when } x = -2, f(x) = -8$$

$$\Rightarrow -8 = a(-2)^2 + b(-2) + c$$

$$\begin{aligned}
 & -8 = 4a - 2b + c && : C \\
 \text{also } A: & a + b - 6 = 1 \\
 C: & 4a - 2b - 6 = -8 \\
 \Rightarrow 2A: & 2a + 2b - 12 = 2 \\
 \text{adding:} & \underline{6a - 18 = -6} \\
 & 6a = 12 \\
 & a = 2 \\
 \text{since } A: & a + b - 6 = 1 \\
 \Rightarrow & 2 + b - 6 = 1 \\
 & b = 1 + 4 = 5 \\
 (a, b, c) = & (2, 5, -6) \\
 \text{solution: } f(x) = & 2x^2 + 5x - 6
 \end{aligned}$$

- 10.** Let x = number of people paying €20

Let y = number of people paying €30

$$\therefore x + y = 44\,000 : A$$

$$\text{also, } 20x + 30y = 1\,200\,000 : B$$

$$\therefore 20A: \underline{20x + 20y = 880\,000}$$

$$B: \underline{20x + 30y = 1\,200\,000}$$

$$\begin{aligned}
 \text{subtracting:} & -10y = -320\,000 \\
 & y = 32\,000
 \end{aligned}$$

\therefore 32 000 paid the higher price.

- 11.** Let x be Lydia's age now.

\therefore five years ago, Lydia was $(x - 5)$ years old.

Let Callum be y years old now.

\therefore three years from now, Callum will be $(y + 3)$ years old

$$\Rightarrow (y + 3) = 2(x - 5) : A$$

$$\text{also, } \frac{x + y}{2} = 16$$

$$\Rightarrow x + y = 32 : B$$

$$\text{From } A: y + 3 = 2x - 10$$

$$13 = 2x - y$$

$$\therefore A: 2x - y = 13$$

$$B: \underline{x + y = 32}$$

$$\begin{aligned}
 \text{adding: } 3x &= 45 \\
 x &= 15
 \end{aligned}$$

$$\text{since } x + y = 32$$

$$\Rightarrow 15 + y = 32$$

$$y = 17$$

Lydia is 15 years old, Callum is 17 years old.

- 12.** Equation of line: $y = ax + b$.

$(6, 7)$ on line \Rightarrow when $x = 6, y = 7$

$$\Rightarrow 7 = a(6) + b$$

$$\Rightarrow 6a + b = 7 : A$$

also, $(-2, 3)$ on line \Rightarrow when $x = -2, y = 3$

$$\Rightarrow 3 = a(-2) + b$$

$$\begin{array}{rcl} & \Rightarrow -2a + b = 3 & :B \\ \text{since } A: & 6a + \cancel{b} = 7 & \\ \text{and } B: & -2a + \cancel{b} = 3 & \\ \text{subtracting:} & 8a & = 4 \\ \Rightarrow & a & = \frac{4}{8} = \frac{1}{2}. \end{array}$$

$$\text{since } A: 6a + b = 7$$

$$\Rightarrow 6\left(\frac{1}{2}\right) + b = 7$$

$$b = 4$$

$$\therefore \text{line } y = \frac{1}{2}x + 4.$$

$$\text{Verify } (4, 6) \text{ is on line } \Rightarrow 6 = \frac{1}{2}(4) + 4$$

$$= 6, \text{ which is true.}$$

$$13. \quad \frac{N_1}{4} - N_2 = 0 \Rightarrow N_1 - 4N_2 = 0 \quad :A$$

$$N_1 + \frac{1}{2}N_2 - 99 = 0 \Rightarrow 2N_1 + N_2 = 198 \quad :B$$

$$\text{since } A: N_1 - 4N_2 = 0$$

$$\text{and } 4B: \frac{8N_1 + 4N_2 = 792}{}$$

$$\text{adding: } 9N_1 = 792$$

$$N_1 = 88$$

$$\text{also, } A: N_1 - 4N_2 = 0$$

$$\Rightarrow 88 - 4N_2 = 0$$

$$-4N_2 = -88$$

$$N_2 = 22$$

$$(N_1, N_2) = (88, 22)$$

$$\begin{aligned} 14. \quad \frac{a}{x-2} + \frac{b}{x+2} &= \frac{4}{(x-2)(x+2)} \\ \Rightarrow \frac{a(x+2) + b(x-2)}{(x-2)(x+2)} &= \frac{4}{(x-2)(x+2)} \\ \Rightarrow ax + 2a + bx - 2b &= 4 \end{aligned}$$

$$(a+b)x + 2a - 2b = 4 + 0x$$

$$\Rightarrow a + b = 0 \quad :A$$

$$\text{and } 2a - 2b = 4 \quad :B$$

$$\text{also } \frac{2a + 2b = 0}{\quad} \quad :2A$$

$$\text{adding: } 4a = 4$$

$$a = 1$$

$$\text{since } a + b = 0$$

$$\Rightarrow 1 + b = 0$$

$$b = -1$$

$$\therefore (a, b) = (1, -1)$$

$$\therefore \frac{1}{x-2} + \frac{-1}{x+2} = \frac{4}{(x-2)(x+2)}$$

$$\frac{x+2 - (x-2)}{(x-2)(x+2)} = \frac{4}{(x-2)(x+2)}$$

$$= \frac{\cancel{x} + 2 - \cancel{x} + 2}{(x-2)(x+2)}$$

$$\frac{4}{(x-2)(x+2)} = \frac{4}{(x-2)(x+2)} \quad \text{qed}$$

$$\begin{aligned}
 15. \quad & \frac{c}{z-3} + \frac{d}{z+2} = \frac{4}{(z-3)(z+2)} \\
 \Rightarrow & \frac{c(z+2) + d(z-3)}{(z-3)(z+2)} = \frac{4}{(z-3)(z+2)} \\
 \Rightarrow & cz + 2c + dz - 3d = 4 \\
 (c+d)z + 2c - 3d &= 0z + 4 \\
 \therefore & \quad c + d = 0 \quad : A \\
 \text{and} \quad & 2c - 3d = 4 \quad : B \\
 \Rightarrow & \quad 2c + 2d = 0 \quad : 2A \\
 \text{subtracting:} \quad & -5d = 4
 \end{aligned}$$

$$d = \frac{-4}{5}$$

$$\text{since } c + d = 0$$

$$\Rightarrow c - \frac{4}{5} = 0$$

$$\Rightarrow c = \frac{4}{5}$$

$$\therefore (c, d) = \left(\frac{4}{5}, -\frac{4}{5}\right)$$

$$\therefore \frac{4}{5(z-3)} + \frac{-4}{5(z+2)} = \frac{4}{(z-3)(z+2)}$$

$$\Rightarrow \frac{4(z+2) - 4(z-3)}{5(z-3)(z+2)}$$

$$= \frac{\cancel{4z} + 8 - \cancel{4z} + 12}{5(z-3)(z+2)}$$

$$= \frac{20}{5(z-3)(z+2)}$$

$$= \frac{4}{(z-3)(z+2)} = \frac{4}{(z-3)(z+2)} \quad \text{qed}$$

16. Let x = number of litres of 70% alcohol

$$\Rightarrow x(70\%) + 50(40\%) = (x + 50)(50\%)$$

$$\Rightarrow 0.7x + 0.4(50) = 0.5x + 0.5(50)$$

$$0.7x + 20 = 0.5x + 25$$

$$0.7x - 0.5x = 25 - 20$$

$$0.2x = 5$$

$$x = 25 \text{ litres.}$$

17. x = bigger number

y = smaller number

$$A: \quad x + y = 26$$

$$B: \quad 4x - 5y = 5$$

$$\Rightarrow 5A: \quad 5x + 5y = 130$$

$$B: \quad 4x - 5y = 5$$

$$\text{adding: } 9x = 135$$

$$x = 15$$

$$\text{since } x + y = 26$$

$$\Rightarrow 15 + y = 26$$

$$y = 11$$

$$\therefore (x, y) = (15, 11)$$

18. $v = u + at$ $v = \text{speed}$
 $t = \text{time.}$

$$\text{at } t = 7, v = 2 \quad \Rightarrow \quad 2 = u + 7a \quad :A$$

$$\text{at } t = 13, v = 5 \quad \Rightarrow \quad 5 = u + 13a \quad :B$$

$$\text{since } A: u + 7a = 2$$

$$B: u + 13a = 5$$

$$\text{subtracting: } -6a = -3$$

$$6a = 3$$

$$a = \frac{3}{6} = \frac{1}{2}$$

$$\text{since } u + 7a = 2$$

$$\Rightarrow u + 7\left(\frac{1}{2}\right) = 2$$

$$u = 2 - 3\frac{1}{2}$$

$$u = -1\frac{1}{2}$$

19. $\therefore 4x + 2y = 60 \quad :A$

$$\text{also, } 2x + 4y = 42 \quad :B$$

$$\therefore 4x + 2y = 60 \quad :A$$

$$\text{also } 4x + 8y = 84 \quad :2B$$

$$\text{subtracting: } -6y = -24$$


$$y = 4$$

$$\text{since } 4x + 2y = 60$$

$$\Rightarrow 4x + 2(4) = 60$$

$$4x = 52$$

$$x = 13$$



\therefore Original pen dimensions: 26×4

new pen dimensions: 13×8

Pen 1 requires $4x$ and $2y$ lengths

Pen 2 requires $2x$ and $4y$ lengths

\Rightarrow they both have $2x$ and $2y$ length in common

\therefore in pen 2, $2x$ lengths are swapped for $2y$ lengths

if $y < x$, less fencing is needed in pen 2.

Note: Area of pen 1 = $(2x) \times (y) = 2xy$

Area of pen 2 = $(x) \times (2y) = 2xy$

\Rightarrow the areas are the same.

20. $y = ax^2 + bx + c$

the point $(0, 1) \in \text{curve} \Rightarrow \text{when } x = 0, y = 1$

$$\Rightarrow 1 = a(0)^2 + b(0) + c$$

$$\Rightarrow 1 = c$$

the point $(2, 9) \in \text{curve} \Rightarrow \text{when } x = 2, y = 9$

$$\Rightarrow 9 = a(2)^2 + b(2) + c$$

$$\Rightarrow 9 = 4a + 2b + c$$

$$\Rightarrow 9 = 4a + 2b + 1 \quad \text{since } c = 1$$

$$\Rightarrow 8 = 4a + 2b$$

$$A: \Rightarrow 4 = 2a + b \quad \text{dividing each term by 2.}$$

the point $(4, 41) \in \text{curve} \Rightarrow \text{when } x = 4, y = 41$

$$\Rightarrow 41 = a(4)^2 + b(4) + c$$

$$\Rightarrow 41 = 16a + 4b + c$$

$$\Rightarrow 41 = 16a + 4b + 1 \quad \text{since } c = 1$$

$$\Rightarrow 40 = 16a + 4b$$

$$B: \Rightarrow 10 = 4a + b \quad \text{dividing each term by 4.}$$

$$\text{since } A: 2a + \cancel{b} = 4$$

$$\text{and } B: \underline{4a + \cancel{b} = 10}$$

$$\text{subtracting} \quad -2a = -6$$

$$a = 3$$

$$\text{since } A: 2a + b = 4$$

$$\Rightarrow \underline{2(3) + b = 4}$$

$$b = -2$$

$$\therefore (a, b, c) = (3, -2, 1)$$

21. (i) A: $y - z = 3$

$$B: x - 2y + z = -4$$

$$C: x + 2y = 11$$

$$\therefore A: \quad y - \cancel{z} = 3$$

$$B: \underline{x - 2y + \cancel{z} = -4}$$

$$D(\text{adding}): \quad \cancel{x} - y = -1$$

$$C: \underline{\cancel{x} + 2y = 11}$$

$$\text{subtracting:} \quad -3y = -12$$

$$y = 4$$

$$\text{since } x - y = -1$$

$$\Rightarrow x - 4 = -1$$

$$x = 3$$

$$\text{also } y - z = 3$$

$$\Rightarrow 4 - z = 3$$

$$-z = -1$$

$$z = 1$$

$$\text{solution } (x, y, z) = (3, 4, 1)$$

(ii) A: $\frac{x}{3} + \frac{y}{2} - z = 7 \Rightarrow 2x + 3y - 6z = 42$

$$B: \frac{x}{4} - \frac{3y}{2} + \frac{z}{2} = -6 \Rightarrow x - 6y + 2z = -24$$

$$C: \frac{x}{6} - \frac{y}{4} - \frac{z}{3} = 1 \Rightarrow 2x - 3y - 4z = 12$$

$$\therefore A: \quad \cancel{2x} + 3y - 6z = 42$$

$$2B: \underline{\cancel{2x} - 12y + 4z = -48}$$

$$D(\text{subtracting}): \quad 15y - 10z = 90$$

$$\text{also } 2B: \quad \cancel{2x} - 12y + 4z = -48$$

$$C: \underline{\cancel{2x} - 3y - 4z = 12}$$

$$E(\text{subtracting}): \quad -9y + 8z = -60$$

$$\therefore 4D: \quad 60y - \cancel{40z} = 360$$

$$5E: \quad \underline{-45y + \cancel{40z} = -300}$$

$$\begin{array}{rcl} \text{adding:} & 15y & = 60 \\ & y & = 4 \end{array}$$

$$\begin{array}{rcl} \text{since } D: & 15 - 10z & = 90 \\ & 15(4) - 10z & = 90 \\ & -10z & = 30 \\ & z & = -3 \end{array}$$

$$\begin{array}{rcl} \text{also } A: & 2x + 3y - 6z & = 42 \\ & 2x + 3(4) - 6(-3) & = 42 \\ & 2x + 12 + 18 & = 42 \\ & 2x & = 12 \\ & x & = 6 \end{array}$$

$$\therefore (x, y, z) = (6, 4, -3)$$

22. Curve: $x^2 + y^2 + ax + by + c = 0$.

$$\begin{array}{l} (1, 0) \in \text{curve} \Rightarrow \text{when } x = 1, y = 0 \\ \Rightarrow 1^2 + 0^2 + a(1) + b(0) + c = 0 \\ \Rightarrow 1 + a + c = 0 \\ \Rightarrow a + c = -1 \end{array} \quad : A$$

$$\begin{array}{l} (1, 2) \in \text{curve} \Rightarrow \text{when } x = 1, y = 2 \\ \Rightarrow 1^2 + 2^2 + a(1) + b(2) + c = 0 \\ \Rightarrow 1 + 4 + a + 2b + c = 0 \\ \Rightarrow a + 2b + c = -5 \end{array} \quad : B$$

$$\begin{array}{l} (2, 1) \in \text{curve} \Rightarrow \text{when } x = 2, y = 1 \\ \Rightarrow 2^2 + 1^2 + a(2) + b(1) + c = 0 \\ \Rightarrow 4 + 1 + 2a + b + c = 0 \\ \Rightarrow 2a + b + c = -5 \end{array} \quad : C$$

$$\begin{array}{l} \text{since } B: \quad a + 2b + c = -5 \\ \text{and } 2C: \quad 4a + 2b + 2c = -10 \end{array}$$

$$\begin{array}{rcl} D(\text{subtracting}): & -3a & -c = 5 \\ A: & \underline{a} & + c = -1 \end{array}$$

$$\begin{array}{rcl} \text{adding:} & -2a & = 4 \\ & a & = -2 \end{array}$$

$$\begin{array}{l} \text{since } A: \quad a + c = -1 \\ \Rightarrow \quad -2 + c = -1 \\ \quad \quad c = 1 \end{array}$$

$$\begin{array}{l} \text{also } B: \quad a + 2b + c = -5 \\ \quad \quad -2 + 2b + 1 = -5 \\ \quad \quad \quad 2b = -4 \\ \quad \quad \quad b = -2 \end{array}$$

$$\therefore (a, b, c) = (-2, -2, 1)$$

23. Let x be the number of small bags, y be the number of medium bags and z be the number of large bags. Then

$$\begin{array}{rcl} x + y + z & = & 15 \quad \dots 1 \\ 3x + 5y + 7z & = & 77 \quad \dots 2 \\ y = x + 2 & & \\ -x + y & = & 2 \quad \dots 3 \end{array}$$

Then

$$1: \quad x + y + z = 15$$

$$\begin{array}{rcl} 3: & \frac{-x + y}{2y + z} & = \frac{6}{17} \end{array} \quad \dots 4$$

Also

$$\begin{array}{rcl} 2: & 3x + 5y + 7z & = 77 \\ 3 \times 3: & \frac{-3x + 3y}{8y + 7z} & = \frac{6}{83} \end{array} \quad \dots 5$$

Then

$$4 \times -4: \quad -8y - 4z = -68$$

$$\begin{array}{rcl} 5: & 8y + 7z & = 83 \\ & \underline{3z} & = 15 \\ & z & = 5 \end{array}$$

$$\begin{array}{rcl} 4: & 2y + 5 & = 17 \\ & 2y & = 12 \\ & y & = 6 \end{array}$$

$$\begin{array}{rcl} 3: & 6 & = x + 2 \\ & x & = 4 \end{array}$$

The shop sells 4 small, 6 medium and 5 large bags of popcorn.

- 24.** Let € x be the amount in the mutual fund, € y be the amount in government bonds and € z be the amount in her local bank. Then

$$x + y + z = 10500 \quad \dots 1$$

$$x = 2z$$

$$x - 2z = 0 \quad \dots 2$$

$$0.11x + 0.07y + 0.05z = 825$$

$$11x + 7y + 5z = 82500 \quad \dots 3$$

Then

$$1 \times 2: \quad 2x + 2y + 2z = 21\,000$$

$$\begin{array}{rcl} 2: & x & - 2z = 0 \\ & \underline{3x + 2y} & = 21\,000 \end{array} \quad \dots 4$$

$$1 \times -5: \quad -5x - 5y - 5z = -52\,500$$

$$\begin{array}{rcl} 3: & \frac{11x + 7y + 5z}{6x + 2y} & = \frac{82\,500}{30\,000} \end{array} \quad \dots 5$$

Then

$$4 \times -1: \quad -3x - 2y = -21\,000$$

$$\begin{array}{rcl} 5: & 6x + 2y & = 30\,000 \\ & \underline{3x} & = 9\,000 \\ & x & = 3\,000 \end{array}$$

$$\begin{array}{rcl} 2: & x & = 2z \\ & 3\,000 & = 2z \\ & z & = 1\,500 \end{array}$$

$$\begin{array}{rcl} 1: & 3\,000 + y + 1\,500 & = 10\,500 \\ & y & = 6\,000 \end{array}$$

She invests €3000 in the mutual fund, €6000 in a government bond and €1500 in her local bank.

- 25.** Let x be the number of white beads, y be the number of blue beads and z be the number of green beads. Then

$$\begin{aligned}
 y &= (x + z) - 4 \\
 x - y + z &= 4 & \dots 1 \\
 z &= x + y \\
 x + y - z &= 0 & \dots 2 \\
 y &= 2x \\
 2x - y &= 0 & \dots 3
 \end{aligned}$$

Then

$$\begin{aligned}
 1: \quad & x - y + z = 4 \\
 2: \quad & \frac{x + y - z = 0}{2x = 4} \\
 & x = 2 \\
 3: \quad & y = 2x \\
 & y = 4 \\
 2: \quad & z = x + y \\
 & z = 2 + 4 \\
 & z = 6
 \end{aligned}$$

He uses 2 white, 4 blue and 6 green beads.

Revision Exercise 1 (Core)

$$\begin{aligned}
 1. \quad (i) \quad & \frac{12m^2n^3}{(6m^4n^5)^2} = \frac{\cancel{12}^1 m^2 n^3}{\cancel{36}_3 m^8 n^{10}} = \frac{1}{3m^6 n^7} \\
 (ii) \quad & \frac{3 + \frac{1}{x}}{\frac{5}{x} + 4} = \frac{\left(\frac{3x+1}{x}\right)}{\left(\frac{5+4x}{x}\right)} = \left(\frac{3x+1}{x}\right) \cdot \left(\frac{x}{5+4x}\right) = \frac{3x+1}{5+4x} \\
 (iii) \quad & \frac{2 + \frac{x}{2}}{x^2 - 16} = \frac{\frac{4+x}{2}}{(x-4)(x+4)} = \frac{\cancel{(x+4)}}{2(x-4)\cancel{(x+4)}} = \frac{1}{2x-8} \\
 2. \quad (i) \quad & \begin{array}{lll} y = x + 4 & \Rightarrow & y - x = 4 \quad : A \\ 5y + 2x = 6 & \Rightarrow & 5y + \cancel{2x} = 6 \quad : B \\ & \therefore & \underline{2y - \cancel{2x} = 8 \quad : 2A} \\ & \text{adding} & 7y = 14 \\ & & y = 2 \end{array}
 \end{aligned}$$

$$\begin{aligned}
 & \text{since } y = x + 4 \\
 & \Rightarrow 2 = x + 4 \\
 & \Rightarrow x = -2 \\
 & \therefore (x, y) = (-2, 2)
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad 3x + y &= 7 \Rightarrow y = 7 - 3x \\
 x^2 + y^2 &= 13 \Rightarrow x^2 + (7 - 3x)^2 = 13 \\
 &\Rightarrow x^2 + [49 - 42x + 9x^2] = 13 \\
 &\quad x^2 + 49 - 42x + 9x^2 = 13 \\
 &\quad 10x^2 - 42x + 36 = 0 \\
 &\quad 5x^2 - 21x + 18 = 0 \\
 &\quad (5x - 6)(x - 3) = 0
 \end{aligned}$$

$$\begin{aligned}
 \therefore x &= 3 \quad \text{or} \quad x = \frac{6}{5} \\
 \Rightarrow y &= 7 - 3(3) \quad \text{or} \quad y = 7 - 3\left(\frac{6}{5}\right) \\
 y &= 7 - 9 \quad \text{or} \quad y = 7 - \frac{18}{5} \\
 y &= -2 \quad \text{or} \quad y = \frac{17}{5}
 \end{aligned}$$

$$\therefore (x, y) = (3, -2) \quad \text{or} \quad \left(\frac{6}{5}, \frac{17}{5}\right)$$

$$\begin{array}{r} 3. \quad x^2 + 2x - 1 \\ x - 3 \overline{) x^3 - x^2 - 7x + 3} \\ \underline{x^3 - 3x^2} \quad \text{(subtracting)} \\ 2x^2 - 7x + 3 \\ \underline{2x^2 - 6x} \quad \text{(subtracting)} \\ -x + 3 \\ \underline{-x + 3} \quad \text{(subtracting)} \\ 0 \end{array}$$

answer: $x^2 + 2x - 1$

$$\begin{array}{r} 4. \quad 3x^3 + 6x^2 + 3x + 33 \\ x - 2 \overline{) 3x^4 \quad - 9x^2 + 27x - 66} \\ \underline{3x^4 - 6x^3} \quad \text{(subtracting)} \\ 6x^3 - 9x^2 + 27x - 66 \\ \underline{6x^3 - 12x^2} \quad \text{(subtracting)} \\ 3x^2 + 27x - 66 \\ \underline{3x^2 - 6x} \quad \text{(subtracting)} \\ 33x - 66 \\ \underline{33x - 66} \quad \text{(subtracting)} \\ 0 \end{array}$$

answer: $3x^3 + 6x^2 + 3x + 33$

$$\begin{aligned} 5. \quad (i) \quad x^4 - 9x^2 &= 0 \\ \Rightarrow x^2(x^2 - 9) &= 0 \\ \Rightarrow x^2(x - 3)(x + 3) &= 0 \\ \therefore x &= 0, 3, -3 \end{aligned}$$

$$\begin{aligned} (ii) \quad (2x - 1)^3(2 - x) &= 0 \\ \Rightarrow (2x - 1)^3 &= 0 \\ \Rightarrow (2x - 1) &= 0 \end{aligned}$$

$$x = \frac{1}{2}$$

$$\begin{aligned} \text{or} \quad 2 - x &= 0 \\ x &= 2 \end{aligned}$$

$$\therefore x = 2, \frac{1}{2}$$

$$\begin{aligned} 6. \quad 4x^2 + 20x + k &\text{ is a perfect square} \\ \Rightarrow (2x + a)(2x + a) &= 4x^2 + 20x + k \\ \Rightarrow 4x^2 + 4ax + a^2 &= 4x^2 + 20x + k \\ \Rightarrow 4a &= 20 \\ a &= 5 \end{aligned}$$

$$\therefore a^2 = 5^2 = 25 = k$$

$$\therefore k = 25(5^2)$$

$$\begin{aligned} 7. \quad (i) \quad (2x + 3)^5 &= \binom{5}{0}(2x)^5 + \binom{5}{1}(2x)^4(3) + \binom{5}{2}(2x)^3(3)^2 + \binom{5}{3}(2x)^2(3)^3 + \binom{5}{4}(2x)(3)^4 + \binom{5}{5}(3)^5 \\ &= (1)(32x^5) + (5)(16x^4)(3) + (10)(8x^3)(9) + (10)(4x^2)(27) + (5)(2x)(81) + (1)(243) \\ &= 32x^5 + 240x^4 + 720x^3 + 1080x^2 + 810x + 243 \end{aligned}$$

$$\begin{aligned} (ii) \quad (x - 2)^6 &= \binom{6}{0}x^6 + \binom{6}{1}x^5(-2) + \binom{6}{2}x^4(-2)^2 + \binom{6}{3}x^3(-2)^3 + \binom{6}{4}x^2(-2)^4 + \binom{6}{5}x(-2)^5 + \binom{6}{6}(-2)^6 \\ &= (1)x^6 + (6)x^5(-2) + (15)x^4(4) + (20)x^3(-8) + (15)x^2(16) + (6)x(-32) + (1)(64) \\ &= x^6 - 12x^5 + 60x^4 - 160x^3 + 240x^2 - 192x + 64 \end{aligned}$$

$$8. \quad x^3 - 27 = x^3 - 3^3 = (x - 3)(x^2 + 3x + 9)$$

9. $p(x - q)^2 + r = 2x^2 - 12x + 5$ for all values of x

$$\Rightarrow p(x^2 - 2xq + q^2) + r =$$

$$\Rightarrow px^2 - 2pqx + pq^2 + r = 2x^2 - 12x + 5.$$

$$\therefore p = 2, -2pq = -12 \quad \text{and} \quad pq^2 + r = 5.$$

$$\Rightarrow -2(2)q = -12$$

$$\Rightarrow q = 3 \quad \text{and} \quad 2(3)^2 + r = 5$$

$$r = 5 - 18$$

$$r = -13$$

$$(p, q, r) = (2, 3, -13)$$

10. A: $3x + 5y - z = -3$

B: $2x + y - 3z = -9$

C: $x + 3y + 2z = 7$

$$\Rightarrow 2A: 6x + 10y - 2z = -6$$

$$C: x + 3y + 2z = 7$$

$$D(\text{adding}): 7x + 13y = 1$$

also $3A: 9x + 15y - 3z = -9$

$$B: 2x + y - 3z = -9$$

$$E(\text{subtracting}): 7x + 14y = 0$$

$$\therefore D: 7x + 13y = 1$$

$$E: 7x + 14y = 0$$

$$(\text{subtracting}): -y = 1$$

$$y = -1$$

since $7x + 14y = 0$

$$7x + 14(-1) = 0$$

$$7x = 14$$

$$x = 2.$$

also, C: $x + 3y + 2z = 7$

$$2 + 3(-1) + 2z = 7$$

$$2z = 8$$

$$z = 4.$$

$$\therefore (x, y, z) = (2, -1, 4)$$

11. $(b + 1)^3 - (b - 1)^3$

$$= b^3 + 3b^2 + 3b + 1 - (b^3 - 3b^2 + 3b - 1)$$

$$= b^3 + 3b^2 + 3b + 1 - b^3 + 3b^2 - 3b + 1$$

$$= 6b^2 + 2.$$

12. (i) 3, 12, 27, 48, 75, ...

first difference: 9, 15, 21, 27, ...

second difference: 6, 6, 6... \Rightarrow quadratic pattern of the form $an^2 + bn + c$

$$\Rightarrow 2a = 6$$

$$a = 3$$

$$\therefore 3n^2 + bn + c \text{ represents the pattern}$$

let $n = 1 \Rightarrow 3(1)^2 + b(1) + c = 3$

$$b + c = 0 \quad : A$$

let $n = 2 \Rightarrow 3(2)^2 + b(2) + c = 12$

$$\begin{array}{rcl}
 & & 2b + c = 0 \quad : B \\
 \Rightarrow & A: & b + c = 0 \\
 & B: & \underline{2b + c = 0} \\
 \text{(subtracting):} & & -b = 0 \\
 \Rightarrow & & b = 0
 \end{array}$$

$$\text{since } b + c = 0$$

$$\Rightarrow 0 + c = 0$$

$$\Rightarrow c = 0$$

$$\therefore an^2 + bn + c = 3n^2$$

(ii) 5, 20, 45, 80, 125, ...

first difference: 15, 25, 35, 45, ...

second difference: 10, 10, 10, ...

\therefore quadratic pattern of the form $an^2 + bn + c$

$$\Rightarrow 2a = 10$$

$$a = 5 \quad \therefore 5n^2 + bn + c \text{ represents the pattern}$$

We note that for $n = 1$: $5n^2 = 5$

$$n = 2: 5n^2 = 20$$

$$n = 3: 5n^2 = 45, \text{ etc.}$$

$\Rightarrow b$ and c must equal zero

[alternatively, set up simultaneous equations in b and c and solve]

\therefore the quadratic pattern is $5n^2$

(iii) 0.5, 2, 4.5, 8, 12.5, ...

first difference = 1.5, 2.5, 3.5, 4.5, ...

second difference = 1, 1, 1

\therefore quadratic pattern of the form $an^2 + bn + c$

$$\Rightarrow 2a = 1$$

$$a = \frac{1}{2} \quad \therefore 0.5n^2 + bn + c \text{ represents this pattern}$$

By comparison to part (ii), we can deduce that the quadratic pattern is $0.5n^2$.

13. 6, 12, 20, 30, 42, ...

first difference: 6, 8, 10, 12, ...

second difference: 2, 2, 2

\therefore quadratic pattern of the form $an^2 + bn + c$

$$\Rightarrow 2a = 2$$

$$a = 1 \Rightarrow n^2 + bn + c \text{ represents this pattern}$$

$$\text{let } n = 1 \Rightarrow 1^2 + b(1) + c = 6$$

$$b + c = 5 \quad : A$$

$$\text{let } n = 2 \Rightarrow 2^2 + b(2) + c = 12$$

$$2b + c = 8 \quad : B$$

$$\text{since } A: b + c = 5$$

$$\text{and } B: \underline{2b + c = 8}$$

$$\text{subtracting: } -b = -3$$

$$b = 3$$

$$\text{also } A: b + c = 5$$

$$3 + c = 5$$

$$c = 2$$

∴ quadratic pattern is $n^2 + 3n + 2$

$$\text{When } n = 100 \Rightarrow 100^2 + 3(100) + 2 = 10,302$$

14. Let the width = x cm.

Let the length = y cm.

$$\Rightarrow A: 3x = 2y + 3$$

$$B: 4y = 2(x + y) + 12$$

$$\Rightarrow B: 4y = 2x + 2y + 12$$

$$\Rightarrow B: 2y = 2x + 12.$$

$$\text{since } A: 3x - 2y = 3$$

$$\text{and } B: \frac{2x - 2y}{x} = -12$$

$$(\text{subtracting}): \frac{2x - 2y}{x} = 15 \Rightarrow \text{width} = 15 \text{ cm.}$$

$$\text{also } B: 2x - 2y = -12$$

$$2(15) - 2y = -12$$

$$-2y = -42$$

$$y = 21 \Rightarrow \text{length} = 21 \text{ cm.}$$

15. $A: \frac{1}{u} + \frac{1}{v} = \frac{2}{r}$

$$B: m = \frac{v-r}{r-u}$$

$$\text{since } A: \frac{1}{u} + \frac{1}{v} = \frac{2}{r}$$

$$\Rightarrow \frac{v+u}{uv} = \frac{2}{r}$$

$$\Rightarrow \frac{2}{r} = \frac{v+u}{uv}$$

$$\Rightarrow \frac{r}{2} = \frac{u \cdot v}{v+u}$$

$$\Rightarrow r = \frac{2uv}{v+u}$$

$$\text{also, } m = \frac{v-r}{r-u} = \frac{v - \frac{2uv}{v+u}}{\frac{2uv}{v+u} - u}$$

$$= \frac{\left(\frac{v(v+u) - 2uv}{v+u} \right)}{\left(\frac{2uv - u(v+u)}{v+u} \right)}$$

$$= \frac{v^2 + vu - 2uv}{2uv - uv - u^2} \cdot \frac{(v+u)}{(v+u)}$$

$$= \frac{v^2 - uv}{uv - u^2} = \frac{v(v-u)}{u(v-u)} = \frac{v}{u}$$

Revision Exercise 1 (Advanced)

1. 1, 3, 6, 10, ...

first difference: 2, 3, 4, ...

second difference: 1, 1, ...

 \Rightarrow quadratic pattern of the form $an^2 + bn + c$ $\Rightarrow 2a = 1$ $a = \frac{1}{2} \Rightarrow \frac{1}{2}n^2 + bn + c$ represents this pattern

$$\text{let } n = 1: \quad \frac{1}{2}(1)^2 + b(1) + c = 1$$

$$\frac{1}{2} + b + c = 1$$

$$b + c = \frac{1}{2} \quad :A$$

$$\text{let } n = 2: \quad \frac{1}{2}(2)^2 + b(2) + c = 3$$

$$2 + 2b + c = 3$$

$$2b + c = 1 \quad :B$$

$$\text{since } A: \quad b + c = \frac{1}{2}$$

$$\text{and } B: \quad 2b + c = 1$$

$$(\text{subtracting}): \quad -b = -\frac{1}{2}$$

$$b = \frac{1}{2}$$

$$\text{also } A: \quad b + c = \frac{1}{2}$$

$$\Rightarrow \quad \frac{1}{2} + c = \frac{1}{2}$$

$$\Rightarrow \quad c = 0$$

\therefore the quadratic pattern is $\frac{1}{2}n^2 + \frac{1}{2}n$

2. If $x \text{ m}^3$ of soil is needed,

$$\Rightarrow x(55\%) + 1(25\%) = (x + 1)(35\%)$$

$$\Rightarrow 0.55x + 0.25 = 0.35x + 0.35$$

$$\Rightarrow 0.55x - 0.35x = 0.35 - 0.25$$

$$0.2x = 0.1$$

$$x = 0.5 \text{ m}^3.$$

3. (i) Let $x \text{ kg}$ of alloy 1 be added to $y \text{ kg}$ of alloy 2.

$$\Rightarrow x + y = 8.4 \quad :A$$

alloy 1 contains 60% gold

$$\Rightarrow \text{the amount of gold in the new alloy} = x(60\%) = 0.6x$$

alloy 2 contains 40% gold

$$\Rightarrow \text{the amount of gold in the new alloy} = y(40\%) = 0.4y$$

$$\text{also, the total amount of gold in the new alloy} = (x + y)(50\%) = 0.5(x + y)$$

$$\Rightarrow 0.6x + 0.4y = 0.5(x + y) \quad :B$$

$$\text{since } A: \quad x + y = 8.4$$

$$\text{and } B: \quad 6x + 4y = 5(x + y)$$

$$\Rightarrow B: \quad x - y = 0 \quad (\text{simplifying } B)$$

$$\begin{aligned}
 \text{and } A: \quad x + y &= 8.4 \\
 \Rightarrow \quad 2x &= 8.4 \\
 x &= 4.2 \text{ kg} \\
 \text{since } B: \quad x - y &= 0 \\
 \Rightarrow \quad 4.2 - y &= 0 \\
 \Rightarrow \quad y &= 4.2 \text{ kg also.}
 \end{aligned}$$

$$\begin{aligned}
 4. \quad (3p - 2t)x + r - 4t^2 &= 0 \quad \text{for all } x. \\
 \Rightarrow (3p - 2t)x + r - 4t^2 &= 0 \cdot x + 0 \\
 \Rightarrow 3p - 2t &= 0 \quad \Rightarrow t = \frac{3p}{2}
 \end{aligned}$$

$$\text{and } r - 4t^2 = 0.$$

$$\Rightarrow r - 4\left(\frac{3p}{2}\right)^2 = 0$$

$$\Rightarrow r - \frac{4 \cdot 9p^2}{4} = 0$$

$$\Rightarrow r = 9p^2.$$

$$\begin{aligned}
 5. \quad \frac{x + y^2}{x^2} + \frac{x - 1}{x} &= -1 \\
 \Rightarrow x + y^2 + x(x - 1) &= -x^2 \quad [\text{multiplying each term by } x^2] \\
 \Rightarrow x + y^2 + x^2 - x &= -x^2 \\
 \Rightarrow x + y^2 + x^2 - x + x^2 &= 0 \\
 2x^2 + y^2 &= 0.
 \end{aligned}$$

$$\text{also, } \frac{2x^2}{y^2} + 1 = 0.$$

$$\frac{2x^2}{y^2} = -1$$

$$\frac{x^2}{y^2} = -\frac{1}{2}$$

$$6. \text{ Let the students take } x \text{ litres of 10\% solution and } y \text{ litres of 30\% solution.}$$

$$\Rightarrow x + y = 10 \text{ litres}$$

$$\text{also, } x(10\%) + y(30\%) = (x + y)15\%$$

$$\Rightarrow 10x + 30y = 15(x + y) \quad [\text{multiplying each term by 100}]$$

$$\therefore x + y = 10 \quad : A$$

$$\text{and } 10x + 30y = 15x + 15y$$

$$\Rightarrow -5x + 15y = 0 \quad : B$$

$$\text{since } A: \quad x + y = 10$$

$$\text{and } B: -5x + 15y = 0$$

$$\Rightarrow 5A: \quad 5x + 5y = 50$$

$$\text{adding:} \quad \frac{5x + 5y = 50}{-5x + 15y = 0}$$

$$y = \frac{50}{20} = 2\frac{1}{2} \text{ litres.}$$

$$\text{also, since } x + y = 10$$

$$\Rightarrow x + 2\frac{1}{2} = 10$$

$$x = 7\frac{1}{2} \text{ litres.}$$

$$(i) 7\frac{1}{2} \text{ litres of 10\% mixed with } (ii) 2\frac{1}{2} \text{ litres of 30\%.$$

$$7. \quad 3x - y = 0 \quad \dots (1)$$

$$x + y + z = 23\,000 \quad \dots (2)$$

$$75x + 55y + 30z = 870\,000$$

$$15x + 11y + 6z = 174\,000 \quad \dots (3)$$

$$(2) \times 6: \quad 6x + 6y + 6z = 138\,000$$

$$15x + 11y + 6z = 174\,000$$

$$\underline{6x + 6y + 6z = 138\,000}$$

$$9x + 5y = 36\,000$$

$$(1) \times 3: \quad \underline{9x - 3y = 0}$$

$$8y = 36\,000$$

$$y = 4500$$

$$\therefore 3x = 4500$$

$$x = 1500$$

$$\therefore x + y + z = 1500 + 4500 + z = 23\,000$$

$$\therefore z = 17\,000$$

$$\therefore x = 1500, y = 4500, z = 17\,000$$

8. Brian's time for 50 m = $50a$ seconds.

Luke's time for 50 m = $50b$ seconds.

Luke faster than Brian $\Rightarrow 50a = 50b + 1$

Brian's time for 47 m = $47a$ seconds.

Luke's time for 50 m = $50b$ seconds.

Luke faster than Brian $\Rightarrow 47a = 50b + 0.1$

Subtracting: $3a = 0.9$

$$a = 0.3$$

$$\Rightarrow 50(0.3) = 50b + 1$$

$$14 = 50b$$

$$0.28 = b$$

$$\text{Luke's speed} = \frac{1}{b} = \frac{1}{0.28} = 3.57 \text{ m/sec.}$$

9. (i) $\left(2 - \frac{x}{3}\right)^9$

$$\begin{aligned} \text{General term: } T_{r+1} &= \binom{9}{r} (2)^{9-r} \left(\frac{-x}{3}\right)^r \\ &= \binom{9}{r} 2^{9-r} \frac{(-1)^r x^r}{3^r} \end{aligned}$$

For the 6th term, $r = 5$. Thus the 6th term is

$$\begin{aligned} &\binom{9}{5} 2^{9-5} \frac{(-1)^5 x^5}{3^5} \\ &= (126)(16) \frac{(-1)x^5}{243} \\ &= -\frac{224}{27} x^5 \end{aligned}$$

- (ii) $(3x + 1)^{12}$

$$\begin{aligned} \text{General term: } T_{r+1} &= \binom{12}{r} (3x)^{12-r} (1)^r \\ &= \binom{12}{r} 3^{12-r} x^{12-r} \end{aligned}$$

For the 8th term, $r = 7$. Thus the 8th term is

$$\begin{aligned} &\binom{12}{7} 3^5 x^5 \\ &= (792)(243)x^5 \\ &= 192\,456x^5 \end{aligned}$$

10. $(3x - 1)^{11}$

$$\begin{aligned}\text{General term: } T_{r+1} &= \binom{11}{r}(3x)^{11-r}(-1)^r \\ &= (-1)^r \binom{11}{r} 3x^{11-r}x^r\end{aligned}$$

For the 6th term, $r = 5$. Thus the 6th term is

$$\begin{aligned}(-1)^5 \binom{11}{5} 3^6 x^6 \\ = (-1)(462)(729)x^6 \\ = -336\,798x^6\end{aligned}$$

11. $(2x - 3)^{14}$

$$\begin{aligned}\text{General term: } T_{r+1} &= \binom{14}{r}(2x)^{14-r}(-3)^r \\ &= \binom{14}{r} 2^{14-r} x^{14-r} (-1)^r 3^r\end{aligned}$$

For the term containing x^{10} ,

$$14 - r = 10$$

$$r = 4$$

The coefficient of x^{10} is then

$$\begin{aligned}\binom{14}{4} 2^{10} (-1)^4 3^4 \\ = (1001)(1024)(1)(81) \\ = 83\,026\,944\end{aligned}$$

Revision Exercise 1 (Extended-Response Questions)

- 1.** (a) Let x be the number of adults
 y be the number of children.

$$\therefore x + y = 548 \quad : A$$

$$\text{also, } 5x + 2.5y = 2460 : B$$

$$\text{since } A: \quad x + y = 548$$

$$\text{and } B: \quad 5x + 2.5y = 2460$$

$$\Rightarrow 5A: \quad 5x + 5y = 2740$$

$$(\text{subtracting}): \quad -2.5y = -280$$

$$2.5y = 280$$

$$y = \frac{280}{2.5} = 112.$$

$$\text{since } A: \quad x + y = 548$$

$$\Rightarrow x + 112 = 548$$

$$\Rightarrow x = 548 - 112$$

$$x = 436$$

$$\Rightarrow \text{(i) Number of adult tickets (x) = 436}$$

$$\text{(ii) Number of children tickets (y) = 112.}$$

$$\text{(iii) Proportion of adult tickets sold} = \frac{436}{548} = 0.7956.$$

$$\text{(b) attendance (predicted) = 13 000}$$

$$\Rightarrow \text{adults} = 0.7956 \times 13\,000 = 10\,343$$

$$\Rightarrow \text{children} = 13\,000 - 10\,343 = 2\,657$$

$$\text{Revenue} = 10\,343 \times (\text{€}5) + 2\,657 \times (\text{€}2.5) = \text{€}58\,357.50$$

- 2.** (i) x standard sofas

y deluxe sofas

Standard sofas require 2 hours of work $\Rightarrow 2x = \text{time for } x \text{ sofas}$

Deluxe sofas require 2.5 hours of work $\Rightarrow 2.5y = \text{time for } y \text{ sofas}$

\Rightarrow with 48 hours of manufacturing time: $2x + 2.5y = 48 : A$

(ii) also, standard sofas need 1 hour finishing $\Rightarrow x = \text{time for } x \text{ sofas}$

deluxe sofas need 1.5 hours finishing $\Rightarrow 1.5y = \text{time for } y \text{ sofas}$

\Rightarrow with 26 hours of finishing time: $x + 1.5y = 26 : B$

since $A: 2x + 2.5y = 48$

and $B: x + 1.5y = 26$

$\Rightarrow 2B: 2x + 3y = 52$

and $A: 2x + 2.5y = 48$

(subtracting): $0.5y = 4$

$y = 8$

also, since $B: x + 1.5y = 26$

$x + 1.5(8) = 26$

$x + 12 = 26$

$x = 14$

(iii) 14 standard sofas and 8 deluxe sofas.

3. Rectangular box with square base of length x cm.

Height of box = h cm.

Volume of box = $l \times b \times h$

$$= x \times x \times h = x^2 h.$$

(i) If $V = 40 \Rightarrow 40 = x^2 h$


$$\Rightarrow h = \frac{40}{x^2}.$$

(ii) Surface Area = $4 \times (x \times h) + 2x^2$

$$= 4 \times \frac{40}{x^2} + 2x^2$$

$$S = \frac{160}{x} + 2x^2.$$

(iii)




(iv) $72 = \frac{160}{x} + 2x^2$

$$72x = 160 + 2x^3$$

$$\Rightarrow 2x^3 - 72x + 160 = 0.$$

Using trail + error at $x = 2$: $2(2)^3 - 72(2) + 160 = 32 > 0$



$$\text{at } x = 2.8 : 2(2.8)^3 - 72(2.8) + 160 = 2.3 > 0$$

$$\text{at } x = 2.9 : 2(2.9)^3 - 72(2.9) + 160 = -0.02 < 0$$

$$\therefore \text{at } x = 2.9 \text{ cm (approximately), } S = 72 \text{ cm}^2$$

$$\text{also, at } x = 4 : 2(4)^3 - 72(4) + 160 = 0$$

$$\therefore \text{at } x = 4 \text{ cm, } S = 72 \text{ cm}^2$$

$$\text{when } x = 4, h = \frac{40}{(4)^2} = 2.5 \text{ cm.}$$

$$\text{when } x = 2.9, h = \frac{40}{(2.9)^2} = 4.76 \text{ cm.}$$

4. Selling price of game = €11.50.

Production cost for each game = €10.50.

Initial production costs = €3500.


(i) Cost of producing x games = $C(x)$

$$\Rightarrow C(x) = 10.5x + 3500$$

(ii) Income = $I(x)$

$$\Rightarrow I(x) = 11.5x.$$

(iii)



(iv) To recoup costs: $11.5x = 10.5x + 3500$

$$\Rightarrow x = 3500 \text{ need to be sold.}$$

(v) $P = I - C \Rightarrow P = \text{profit.}$

(vi) To make a profit of a 2000:

$$\Rightarrow P = I - C$$

$$\therefore 2000 = 11.5x - [10.5x + 3500]$$

$$2000 = 11.5x - 10.5x - 3500$$

$$5500 = x$$

$\therefore 5500$ need to be sold.

5. 15 days to complete quilt.

x blue squares at a rate of 4 squares a day.

y white squares at a rate of 7 squares a day.

96 squares in quilt $\Rightarrow x + y = 96 : A$

15 days to finish $\Rightarrow \frac{x}{4} + \frac{y}{7} = 15 : B$

$$\therefore A : x + y = 96$$

$$28B : 7x + 4y = 420$$

$$\text{also } 4A : 4x + 4y = 384$$

$$(\text{subtracting}): \frac{3x}{} = 36$$

$$x = 12$$


since $A : x + y = 96$

$$\Rightarrow 12 + y = 96$$

$$y = 84$$

$$\begin{aligned} \text{(a) Cost} &= x(0.8) + y(1.20) \\ &= 12(0.8) + 84(1.20) \\ &= \text{€}110.40 \end{aligned}$$

$$\begin{aligned} \text{(b) } l:w &= 3x:2x \\ \Rightarrow 3x \cdot 2x &= 96 \\ 6x^2 &= 96 \\ x^2 &= 16 \\ x &= 4 \\ \therefore l &= 3 \times 4 = 12 \\ w &= 2 \times 4 = 8 \end{aligned}$$



The 12 blue squares could form 2 rows of 6 in the centre. (There are many different possibilities.)

6. Overheads = €30 000 per year

Cost of manufacture = €40 per wheelbarrow

$$\text{(i) } C(x) = 40x + 30\,000$$

$$\text{(ii) } 6000 \text{ wheelbarrows per year} \Rightarrow \frac{30\,000}{6000} = \text{€}5 \text{ overhead per wheelbarrow}$$

$$\Rightarrow \text{Total cost per wheelbarrow} = \text{€}40 + \text{€}5 = \text{€}45$$

(iii) To get a cost of €46 per wheelbarrow:

$$\frac{30\,000}{x} + 40 = 46.$$


$$\Rightarrow 30\,000 + 40x = 46x$$

$$\Rightarrow 30\,000 = 6x$$

$$x = 5000 \text{ wheelbarrows}$$

(iv) Selling price = €80 per wheelbarrow

$$\Rightarrow \text{Revenue} = \text{€}80x$$



- (vi) To make a profit $80x > 40x + 30\,000$
 $\Rightarrow 40x > 30\,000$

$$x > \frac{30\,000}{40} = 750$$

The minimum number of wheelbarrows = 751.

- (vii) Profit $\text{€}P = 80x - [40x + 30\,000]$
 $\text{€}P = 40x - 30\,000$

Chapter 2

Exercise 2.1

1. (a) (i) $x = 4, x = -5$

(ii) $(x - 3)(x - 4) = 0$

$\Rightarrow x = 3, x = 4$

(iii) $(x + 1)(x - 5) = 0$

$\Rightarrow x = -1, x = 5$

(b) (i) $(x + 3)(x - 5) = 0$

$\Rightarrow x = -3, x = 5$

(ii) $(2x - 3)(x + 5) = 0$

$\Rightarrow 2x = 3, x = -5$

$\Rightarrow x = \frac{3}{2}, x = -5$

(iii) $(3x + 2)(x - 5) = 0$

$\Rightarrow 3x = -2, x = 5$

$\Rightarrow x = \frac{-2}{3}, x = 5$

(c) (i) $(5x + 2)(x - 3) = 0$

$\Rightarrow 5x = -2, x = 3$

$\Rightarrow x = \frac{-2}{5}, x = 3$

(ii) $(3x + 5)(3x - 4) = 0$

$\Rightarrow 3x = -5, 3x = 4$

$\Rightarrow x = \frac{-5}{3}, x = \frac{4}{3}$

(iii) $(4x + 5)(2x - 3) = 0$

$\Rightarrow 4x = -5, 2x = 3$

$\Rightarrow x = \frac{-5}{4}, x = \frac{3}{2}$

(d) (i) $(x + 3)(x - 3) = 0$

$\Rightarrow x = -3, x = 3$

(ii) $x(3x - 10) = 0$

$\Rightarrow x = 0, 3x = 10$

$\Rightarrow x = 0, x = \frac{10}{3}$

(iii) $x(5x - 8) = 0$

$\Rightarrow x = 0, 5x = 8$

$\Rightarrow x = 0, x = \frac{8}{5}$

(e) (i) $(5 + x)(3 - 2x) = 0$

$\Rightarrow x = -5, 2x = 3$

$\Rightarrow x = -5, x = \frac{3}{2}$

(ii) $(5 + 3x)(2 - x) = 0$

$\Rightarrow 3x = -5, x = 2$

$\Rightarrow x = \frac{-5}{3}, x = 2$

(iii) $(2 + x)(1 - x) = 0$

$\Rightarrow x = -2, x = 1$

(f) (i) $(x + 5)(x + 4)(x - 4) = 0$

$\Rightarrow x = -5, x = -4, x = 4$

(ii) $(x - 3)(x + 1)(x - 1) = 0$

$\Rightarrow x = 3, x = -1, x = 1$

(g) (i) $(x + 2)(x - 2)(3x + 4) = 0$

$\Rightarrow x = -2, x = 2, 3x = -4$

$\Rightarrow x = -2, x = 2, x = \frac{-4}{3}$

(ii) $(x + 4)(x + 3)(x - 5) = 0$

$\Rightarrow x = -4, x = -3, x = 5$

2. (a) (i) $x = \frac{2 \pm \sqrt{4 - 4(1)(-2)}}{2(1)}$

$= \frac{2 \pm \sqrt{12}}{2}$

$= \frac{2 \pm \sqrt{4\sqrt{3}}}{2}$

$= 1 \pm \sqrt{3}$

$= 2.7, 20.7$

(ii) $x = \frac{-3 \pm \sqrt{9 - 4(+1)(-2)}}{2(1)}$

$= \frac{-3 \pm \sqrt{17}}{2}$

$= 0.56, -3.56$

$= 0.6, -3.6$

(iii) $x = \frac{6 \pm \sqrt{36 - 4(2)(3)}}{2(2)}$

$= \frac{6 \pm \sqrt{12}}{(4)}$

$= 2.36, 0.635$

$= 2.4, 0.6$

$$\begin{aligned}
 \text{(b) (i) } x &= \frac{6 \pm \sqrt{36 - 4(1)(3)}}{2(1)} \\
 &= \frac{6 \pm \sqrt{24}}{2} \\
 &= 5.44, 0.55 \\
 &= 5.4, 0.6
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } x &= \frac{8 \pm \sqrt{64 - 4(3)(1)}}{2(3)} \\
 &= \frac{8 \pm \sqrt{52}}{6} \\
 &= 2.535, 0.13 \\
 &= 2.5, 0.1
 \end{aligned}$$

$$\begin{aligned}
 \text{3. (a) (i) } x &= \frac{-4 \pm \sqrt{16 - 4(3)(-5)}}{2(3)} \\
 &= \frac{-4 \pm \sqrt{76}}{6} \\
 &= \frac{-2 \pm \sqrt{19}}{3} \\
 \text{(ii) } x &= \frac{12 \pm \sqrt{144 - 4(2)(-5)}}{2(2)} \\
 &= \frac{12 \pm \sqrt{184}}{4} \\
 &= \frac{6 \pm \sqrt{46}}{2} \\
 \text{(iii) } 4x^2 - 12x + 9 &= 8 \\
 \Rightarrow 4x^2 - 12x + 1 &= 0 \\
 \Rightarrow x &= \frac{12 \pm \sqrt{144 - 4(4)(1)}}{2(4)} \\
 &= \frac{12 \pm \sqrt{128}}{8} \\
 &= \frac{3 \pm 2\sqrt{2}}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{4. (a) (i) } x(x + 7) + 2(3) &= 4(3)(x) \\
 \Rightarrow x^2 + 7x + 6 &= 12x \\
 \Rightarrow x^2 - 5x + 6 &= 0 \\
 \Rightarrow (x - 2)(x - 3) &= 0 \\
 \Rightarrow x = 2, x = 3 \\
 \text{(ii) } 1(x) + 4(x - 1) &= 3(x)(x - 1) \\
 \Rightarrow x + 4x - 4 &= 3x^2 - 3x \\
 \Rightarrow 3x^2 - 8x + 4 &= 0 \\
 \Rightarrow (3x - 2)(x - 2) &= 0 \\
 \Rightarrow 3x = 2, x = 2 \\
 \Rightarrow x = \frac{2}{3}, x = 2 \\
 \text{(iii) } 3(x + 1) - 2(x - 1) &= 1(x - 1)(x + 1) \\
 \Rightarrow 3x + 3 - 2x + 2 &= x^2 - 1 \\
 \Rightarrow x^2 - x - 6 &= 0 \\
 \Rightarrow (x + 2)(x - 3) &= 0 \\
 \Rightarrow x = -2, x = 3
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } x &= \frac{-4 \pm \sqrt{16 - 4(2)(-5)}}{2(2)} \\
 &= \frac{-4 \pm \sqrt{56}}{4} \\
 &= 0.87, -2.87 \\
 &= 0.9, -2.9
 \end{aligned}$$

$$\begin{aligned}
 \text{(b) (i) } x &= \frac{-4 \pm \sqrt{16 - 4(1)(-8)}}{2(1)} \\
 &= \frac{-4 \pm \sqrt{48}}{2} \\
 &= \frac{-4 \pm 4\sqrt{3}}{2} \\
 &= -2 \pm 2\sqrt{3} \\
 \text{(ii) } x &= \frac{-4 \pm \sqrt{16 - 4(5)(-2)}}{2(5)} \\
 &= \frac{-4 \pm \sqrt{56}}{10} \\
 &= \frac{-2 \pm \sqrt{14}}{5} \\
 \text{(iii) } x &= \frac{1 \pm \sqrt{1 - 4(1)(-1)}}{2(1)} \\
 &= \frac{1 \pm \sqrt{5}}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{(b) (i) } 1(x - 2) + 2(x) &= 3(x)(x - 2) \\
 \Rightarrow x - 2 + 2x &= 3x^2 - 6x \\
 \Rightarrow 3x^2 - 9x + 2 &= 0 \\
 \Rightarrow x &= \frac{9 \pm \sqrt{81 - 4(3)(2)}}{2(3)} = \frac{9 \pm \sqrt{57}}{6} \\
 \text{(ii) } (x + 2)(x - 2) &= (2x + 1)(x - 4) \\
 \Rightarrow x^2 - 4 &= 2x^2 - 7x - 4 \\
 \Rightarrow x^2 - 7x &= 0 \\
 \Rightarrow x(x - 7) &= 0 \\
 \Rightarrow x = 0, x = 7 \\
 \text{(iii) } 2(x)(x - 4) + 3(x - 2)(x - 4) &= 5(x)(x - 2) \\
 \Rightarrow 2x^2 - 8x + 3x^2 - 12x - 6x + 24 &= 5x^2 - 10x \\
 \Rightarrow 5x^2 - 26x + 24 &= 5x^2 - 10x \\
 \Rightarrow -16x &= -24 \\
 \Rightarrow x &= \frac{-24}{-16} = \frac{3}{2}
 \end{aligned}$$

5. (a) (i) $y = x^2 \Rightarrow y^2 - 7y + 10 = 0$

$$\Rightarrow (y - 2)(y - 5) = 0$$

$$\Rightarrow y = 2, y = 5$$

$$\text{Hence: } x^2 = 2, x^2 = 5$$

$$\Rightarrow x = \pm\sqrt{2}, x = \pm\sqrt{5}$$

(ii) $y = x + 1 \Rightarrow y^2 + 3y - 2 = 0$

$$\Rightarrow y = \frac{-3 \pm \sqrt{9 - 4(1)(-2)}}{2}$$

$$\Rightarrow y = \frac{-3 \pm \sqrt{17}}{2}$$

$$\Rightarrow x + 1 = \frac{-3 \pm \sqrt{17}}{2}$$

$$\Rightarrow x = \frac{-5 \pm \sqrt{17}}{2}$$

(iii) $y = x^2 \Rightarrow y^2 - 2y - 2 = 0$

$$\Rightarrow y = \frac{2 \pm \sqrt{4 - 4(1)(-2)}}{2}$$

$$\Rightarrow y = \frac{2 \pm \sqrt{12}}{2} = 1 \pm \sqrt{3}$$

$$\text{Hence: } x^2 = 1 \pm \sqrt{3}$$

$$\Rightarrow x = \pm\sqrt{1 \pm \sqrt{3}}$$

(iv) $y = k - 2 \Rightarrow 2y^2 - 3y - 4 = 0$

$$\Rightarrow y = \frac{3 \pm \sqrt{9 - 4(2)(-4)}}{2(2)}$$

$$= \frac{3 \pm \sqrt{41}}{4}$$

$$\text{Hence: } k - 2 = \frac{3 \pm \sqrt{41}}{4}$$

$$\Rightarrow k = \frac{11 \pm \sqrt{41}}{4}$$

(b) (i) $k = 2y - 1 \Rightarrow k^2 - 3k - 28 = 0$

$$\Rightarrow (k + 4)(k - 7) = 0$$

$$\Rightarrow k = -4, k = 7$$

$$\text{Hence: } 2y - 1 = -4, 2y - 1 = 7$$

$$\Rightarrow 2y = -3, 2y = 8$$

$$\Rightarrow y = -\frac{3}{2}, y = 4$$

6. $x = \frac{\sqrt{3} \pm \sqrt{3 - 4(2)(-3)}}{2(2)}$

$$= \frac{\sqrt{3} \pm \sqrt{27}}{4}$$

$$= \frac{\sqrt{3} \pm 3\sqrt{3}}{4}$$

$$= \frac{4\sqrt{3}}{4}, -\frac{2\sqrt{3}}{4}$$

$$= \sqrt{3}, -\frac{\sqrt{3}}{2}$$

(ii) $k = 2y - 3 \Rightarrow k^2 - 1 = 0$

$$\Rightarrow (k + 1)(k - 1) = 0$$

$$\Rightarrow k = -1, k = 1$$

$$\text{Hence: } 2y - 3 = -1, 2y - 3 = 1$$

$$\Rightarrow 2y = 2, 2y = 4$$

$$\Rightarrow y = 1, y = 2$$

(c) $k = y + \frac{4}{y} \Rightarrow k^2 - 9k + 20 = 0$

$$\Rightarrow (k - 4)(k - 5) = 0$$

$$\Rightarrow k = 4, k = 5$$

$$\text{Hence: } y + \frac{4}{y} = 4, y + \frac{4}{y} = 5$$

$$\Rightarrow y^2 + 4 = 4y, y^2 + 4 = 5y$$

$$\Rightarrow y^2 - 4y + 4 = 0,$$

$$y^2 - 5y + 4 = 0$$

$$\Rightarrow (y - 2)(y - 2) = 0,$$

$$(y - 1)(y - 4) = 0$$

$$\Rightarrow y = 2, y = 1, y = 4$$

(d) $x = 2t - \frac{5}{t} = 3 \Rightarrow x^2 - 12x + 27 = 0$

$$\Rightarrow (x - 3)(x - 9) = 0$$

$$\Rightarrow x = 3, x = 9$$

$$\text{Hence: } 2t - \frac{5}{t} = 3, 2t - \frac{5}{t} = 9$$

$$\Rightarrow 2t^2 - 5 = 3t, 2t^2 - 5 = 9t$$

$$\Rightarrow 2t^2 - 3t - 5 = 0, 2t^2 - 9t - 5 = 0$$

$$\Rightarrow (t + 1)(2t - 5) = 0, (2t + 1)(t - 5) = 0,$$

$$\Rightarrow t = -1, 2t = 5, 2t = -1, t = 5$$

$$\Rightarrow t = -1, t = \frac{5}{2}, t = -\frac{1}{2}, t = 5$$

7. (a) $x = -4.2, 1.2$

(b) $x = -2.3, 1.3$

(c) $x = -9.5, -0.5$

(d) $-1.6 < x < 0.6$

(e) $x = -1.6, 0.6$

(f) $x = -3, 1$

(g) $x = -3.6, 0.6$

(h) $-4.1 < x < -0.9$

8. Graph does not intersect the x-axis

9. (a) $x_2 - x_1 = -22.5 + 2.5 = -20$
 (b) $x_2 + x_1 = -22.5 - 2.5 = -25$
10. (a) $x = -0.5, 2$
 (b) $x = -0.8$
 (c) $x = -0.5, 2.4$

11. (i) $u = \sqrt{x}$ Equation is:

$$\begin{aligned} 2u^2 + 3u - 5 &= 0 \\ (2u + 5)(u - 1) &= 0 \\ 2u + 5 &= 0 \quad \text{or} \quad u - 1 = 0 \\ u &= -\frac{5}{2} \quad \text{or} \quad u = 1 \end{aligned}$$

As $u = \sqrt{x}$ must be positive, $u = -\frac{5}{2}$ is not possible. Thus

$$u = \sqrt{x} = 1$$

$$u^2 = x = 1$$

is the only solution for x .

- (ii) $u = \sqrt{x}$ Equation is:

$$\begin{aligned} u^2 - 3u - 4 &= 0 \\ (u + 1)(u - 4) &= 0 \\ u + 1 &= 0 \quad \text{or} \quad u - 4 = 0 \\ u &= -1 \quad \text{or} \quad u = 4 \end{aligned}$$

As $u = \sqrt{x}$ must be positive, $u = -1$ is not possible. Thus

$$u = \sqrt{x} = 4$$

$$u^2 = x = 16$$

is the only solution for x .

$$\begin{aligned} 12. \quad (i) \quad x &= \frac{\sqrt{7} \pm \sqrt{7 - 4(1)(-14)}}{2(1)} \\ &= \frac{\sqrt{7} \pm \sqrt{63}}{2} \\ &= \frac{\sqrt{7} + 3\sqrt{7}}{2}, \frac{\sqrt{7} - 3\sqrt{7}}{2} \\ &= 2\sqrt{7}, -\sqrt{7} \end{aligned}$$

$$\begin{aligned} (ii) \quad x &= \frac{-7\sqrt{5} \pm \sqrt{245 - 4(2)(15)}}{2(2)} \\ &= \frac{-7\sqrt{5} \pm \sqrt{125}}{4} \\ &= \frac{-7\sqrt{5} + 5\sqrt{5}}{4}, \frac{-7\sqrt{5} - 5\sqrt{5}}{4} \\ &= -\frac{\sqrt{5}}{2}, -3\sqrt{5} \end{aligned}$$

Exercise 2.2


1. (i) f
 (ii) h
 (iii) g
 (iv) Curve f has roots = 1.6, 4.4
 Curve h has root = 3

$$\begin{aligned} 2. \quad y = 0 &\Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &\Rightarrow A = \left(\frac{-b - \sqrt{b^2 - 4ac}}{2a}, 0 \right) \\ &\Rightarrow B = \left(\frac{-b + \sqrt{b^2 - 4ac}}{2a}, 0 \right) \end{aligned}$$

3. (i) $b^2 - 4ac = (1)^2 - 4(2)(5) = -39 < 0 \Rightarrow$ Imaginary roots
 (ii) $b^2 - 4ac = (3)^2 - 4(-2)(1) = 17 > 0$
 \Rightarrow Roots are real and different
 (iii) $b^2 - 4ac = (2)^2 - 4(3)(-1) = 16 > 0$
 \Rightarrow Roots are real and different

- (iv) $b^2 - 4ac = (2)^2 - 4(-1)(-3) = -8 < 0$
 \Rightarrow Imaginary roots
- (v) $b^2 - 4ac = (8)^2 - 4(1)(16) = 0$
 \Rightarrow Roots are real and equal
- (vi) $b^2 - 4ac = (-10)^2 - 4(1)(25) = 0$
 \Rightarrow Roots are real and equal

4. $y = x^2 - 2x + 5$ is positive for all values of x
 $3x^2 - kx + 12$ is positive
 \Rightarrow Roots are imaginary
 $\Rightarrow b^2 - 4ac < 0$
 $\Rightarrow (-k^2) - 4(3)(12) < 0$
 $\Rightarrow k^2 - 144 < 0$
 \Rightarrow Factors: $(k + 12)(k - 12)$
 \Rightarrow Roots: $k = -12, 12$
 \Rightarrow Solution for $k^2 - 144 < 0 \Rightarrow -12 < k < 12$



5. Equal Roots $\Rightarrow b^2 - 4ac = 0$

(i) $(-10)^2 - 4(1)(k) = 0$
 $\Rightarrow 100 - 4k = 0$
 $\Rightarrow -4k = -100$
 $\Rightarrow k = 25$

(ii) $(k)^2 - 4(4)(9) = 0$
 $\Rightarrow k^2 - 144 = 0$
 $\Rightarrow (k + 12)(k - 12) = 0$
 $\Rightarrow k = -12, 12$

(iii) $(-2k - 2)^2 - 4(1)(5k + 1) = 0$
 $\Rightarrow 4k^2 + 8k + 4 - 20k - 4 = 0$
 $\Rightarrow 4k^2 - 12k = 0$
 $\Rightarrow k^2 - 3k = 0$
 $\Rightarrow k(k - 3) = 0$
 $\Rightarrow k = 0, k = 3$

6. Equal Roots $\Rightarrow (2k + 2)^2 - 4(k^2)(4) = 0$
 $\Rightarrow 4k^2 + 8k + 4 - 16k^2 = 0$
 $\Rightarrow 12k^2 - 8k - 4 = 0$
 $\Rightarrow 3k^2 - 2k - 1 = 0$
 $\Rightarrow (3k + 1)(k - 1) = 0$
 $\Rightarrow k = -\frac{1}{3}, k = 1$

7. (i) Real Roots $\Rightarrow (-3k)^2 - 4(1)(-k^2) \geq 0$
 $\Rightarrow 9k^2 + 4k^2 \geq 0$
 $\Rightarrow 13k^2 \geq 0$ True
- (ii) Real Roots $\Rightarrow (2)^2 - 4(k)(2 - k) \geq 0$
 $\Rightarrow 4 - 8k + 4k^2 \geq 0$
 $\Rightarrow k^2 - 2k + 1 \geq 0$
 $\Rightarrow (k - 1)^2 \geq 0$ True

8. Real Roots $\Rightarrow (-3)^2 - 4(1)(2 - c^2) \geq 0$
 $\Rightarrow 9 - 8 + 4c^2 \geq 0$
 $\Rightarrow 1 + 4c^2 \geq 0$ True

9. Real Roots $\Rightarrow (2)^2 - 4(k - 2)(-k) \geq 0$
 $\Rightarrow 4 + 4k^2 - 8k \geq 0$
 $\Rightarrow k^2 - 2k + 1 \geq 0$
 $\Rightarrow (k - 1)^2 \geq 0$ True

$$\begin{aligned}
 10. \text{ Equal Roots } &\Rightarrow (2k + 1)^2 - 4(k - 2)(k) = 0 \\
 &\Rightarrow 4k^2 + 4k + 1 - 4k^2 + 8k = 0 \\
 &\Rightarrow 12k = -1 \\
 &\Rightarrow k = -\frac{1}{12}
 \end{aligned}$$

$$\begin{aligned}
 11. \text{ Equal Roots } &\Rightarrow (6 - 2m)^2 - 4(m + 3)(m - 1) = 0 \\
 &\Rightarrow 36 - 24m + 4m^2 - 4m^2 - 12m + 4m + 12 = 0 \\
 &\Rightarrow -32m + 48 = 0 \\
 &\Rightarrow -2m + 3 = 0 \\
 &\Rightarrow -2m = -3 \\
 &\Rightarrow m = \frac{3}{2}
 \end{aligned}$$

$$\begin{aligned}
 12. \text{ Equal Roots } &\Rightarrow b^2 - 4(a)(1) = 0 \\
 &\Rightarrow b^2 - 4a = 0 \\
 &\Rightarrow -4a = -b^2 \\
 &\Rightarrow a = \frac{b^2}{4}
 \end{aligned}$$

$$\begin{aligned}
 \text{Hence, } x &= \frac{-b \pm \sqrt{0}}{2\left(\frac{b^2}{4}\right)} \\
 &= -b \cdot \frac{2}{b^2} = -\frac{2}{b}
 \end{aligned}$$

$$\begin{aligned}
 13. \text{ No Real Roots } &\Rightarrow (-2p)^2 - 4(1)(3p^2 + q^2) < 0 \\
 &\Rightarrow 4p^2 - 12p^2 - 4q^2 < 0 \\
 &\Rightarrow -8p^2 - 4q^2 < 0 \\
 &\Rightarrow -2p^2 - q^2 < 0 \quad \text{True}
 \end{aligned}$$

Exercises 2.3

$$\begin{aligned}
 1. \quad y = -2x + 3 \cap y = x^2 \\
 &\Rightarrow x^2 = -2x + 3 \\
 &\Rightarrow x^2 + 2x - 3 = 0 \\
 &\Rightarrow (x + 3)(x - 1) = 0 \\
 &\Rightarrow x = -3, \quad x = 1 \\
 &\Rightarrow y = (-3)^2, y = (1)^2 \\
 &\Rightarrow y = 9, \quad y = 1 \\
 \text{ANS} &= (-3, 9), (1, 1)
 \end{aligned}$$

$$\begin{aligned}
 2. \quad x = y - 1 \cap x^2 + y^2 = 5 \\
 &\Rightarrow (y - 1)^2 + y^2 - 5 = 0 \\
 &\Rightarrow y^2 - 2y + 1 + y^2 - 5 = 0 \\
 &\Rightarrow 2y^2 - 2y - 4 = 0 \\
 &\Rightarrow y^2 - y - 2 = 0 \\
 &\Rightarrow (y + 1)(y - 2) = 0 \\
 &\Rightarrow y = -1, \quad y = 2 \\
 &\Rightarrow x = -1 - 1, x = 2 - 1 \\
 &\Rightarrow x = -2, x = 1 \\
 \text{ANS} &= (-2, -1), (1, 2)
 \end{aligned}$$

$$\begin{aligned}
 3. \quad y = -2x + 2 \cap 4x^2 = y \\
 &\Rightarrow 4x^2 = -2x + 2 \\
 &\Rightarrow 4x^2 + 2x - 2 = 0 \\
 &\Rightarrow 2x^2 + x - 1 = 0 \\
 &\Rightarrow (x + 1)(2x - 1) = 0 \\
 &\Rightarrow x = -1, x = \frac{1}{2} \\
 &\Rightarrow y = -2(-1) + 2, y = -2\left(\frac{1}{2}\right) + 2 \\
 &\Rightarrow y = 4, \quad y = 1 \\
 \text{ANS} &= (-1, 4), \left(\frac{1}{2}, 1\right)
 \end{aligned}$$

$$\begin{aligned}
 4. \quad y = -x + 1 \cap y = x^2 - 6x + 5 \\
 &\Rightarrow x^2 - 6x + 5 = -x + 1 \\
 &\Rightarrow x^2 - 5x + 4 = 0 \\
 &\Rightarrow (x - 1)(x - 4) = 0 \\
 &\Rightarrow x = 1, \quad x = 4 \\
 &\Rightarrow y = -1 + 1, \quad y = -4 + 1 \\
 &\Rightarrow y = 0, \quad y = -3 \\
 \text{ANS} &= (1, 0), (4, -3)
 \end{aligned}$$

$$\begin{aligned}
 5. \quad & y = -x + 7 \cap x^2 + y^2 = 25 \\
 & \Rightarrow x^2 + (x + 7)^2 - 25 = 0 \\
 & \Rightarrow x^2 + x^2 - 14x + 49 - 25 = 0 \\
 & \Rightarrow 2x^2 - 14x + 24 = 0 \\
 & \Rightarrow x^2 - 7x + 12 = 0 \\
 & \Rightarrow (x - 3)(x - 4) = 0 \\
 & \Rightarrow x = 3, x = 4 \\
 & \Rightarrow y = -3 + 7, y = -4 + 7 \\
 & \Rightarrow y = 4, y = 3 \\
 & \text{ANS} = (3, 4), (4, 3)
 \end{aligned}$$

$$\begin{aligned}
 6. \quad & 2x + 1 = y \cap 3x^2 - y^2 = 3 \\
 & \Rightarrow 3x^2 - (2x + 1)^2 - 3 = 0 \\
 & \Rightarrow 3x^2 - 4x^2 - 4x - 1 - 3 = 0 \\
 & \Rightarrow -x^2 - 4x - 4 = 0 \\
 & \Rightarrow x^2 + 4x + 4 = 0 \\
 & \Rightarrow (x + 2)(x + 2) = 0 \\
 & \Rightarrow x = -2 \\
 & \Rightarrow y = 2(-2) + 1 = -3 \\
 & \text{ANS} = (-2, -3)
 \end{aligned}$$

$$\begin{aligned}
 7. \quad & y = 3x - 4 \cap y = x^2 - 4x + 6 \\
 & \Rightarrow x^2 - 4x + 6 = 3x - 4 \\
 & \Rightarrow x^2 - 7x + 10 = 0 \\
 & \Rightarrow (x - 2)(x - 5) = 0 \\
 & \Rightarrow x = 2, x = 5 \\
 & \Rightarrow y = 3(2) - 4, y = 3(5) - 4 \\
 & \Rightarrow y = 2, y = 11 \\
 & \text{ANS} = (2, 2), (5, 11)
 \end{aligned}$$

$$\begin{aligned}
 8. \quad & y = -x + 4 \cap x^2 + y^2 - 4x + 2 = 0 \\
 & \Rightarrow x^2 + (-x + 4)^2 - 4x + 2 = 0 \\
 & \Rightarrow x^2 + x^2 - 8x + 16 - 4x + 2 = 0 \\
 & \Rightarrow 2x^2 - 12x + 18 = 0 \\
 & \Rightarrow x^2 - 6x + 9 = 0 \\
 & \Rightarrow (x - 3)(x - 3) = 0 \\
 & \Rightarrow x = 3 \\
 & \Rightarrow y = -3 + 4 = 1 \\
 & \text{ANS} = (3, 1)
 \end{aligned}$$

$$\begin{aligned}
 9. \quad & x = -2y + 2 \cap x^2 + 4y^2 = 4 \\
 & \Rightarrow (-2y + 2)^2 + 4y^2 - 4 = 0 \\
 & \Rightarrow 4y^2 - 8y + 4 + 4y^2 - 4 = 0 \\
 & \Rightarrow 8y^2 - 8y = 0 \\
 & \Rightarrow y^2 - y = 0 \\
 & \Rightarrow y(y - 1) = 0 \\
 & \Rightarrow y = 0, y = 1 \\
 & \Rightarrow x = -2(0) + 2, x = -2(1) + 2 \\
 & \Rightarrow x = 2, x = 0 \\
 & \text{ANS} = (2, 0), (0, 1)
 \end{aligned}$$

$$\begin{aligned}
 10. \quad & 2x + 2 = y \cap xy = 4 \\
 & \Rightarrow x(2x + 2) = 4 \\
 & \Rightarrow 2x^2 + 2x - 4 = 0 \\
 & \Rightarrow x^2 + x - 2 = 0 \\
 & \Rightarrow (x + 2)(x - 1) = 0 \\
 & \Rightarrow x = -2, x = 1 \\
 & \Rightarrow y = 2(-2) + 2, y = 2(1) + 2 \\
 & \Rightarrow y = -2, y = 4 \\
 & \text{ANS} = (-2, -2), (1, 4)
 \end{aligned}$$

$$\begin{aligned}
 11. \quad & y = -2x + 3 \cap y^2 + xy = 2 \\
 & \Rightarrow (-2x + 3)^2 + x(-2x + 3) - 2 = 0 \\
 & \Rightarrow 4x^2 - 12x + 9 - 2x^2 + 3x - 2 = 0 \\
 & \Rightarrow 2x^2 - 9x + 7 = 0 \\
 & \Rightarrow (2x - 7)(x - 1) = 0 \\
 & \Rightarrow x = \frac{7}{2}, x = 1 \\
 & \Rightarrow y = -2\left(\frac{7}{2}\right) + 3, y = -2(1) + 3 \\
 & \Rightarrow y = -4, y = 1 \\
 & \text{ANS} = \left(\frac{7}{2}, -4\right), (1, 1)
 \end{aligned}$$

$$\begin{aligned}
 12. \quad & x = y - 3 \cap x^2 + y^2 + 2x - 4y + 3 = 0 \\
 & \Rightarrow (y - 3)^2 + y^2 + 2(y - 3) - 4y + 3 = 0 \\
 & \Rightarrow y^2 - 6y + 9 + y^2 + 2y - 6 - 4y + 3 = 0 \\
 & \Rightarrow 2y^2 - 8y + 6 = 0 \\
 & \Rightarrow y^2 - 4y + 3 = 0 \\
 & \Rightarrow (y - 1)(y - 3) = 0 \\
 & \Rightarrow y = 1, y = 3 \\
 & \Rightarrow x = 1 - 3, x = 3 - 3 \\
 & \Rightarrow x = -2, x = 0 \\
 & \text{ANS} = (-2, 1), (0, 3)
 \end{aligned}$$

$$\begin{aligned}
 13. \quad & s = 2t - 1 \cap 3t^2 - 2ts + s^2 = 9 \\
 & \Rightarrow 3t^2 - 2t(2t - 1) + (2t - 1)^2 = 9 \\
 & \Rightarrow 3t^2 - 4t^2 + 2t + 4t^2 - 4t + 1 - 9 = 0 \\
 & \Rightarrow 3t^2 - 2t - 8 = 0 \\
 & \Rightarrow (3t + 4)(t - 2) = 0 \\
 & \Rightarrow t = -\frac{4}{3}, t = 2 \\
 & \Rightarrow s = 2\left(-\frac{4}{3}\right) - 1, s = 2(2) - 1 \\
 & \Rightarrow s = -\frac{11}{3}, s = 3 \\
 & \text{ANS} = \left(-\frac{11}{3}, -\frac{4}{3}\right), (3, 2)
 \end{aligned}$$

$$\begin{aligned}
14. \quad & 2s + 3 = t \cap 2s^2 = t^2 + 1 \\
& \Rightarrow 2s^2 = (2s + 3)^2 + 1 \\
& \Rightarrow 2s^2 = 4s^2 + 12s + 9 + 1 \\
& \Rightarrow 2s^2 + 12s + 10 = 0 \\
& \Rightarrow s^2 + 6s + 5 = 0 \\
& \Rightarrow (s + 1)(s + 5) = 0 \\
& \Rightarrow s = -1, s = -5 \\
& \Rightarrow t = 2(-1) + 3, t = 2(-5) + 3 \\
& \Rightarrow t = 1, t = -7 \\
& \text{ANS} = (-1, 1), (-5, -7)
\end{aligned}$$

$$\begin{aligned}
15. \quad & 2t = 3s + 1 \cap t^2 + ts - 4s^2 = 2 \\
& \Rightarrow t = \frac{3s + 1}{2} \Rightarrow \left(\frac{3s + 1}{2}\right)^2 + s\left(\frac{3s + 1}{2}\right) - 4s^2 - 2 = 0 \\
& \Rightarrow \frac{9s^2 + 6s + 1}{2} + \frac{3s^2 + s}{2} - 4s^2 - 2 = 0 \\
& \Rightarrow 9s^2 + 6s + 1 + 3s^2 + s - 8s^2 - 4 = 0 \\
& \Rightarrow s^2 + 7s - 3 = 0 \\
& \Rightarrow (s - 1)(s + 3) = 0 \\
& \Rightarrow s = 1, s = -3 \\
& \Rightarrow t = \frac{3(1) + 1}{2}, t = \frac{3(-3) + 1}{2} \\
& \Rightarrow t = 2, t = -4 \\
& \text{ANS} = (1, 2), (-3, -4)
\end{aligned}$$

Exercise 2.4

$$\begin{aligned}
1. \quad & x, x + 1 \Rightarrow (x)^2 + (x + 1)^2 = 61 \\
& \Rightarrow x^2 + x^2 + 2x + 1 = 61 \\
& \Rightarrow 2x^2 + 2x - 60 = 0 \\
& \Rightarrow x^2 + x - 30 = 0 \\
& \Rightarrow (x + 6)(x - 5) = 0 \\
& \Rightarrow x = -6, x = 5 \\
& \Rightarrow x + 1 = -5, x + 1 = 6
\end{aligned}$$

Numbers are $-6, -5$ or $5, 6$

$$\begin{aligned}
2. \quad & x, x + 2 \Rightarrow (x)^2 + (x + 2)^2 = 52 \\
& \Rightarrow x^2 + x^2 + 4x + 4 = 52 \\
& \Rightarrow 2x^2 + 4x - 48 = 0 \\
& \Rightarrow x^2 + 2x - 24 = 0 \\
& \Rightarrow (x + 6)(x - 4) = 0 \\
& \Rightarrow x = -6, x = 4 \\
& \Rightarrow x + 2 = -4, x + 2 = 6
\end{aligned}$$

Numbers are $-6, -4$ or $4, 6$

$$\begin{aligned}
3. \quad & \text{(i) Perimeter} = 2x + 2y = 62 \\
& \Rightarrow x + y = 31 \\
& \Rightarrow y = -x + 31 \\
& \text{Area} = (x)(y) = 198 \\
& \Rightarrow xy = 198 \\
& \text{(ii) } \Rightarrow x(-x + 31) - 198 = 0 \\
& \Rightarrow -x^2 + 31x - 198 = 0 \\
& \Rightarrow x^2 - 31x + 198 = 0 \\
& \Rightarrow (x - 9)(x - 22) = 0 \\
& \Rightarrow x = 9 \text{ m}, x = 22 \text{ m} \\
& \Rightarrow y = -9 + 31, y = -22 + 31 \\
& \quad = 22 \text{ m}, \quad = 9 \text{ m} \\
& \text{Dimensions} = 9 \text{ m}, 22 \text{ m}
\end{aligned}$$

4. Sides = $x, x + 1, x + 2 \Rightarrow$ Pythagoras $\Rightarrow (x + 2)^2 = (x)^2 + (x + 1)^2$
 $\Rightarrow x^2 + 4x + 4 = x^2 + x^2 + 2x + 1$
 $\Rightarrow x^2 - 2x - 3 = 0$
 $\Rightarrow (x - 3)(x + 1) = 0$
 $\Rightarrow x = 3, x = -1$ (Not valid)
 $\Rightarrow x + 1 = 4$
 $\Rightarrow x + 2 = 5$


Perimeter = $3 + 4 + 5 = 12$ units

5. $s = 25 \Rightarrow 12t - t^2 = 25$
 $\Rightarrow t^2 - 12t + 25 = 0$
 $\Rightarrow t = \frac{12 \pm \sqrt{(-12)^2 - 4(1)(25)}}{2(1)}$
 $= \frac{12 \pm \sqrt{144 - 100}}{2}$
 $= \frac{12 \pm \sqrt{44}}{2}$
 $= \frac{12 + 6.633}{2}, \frac{12 - 6.633}{2}$
 $= 9.316, 2.683$
 $= 9.32, 2.68$

6. $x^2 - 15 = 2x$
 $\Rightarrow x^2 - 2x - 15 = 0$
 $\Rightarrow (x + 3)(x - 5) = 0$
 $\Rightarrow x = -3, x = 5$

7. $h = 6 \Rightarrow -16t^2 + 24t + 1 = 6$
 $\Rightarrow 16t^2 - 24t + 5 = 0$
 $\Rightarrow (4t - 1)(4t - 5) = 0$
 $\Rightarrow 4t = 1, 4t = 5$
 $\Rightarrow t = \frac{1}{4} \text{ sec}, t = \frac{5}{4} \text{ sec}$


8. Pythagoras $\Rightarrow (x)^2 + (x + 4)^2 = (20)^2$
 $\Rightarrow x^2 + x^2 + 8x + 16 = 400$
 $\Rightarrow 2x^2 + 8x - 384 = 0$
 $\Rightarrow x^2 + 4x - 192 = 0$
 $\Rightarrow (x - 12)(x + 16) = 0$
 $\Rightarrow x = 12, x = -16$ (Not valid)



9. $x, x + 2 \Rightarrow x(x + 2) = 4(2x + 2) - 1$
 $\Rightarrow x^2 + 2x = 8x + 8 - 1$
 $\Rightarrow x^2 - 6x - 7 = 0$
 $\Rightarrow (x + 1)(x - 7) = 0$
 $\Rightarrow x = -1, x = 7$
 $\Rightarrow x + 2 = 1, x + 2 = 9$

ANS = $-1, 1$ or $7, 9$

10. Pythagoras $\Rightarrow (x)^2 + (x + 3)^2 = (x + 6)^2$
 $\Rightarrow x^2 + x^2 + 6x + 9 = x^2 + 12x + 36$
 $\Rightarrow x^2 - 6x - 27 = 0$
 $\Rightarrow (x - 9)(x + 3) = 0$
 $\Rightarrow x = 9, x = -3$ (Not valid)




$$\begin{aligned}
 11. \quad x, x + 4 &\Rightarrow \text{Area} = x(x + 4) = 60 \\
 &\Rightarrow x^2 + 4x - 60 = 0 \\
 &\Rightarrow (x - 6)(x + 10) = 0 \\
 &\Rightarrow x = 6, x = -10 \text{ (Not valid)}
 \end{aligned}$$

Dimensions are 6 m, 10 m

$$\begin{aligned}
 12. \quad x, x + 1, x + 2 &\Rightarrow 3(x + x + 1 + x + 2) = (x + 1)(x + 2) \\
 &\Rightarrow 9x + 9 = x^2 + 3x + 2 \\
 &\Rightarrow x^2 - 6x - 7 = 0 \\
 &\Rightarrow (x + 1)(x - 7) = 0 \\
 &\Rightarrow x = -1, x = 7
 \end{aligned}$$


ANS = -1, 0, 1 or 7, 8, 9

$$\begin{aligned}
 13. \quad \text{Area} &= \pi(x + 14)^2 - \pi(14)^2 = 60\pi \\
 &\Rightarrow x^2 + 28x + 196 - 196 = 60 \\
 &\Rightarrow x^2 + 28x - 60 = 0 \\
 &\Rightarrow (x - 2)(x + 30) = 0 \\
 &\Rightarrow x = 2, x = -30 \text{ (Not valid)} \\
 &\text{width of desk} = 2 \text{ m}
 \end{aligned}$$



$$\begin{aligned}
 14. \quad \text{Area} &= x^2 + 96 = (2x)(x - 2) \\
 &\Rightarrow x^2 + 96 = 2x^2 - 4x \\
 &\Rightarrow x^2 - 4x - 96 = 0 \\
 &\Rightarrow (x - 12)(x + 8) = 0 \\
 &\Rightarrow x = 12, x = -8 \text{ (Not valid)}
 \end{aligned}$$

Dimensions = 10 m, 24 m



$$\begin{aligned}
 15. \quad \text{Point G} &\Rightarrow 0.1x^2 - x + 2.5 = 1.5 \\
 &\Rightarrow 0.1x^2 - x + 1 = 0 \\
 &\Rightarrow x^2 - 10x + 10 = 0 \\
 &\Rightarrow x = \frac{10 \pm \sqrt{100 - 4(1)(10)}}{2(1)} \\
 &= \frac{10 \pm \sqrt{60}}{2} \\
 &= \frac{10 - \sqrt{60}}{2}, \frac{10 + \sqrt{60}}{2} \\
 &= 1.127, 8.873 \text{ (Not valid)}
 \end{aligned}$$

$$\begin{aligned}
 \text{Point C} &\Rightarrow 0.1x^2 - x + 2.5 = 3 \\
 &\Rightarrow 0.1x^2 - x - 0.5 = 0 \\
 &\Rightarrow x^2 - 10x - 5 = 0 \\
 &\Rightarrow x = \frac{10 \pm \sqrt{100 - 4(1)(-5)}}{2(1)} \\
 &= \frac{10 \pm \sqrt{120}}{2} \\
 &= \frac{10 + \sqrt{120}}{2}, \frac{10 - \sqrt{120}}{2} \\
 &= 10.477, -0.477 \text{ (Not valid)}
 \end{aligned}$$

Distance = $10.477 - 1.127 = 9.35 \text{ m}$

$$\begin{aligned}
 16. \quad 3t - 4 = s \cap 2t^2 + s^2 &= 43 \\
 \Rightarrow 2t^2 + (3t - 4)^2 - 43 &= 0 \\
 \Rightarrow 2t^2 + 9t^2 - 24t + 16 - 43 &= 0 \\
 \Rightarrow 11t^2 - 24t - 27 &= 0 \\
 \Rightarrow (t - 3)(11t + 9) &= 0 \\
 \Rightarrow t = 3, t = -\frac{9}{11} & \text{ (Not valid)} \\
 \Rightarrow s = 3(3) - 4 & \\
 &= 5
 \end{aligned}$$

Point = (3, 5)

$$\begin{aligned}
 17. \quad x + 3y = 5 \cap x^2 + 6y^2 &= 40 \\
 \Rightarrow x = -3y + 5 \Rightarrow (-3y + 5)^2 + 6y^2 - 40 &= 0 \\
 \Rightarrow 9y^2 - 30y + 25 + 6y^2 - 40 &= 0 \\
 \Rightarrow 15y^2 - 30y - 15 &= 0 \\
 \Rightarrow y^2 - 2y - 1 &= 0 \\
 \Rightarrow y = \frac{2 \pm \sqrt{4 - 4(1)(-1)}}{2(1)} & \\
 \Rightarrow y = \frac{2 \pm \sqrt{8}}{2} & \\
 \Rightarrow y = 1 + \sqrt{2}, 1 - \sqrt{2} & \\
 \Rightarrow y = 2.414, -0.414 & \\
 \Rightarrow x = -3(2.414) + 5, x = -3(-0.414) + 5 & \\
 &= -2.242, \quad = 6.242
 \end{aligned}$$

Plane crosses front at 2 points; (-2.2, 2.4) and (6.2, -0.4)

$$\begin{aligned}
 k = 10 \Rightarrow x + 3y = 10 \cap x^2 + 6y^2 &= 40 \\
 \Rightarrow x = -3y + 10 \Rightarrow (-3y + 10)^2 + 6y^2 - 40 &= 0 \\
 \Rightarrow 15y^2 - 60y + 60 &= 0 \\
 \Rightarrow y^2 - 4y + 4 &= 0 \\
 \Rightarrow (y - 2)(y - 2) &= 0 \\
 \Rightarrow y = 2 \text{ and } x = -3(2) + 10 &= 4
 \end{aligned}$$

Plane crosses front at one point; (4, 2)

\Rightarrow Plane will avoid weather front when $k > 10$

\Rightarrow Minimum value of $k = 11$

Exercise 2.5

1. (a) (i) sum = -9 (ii) product = 4
- (b) (i) sum = 2 (ii) product = -5
- (c) (i) sum = 7 (ii) product = 2
- (d) (i) sum = 9 (ii) product = -3
- (e) (i) sum = $\frac{7}{2}$ (ii) product = $-\frac{1}{2}$
- (f) (i) sum = $-\frac{1}{7}$ (iii) product = $-\frac{1}{7}$
- (g) (i) sum = $-\frac{10}{3}$ (ii) product = $-\frac{2}{3}$
- (h) (i) sum = $-\frac{10}{5} = -2$ (ii) product = $\frac{1}{5}$
- (i) (i) sum = -2 (ii) product = -3
- (j) (i) sum = $\frac{3}{4}$ (ii) product = $\frac{5}{4}$

2. (a) $x^2 + 3x - 1 = 0$

(b) $x^2 - 6x - 4 = 0$

(c) $x^2 - 7x - 5 = 0$

(d) $x^2 + \frac{2}{3}x - \frac{7}{3} = 0$

$$\Rightarrow 3x^2 + 2x - 7 = 0$$

(e) $x^2 + \frac{5}{2}x - 2 = 0$

$$\Rightarrow 2x^2 + 5x - 4 = 0$$

3. (i) $x^2 - (4 + 6)x + (4)(6) = 0$

$$\Rightarrow x^2 - 10x + 24 = 0$$

(ii) $x^2 - (2 - 3)x + (2)(-3) = 0$

$$\Rightarrow x^2 + x - 6 = 0$$

(iii) $x^2 - (-5 - 1)x + (-5)(-1) = 0$

$$\Rightarrow x^2 + 6x + 5 = 0$$

(iv) $x^2 - (4 + \sqrt{5})x + (4)(\sqrt{5}) = 0$

$$\Rightarrow x^2 - (4 + \sqrt{5})x + 4\sqrt{5} = 0$$

(v) $x^2 - (a + 3a)x + (a)(3a) = 0$

$$\Rightarrow x^2 - 4ax + 3a^2 = 0$$

(vi) $x^2 - \left(\frac{2}{5} + \frac{3}{5}\right)x + \left(\frac{2}{5}\right)\left(\frac{3}{5}\right) = 0$

$$\Rightarrow x^2 - x + \frac{6}{25} = 0$$

$$\Rightarrow 25x^2 - 25x + 6 = 0$$

(f) $x^2 + \frac{3}{2}x - 5 = 0$

$$\Rightarrow 2x^2 + 3x - 10 = 0$$

(g) $x^2 + \frac{1}{4}x - \frac{1}{3} = 0$

$$\Rightarrow 12x^2 + 3x - 4 = 0$$

(h) $x^2 + 1\frac{2}{3}x + \frac{1}{2} = 0$

$$\Rightarrow 6x^2 + 10x + 3 = 0$$

(vii) $x^2 - \left(\frac{2}{b} + \frac{3}{b}\right)x + \left(\frac{2}{b}\right)\left(\frac{3}{b}\right) = 0$

$$\Rightarrow x^2 - \frac{5}{b}x + \frac{6}{b^2} = 0$$

$$\Rightarrow b^2x^2 - 5bx + 6 = 0$$

(viii) $x^2 - \left(\frac{5}{2} + \frac{3}{5}\right)x + \left(\frac{5}{2}\right)\left(\frac{3}{5}\right) = 0$

$$\Rightarrow x^2 - \frac{31}{10}x + \frac{3}{2} = 0$$

$$\Rightarrow 10x^2 - 31x + 15 = 0$$

Exercise 2.6

1. (i) $c = \left(\frac{28}{2}\right)^2 = (14)^2 = 196$ (ii) $c = \left(\frac{-6}{2}\right)^2 = (-3)^2 = 9$ (iii) $c = \left(\frac{-5}{2}\right)^2 = \frac{25}{4}$

2. (i) $x^2 - 8x - 3 = 0$

$$\Rightarrow x^2 - 8x + 16 - 16 - 3 = 0$$

$$\Rightarrow (x - 4)^2 - 19 = 0$$

(ii) $x^2 - 2x - 5 = 0$

$$\Rightarrow x^2 - 2x + 1 - 1 - 5 = 0$$

$$\Rightarrow (x - 1)^2 - 6 = 0$$

(iii) $x^2 - 2x + 1 = 0$

$$\Rightarrow (x - 1)^2 = 0$$

3. (i) $x^2 + 4x - 6 = 0$

$$\Rightarrow x^2 + 4x + 4 - 4 - 6 = 0$$

$$\Rightarrow (x + 2)^2 - 10 = 0$$

(ii) $x^2 + 9x + 4 = 0$

$$\Rightarrow x^2 + 9x + 20\frac{1}{4} - 20\frac{1}{4} + 4 = 0$$

$$\Rightarrow \left(x + 4\frac{1}{2}\right)^2 - 16\frac{1}{4} = 0$$

(iii) $x^2 - 7x - 3 = 0$

$$\Rightarrow x^2 - 7x + 12\frac{1}{4} - 12\frac{1}{4} - 3 = 0$$

$$\Rightarrow \left(x - 3\frac{1}{2}\right)^2 - 15\frac{1}{4} = 0$$

4. (i) $2x^2 + 4x - 5 = 0$

$$= 2\left(x^2 + 2x - \frac{5}{2}\right) = 0$$

$$= 2\left(x^2 + 2x + 1 - 1 - \frac{5}{2}\right) = 0$$

$$= 2\left[(x+1)^2 - \frac{7}{2}\right] = 0$$

$$= 2(x+1)^2 - 7 = 0$$

$$\Rightarrow \text{Minimum Point} = (-1, -7)$$

(ii) $3x^2 - 6x - 1 = 0$

$$\Rightarrow 3\left(x^2 - 2x - \frac{1}{3}\right) = 0$$

$$\Rightarrow 3\left(x^2 - 2x + 1 - 1 - \frac{1}{3}\right) = 0$$

$$\Rightarrow 3\left[(x-1)^2 - 1\frac{1}{3}\right] = 0$$

$$\Rightarrow 3(x-1)^2 - 4 = 0$$

$$\Rightarrow \text{Minimum Point} = (1, -4)$$

(iii) $4x^2 + x + 3 = 0$

$$\Rightarrow 4\left(x^2 + \frac{1}{4}x + \frac{3}{4}\right) = 0$$

$$\Rightarrow 4\left(x^2 + \frac{1}{4}x + \frac{1}{64} - \frac{1}{64} + \frac{3}{4}\right) = 0$$

$$\Rightarrow 4\left[\left(x + \frac{1}{8}\right)^2 + \frac{47}{64}\right] = 0$$

$$\Rightarrow 4\left(x + \frac{1}{8}\right)^2 + \frac{47}{16} = 0$$

$$\Rightarrow \text{Minimum Point} = \left(-\frac{1}{8}, \frac{47}{16}\right)$$

5. $x^2 - 6x + k = 0$

$$= x^2 - 6x + 9 - 9 + k$$

$$= (x-3)^2 + k - 9$$

$$\text{hence, } k - 9 > 0$$

$$\Rightarrow k > 9$$

$$\text{Minimum value of } k = 10$$

6. $2x^2 - 12x + 7$

$$= 2\left(x^2 - 6x + \frac{7}{2}\right)$$

$$= 2\left(x^2 - 6x + 9 - 9 + \frac{7}{2}\right)$$

$$= 2\left[(x-3)^2 - \frac{11}{2}\right]$$

$$= 2(x-3)^2 - 11$$

7. $g(x) = x^2 + 8x + 20$

$$= x^2 + 8x + 16 - 16 + 20$$

$$= (x+4)^2 + 4$$

$$\text{Since } (x+4)^2 \geq 0$$

$$\Rightarrow g(x) = (x+4)^2 + 4 \geq 4$$

8. (i) Minimum for $f(x) = (-1, -5)$

$$\Rightarrow f(x) = (x+1)^2 - 5$$

$$= x^2 + 2x + 1 - 5$$

$$= x^2 + 2x - 4$$

$$\text{Choose } (-2, -4) \Rightarrow f(-2) = (-2)^2 + 2(-2) - 4$$

$$= 4 - 4 - 4$$

$$= -4 \quad \text{True}$$

$$\text{Minimum for } g(x) = (2, -1)$$

$$\Rightarrow g(x) = (x-2)^2 - 1$$

$$= x^2 - 4x + 4 - 1$$

$$= x^2 - 4x + 3$$

$$\begin{aligned}\text{Choose } (3, 0) \Rightarrow g(3) &= (3)^2 - 4(3) + 3 \\ &= 9 - 12 + 3 \\ &= 0 \quad \text{True}\end{aligned}$$

$$\begin{aligned}\text{Minimum for } h(x) &= (4, 1) \\ \Rightarrow h(x) &= (x - 4)^2 + 1 \\ &= x^2 - 8x + 16 + 1 \\ &= x^2 - 8x + 17\end{aligned}$$

$$\begin{aligned}\text{Choose } (5, 2) \Rightarrow h(5) &= (5)^2 - 8(5) + 17 \\ &= 25 - 40 + 17 \\ &= 42 - 40 = 2 \quad \text{True}\end{aligned}$$


$$\begin{aligned}9. \quad f(x) &= x^2 + 4x + 7 \\ &= x^2 + 4x + 4 + 3 \\ &= (x + 2)^2 + 3\end{aligned}$$

(i) smallest value of $f(x) = 3$ (ii) $x = -2$

$$(iii) \frac{1}{(x^2 + 4x + 7)} = \frac{1}{3} \text{ (greatest value)}$$

$$\begin{aligned}10. \quad y &= -x^2 + 6x \\ &= -(x^2 - 6x) \\ &= -(x^2 - 6x + 9 - 9) \\ &= -[(x - 3)^2 - 9] \\ &= -(x - 3)^2 + 9\end{aligned}$$

maximum point = (3, 9)
greatest height = 9



$$\begin{aligned}11. \quad (i) \quad C: x^2 - 6x + 8 \\ &= x^2 - 6x + 9 - 9 + 8 \\ &= (x - 3)^2 - 1\end{aligned}$$

$$\begin{aligned}(ii) \quad B: x^2 - 6x + 9 \\ &= (x - 3)^2\end{aligned}$$

$$\begin{aligned}(iii) \quad A: x^2 - 6x + 10 \\ &= x^2 - 6x + 9 - 9 + 10 \\ &= (x - 3)^2 + 1\end{aligned}$$

$$\begin{aligned}12. \quad \text{Curve C: Maximum point} &= (2, 4) = (p, q) \\ \Rightarrow y &= 4 - a(x - 2)^2 \\ \text{point } (0, 3) \Rightarrow 3 &= 4 - a(0 - 2)^2 \\ \Rightarrow 3 &= 4 - a(4) \\ \Rightarrow 4a &= 1 \\ \Rightarrow a &= \frac{1}{4}\end{aligned}$$

$$\begin{aligned}\text{Curve D: Maximum point} &= (2, 4) = (p, q) \\ \Rightarrow y &= 4 - a(x - 2)^2 \\ \text{point } (1, 3) \Rightarrow 3 &= 4 - a(1 - 2)^2 \\ \Rightarrow 3 &= 4 - a(1) \\ \Rightarrow a &= 1\end{aligned}$$

$$\begin{aligned}13. \quad \text{Roots are } x = 6, x = -3 \\ \text{Hence, } y &= a(x - 6)(x + 3) \\ \Rightarrow y &= a(x^2 - 3x - 18) \\ \text{point } (1, 10) \Rightarrow 10 &= a[(1)^2 - 3(1) - 18] \\ \Rightarrow 10 &= a(1 - 3 - 18) \\ \Rightarrow 10 &= -20a \\ \Rightarrow a &= -\frac{1}{2}\end{aligned}$$

$$\Rightarrow y = -\frac{1}{2}(x^2 - 3x - 18) = -\frac{1}{2}x^2 + \frac{3}{2}x + 9$$

- 14.** Minimum point = $(-1, 3)$

$$\Rightarrow y = a(x + 1)^2 + 3$$

$$\text{point } (0, 4) \Rightarrow 4 = a(0 + 1)^2 + 3$$

$$\Rightarrow 4 = a + 3$$

$$\Rightarrow a = 1$$

$$\Rightarrow y = 1(x + 1)^2 + 3 = x^2 + 2x + 4$$

- 15.** (i) Maximum point = $(6, 4)$

$$\Rightarrow f(x) = 4 - (0.1)(x - 6)^2$$

$$(ii) f(x) = 4 - (0.1)(x^2 - 12x + 36) = 0$$

$$\Rightarrow 40 - (x^2 - 12x + 36) = 0$$

$$\Rightarrow -x^2 + 12x + 4 = 0$$

$$\Rightarrow x = \frac{-12 \pm \sqrt{144 - 4(-1)(4)}}{2(-1)}$$

$$= \frac{-12 \pm \sqrt{160}}{-2}$$

$$= \frac{-12 \pm 4\sqrt{10}}{-2}$$

$$= 6 - 2\sqrt{10}, 6 + 2\sqrt{10}$$

point where ball started is $(6 - 2\sqrt{10}, 0)$

and point where ball finished is $(6 + 2\sqrt{10}, 0)$

$$(iii) \text{ Horizontal Distance} = 6 + 2\sqrt{10} - (6 - 2\sqrt{10})$$

$$= 6 + 2\sqrt{10} - 6 + 2\sqrt{10}$$

$$= 4\sqrt{10}$$

- 16.** (i) Area of trapezium = $\frac{1}{2}(a + b)h$

$$= \frac{1}{2}(x + 20)(2x)$$

$$= x(x + 20)$$

$$= x^2 + 20x$$

$$(ii) \text{ Given: } x^2 + 20x = 400$$

$$x^2 + 20x + 100 = 400 + 100$$

$$(x + 10)^2 = 500$$

$$x + 10 = \pm\sqrt{500}$$

$$x + 10 = \pm 10\sqrt{5}$$

$$x = -10 + 10\sqrt{5}$$

$$x = 10(\sqrt{5} - 1)$$

(as $x = -10 - 10\sqrt{5}$ is not possible)

- 17.** $T = 36 + 4t - t^2$

$$t^2 - 4t = -T + 36$$

$$t^2 - 4t + 4 = -T + 36 + 4$$

$$(t - 2)^2 = -T + 40$$

$$T = -(t - 2)^2 + 40$$

The maximum temperature is then 40° .

Then

$$T = 36$$

$$-(t - 2)^2 + 40 = 36$$

$$-(t - 2)^2 = -4$$

$$(t - 2)^2 = 4$$

$$(t - 2) = \pm\sqrt{4}$$

$$t = 2 \pm 2$$

$$t = 0 \quad \text{or} \quad t = 4$$

Thus it will take 4 hours for temperature returns to 36°

Exercise 2.7

1. (i) $\sqrt{8} = \sqrt{4} \cdot \sqrt{2} = 2\sqrt{2}$
 (ii) $\sqrt{27} = \sqrt{9} \cdot \sqrt{3} = 3\sqrt{3}$
 (iii) $\sqrt{45} = \sqrt{9} \cdot \sqrt{5} = 3\sqrt{5}$
2. (i) $2\sqrt{2} + 6\sqrt{2} - 3\sqrt{2} = 5\sqrt{2}$
 (ii) $2\sqrt{2} + \sqrt{18}$
 $= 2\sqrt{2} + \sqrt{9} \cdot \sqrt{2}$
 $= 2\sqrt{2} + 3\sqrt{2} = 5\sqrt{2}$
 (iii) $\sqrt{32} + \sqrt{18}$
 $= \sqrt{16} \cdot \sqrt{2} + \sqrt{9} \cdot \sqrt{2}$
 $= 4\sqrt{2} + 3\sqrt{2}$
 $= 7\sqrt{2}$
 (iv) $\sqrt{27} + \sqrt{48} - 2\sqrt{3}$
 $= \sqrt{9} \cdot \sqrt{3} + \sqrt{16} \cdot \sqrt{3} - 2\sqrt{3}$
 $= 3\sqrt{3} + 4\sqrt{3} - 2\sqrt{3} = 5\sqrt{3}$
3. (i) $\frac{1}{\sqrt{3}} = \frac{1}{\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{\sqrt{3}}{3}$
 (ii) $\frac{2}{\sqrt{8}} = \frac{2}{\sqrt{4} \cdot \sqrt{2}} = \frac{2}{2\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2}$
 (iii) $\frac{2}{5\sqrt{2}} = \frac{2}{5\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{2\sqrt{2}}{10} = \frac{\sqrt{2}}{5}$
 (iv) $\frac{20}{\sqrt{50}} = \frac{20}{\sqrt{25} \cdot \sqrt{2}} = \frac{20}{5\sqrt{2}} = \frac{4}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{4\sqrt{2}}{2} = 2\sqrt{2}$
 (v) $\frac{8}{\sqrt{128}} = \frac{8}{\sqrt{64} \cdot \sqrt{2}} = \frac{8}{8\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2}$
4. (i) $\sqrt{8} \times \sqrt{12} = \sqrt{96} = \sqrt{16} \sqrt{6} = 4\sqrt{6}$
 (ii) $3\sqrt{2} \times 5\sqrt{2} = 15 \cdot \sqrt{4} = 15 \cdot 2 = 30$
 (iii) $\sqrt{2}(\sqrt{6} + 3\sqrt{2}) = \sqrt{12} + 3\sqrt{4} = \sqrt{4} \sqrt{3} + 3 \cdot 2 = 2\sqrt{3} + 6$
 (iv) $(5 - \sqrt{3})(5 + \sqrt{3})$
 $= 5(5 + \sqrt{3}) - \sqrt{3}(5 + \sqrt{3})$
 $= 25 + 5\sqrt{3} - 5\sqrt{3} - \sqrt{9}$
 $= 25 - 3 = 22$
 (v) $(\sqrt{7} + \sqrt{5})(\sqrt{7} - \sqrt{5})$
 $= \sqrt{7}(\sqrt{7} - \sqrt{5}) + \sqrt{5}(\sqrt{7} - \sqrt{5})$
 $= \sqrt{49} - \sqrt{35} + \sqrt{35} - \sqrt{25} = 7 - 5 = 2$
 (vi) $(a + 2\sqrt{b})(a - 2\sqrt{b})$
 $= a(a - 2\sqrt{b}) + 2\sqrt{b}(a - 2\sqrt{b})$
 $= a^2 - 2a\sqrt{b} + 2a\sqrt{b} - 4\sqrt{b^2} = a^2 - 4b$
5. (i) $\frac{4}{\sqrt{5} + 1} \cdot \frac{\sqrt{5} - 1}{\sqrt{5} - 1} = \frac{4(\sqrt{5} - 1)}{\sqrt{25} - 1} = \frac{4(\sqrt{5} - 1)}{5 - 1} = \frac{4(\sqrt{5} - 1)}{4} = \sqrt{5} - 1$
 (ii) $\frac{12}{3 - \sqrt{2}} \cdot \frac{3 + \sqrt{2}}{3 + \sqrt{2}} = \frac{36 + 12\sqrt{2}}{9 - \sqrt{4}} = \frac{36 + 12\sqrt{2}}{7}$
 (iii) $\frac{2 - \sqrt{5}}{2 + \sqrt{5}} \cdot \frac{2 - \sqrt{5}}{2 - \sqrt{5}} = \frac{4 - 2\sqrt{5} - 2\sqrt{5} + \sqrt{25}}{4 - \sqrt{25}}$
 $= \frac{9 - 4\sqrt{5}}{-1} = -9 + 4\sqrt{5}$
 (iv) $\frac{1}{\sqrt{8} - \sqrt{2}} = \frac{1}{\sqrt{4} \sqrt{2} - \sqrt{2}} = \frac{1}{2\sqrt{2} - \sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{4}} = \frac{\sqrt{2}}{2}$

$$\begin{aligned}
 6. \quad (i) \quad \frac{1}{\sqrt{2}-1} - \frac{1}{\sqrt{2}+1} &= \frac{1(\sqrt{2}+1) - 1(\sqrt{2}-1)}{(\sqrt{2}-1)(\sqrt{2}+1)} \\
 &= \frac{\sqrt{2}+1 - \sqrt{2}+1}{\sqrt{4} + \sqrt{2} - \sqrt{2} - 1} = \frac{2}{2-1} = 2
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad \frac{1}{2+\sqrt{3}} + \frac{1}{2-\sqrt{3}} &= \frac{1(2-\sqrt{3}) + 1(2+\sqrt{3})}{(2+\sqrt{3})(2-\sqrt{3})} \\
 &= \frac{2-\sqrt{3} + 2+\sqrt{3}}{4-2\sqrt{3} + 2\sqrt{3} - \sqrt{9}} \\
 &= \frac{4}{4-3} = 4
 \end{aligned}$$

$$\begin{aligned}
 7. \quad (i) \quad (2\sqrt{3} - \sqrt{5})(2\sqrt{3} + \sqrt{5}) \\
 &= 2\sqrt{3}(2\sqrt{3} + \sqrt{5}) - \sqrt{5}(2\sqrt{3} + \sqrt{5}) \\
 &= 4.3 + 2\sqrt{15} - 2\sqrt{15} - \sqrt{25} = 12 - 5 = 7
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad \frac{4}{2-\sqrt{5}} + \frac{2}{2+\sqrt{5}} &= \frac{4(2+\sqrt{5}) + 2(2-\sqrt{5})}{(2-\sqrt{5})(2+\sqrt{5})} \\
 &= \frac{8 + 4\sqrt{5} + 4 - 2\sqrt{5}}{4 + 2\sqrt{5} - 2\sqrt{5} - \sqrt{25}} \\
 &= \frac{12 + 2\sqrt{5}}{-1} \\
 &= -12 - 2\sqrt{5}
 \end{aligned}$$

$$8. \quad (i) \quad X + Y = \frac{4+\sqrt{3}}{\sqrt{2}} + \frac{4-\sqrt{3}}{\sqrt{2}} = \frac{8}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{8\sqrt{2}}{2} = 4\sqrt{2}$$

$$\begin{aligned}
 (ii) \quad X - Y &= \frac{4+\sqrt{3}}{\sqrt{2}} - \frac{4-\sqrt{3}}{\sqrt{2}} = \frac{4+\sqrt{3} - 4 + \sqrt{3}}{\sqrt{2}} \\
 &= \frac{2\sqrt{3}}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{2\sqrt{6}}{2} = \sqrt{6}
 \end{aligned}$$

$$(iii) \quad XY = \frac{4+\sqrt{3}}{\sqrt{2}} \cdot \frac{4-\sqrt{3}}{\sqrt{2}} = \frac{16 - 4\sqrt{3} + 4\sqrt{3} - \sqrt{9}}{2} = \frac{16-3}{2} = \frac{13}{2}$$

$$\begin{aligned}
 (iv) \quad \frac{X}{Y} &= \frac{\frac{4+\sqrt{3}}{\sqrt{2}}}{\frac{4-\sqrt{3}}{\sqrt{2}}} = \frac{4+\sqrt{3}}{4-\sqrt{3}} \cdot \frac{4+\sqrt{3}}{4+\sqrt{3}} = \frac{16 + 4\sqrt{3} + 4\sqrt{3} + \sqrt{9}}{16 + 4\sqrt{3} - 4\sqrt{3} - \sqrt{9}} \\
 &= \frac{19 + 8\sqrt{3}}{13}
 \end{aligned}$$

$$\begin{aligned}
 9. \quad (2\sqrt{5} - 3\sqrt{2})(2\sqrt{5} + 3\sqrt{2}) \\
 &= 2\sqrt{5}(2\sqrt{5} + 3\sqrt{2}) - 3\sqrt{2}(2\sqrt{5} + 3\sqrt{2}) \\
 &= 4\sqrt{25} + 6\sqrt{10} - 6\sqrt{10} - 9\sqrt{4} \\
 &= 20 - 18 = 2
 \end{aligned}$$

$$\begin{aligned}
 10. \quad \frac{5}{2+\sqrt{3}} &= \frac{5}{2+\sqrt{3}} \cdot \frac{2-\sqrt{3}}{2-\sqrt{3}} \\
 &= \frac{5(2-\sqrt{3})}{4-3} = 5(2-\sqrt{3})
 \end{aligned}$$

$$\begin{aligned}
 11. \quad \frac{2+\sqrt{2}}{1+\sqrt{2}} \cdot \frac{1-\sqrt{2}}{1-\sqrt{2}} &= \frac{2-2\sqrt{2}+\sqrt{2}-2}{1+\sqrt{2}-\sqrt{2}-2} = \frac{-\sqrt{2}}{-1} = \sqrt{2} \\
 \frac{\sqrt{5}-2}{\sqrt{5}-1} \cdot \frac{\sqrt{5}+1}{\sqrt{5}+1} &= \frac{5+\sqrt{5}-2\sqrt{5}-2}{5+\sqrt{5}-\sqrt{5}-1} = \frac{3-\sqrt{5}}{4}
 \end{aligned}$$


$$\begin{aligned}
 \text{Hence, } \sqrt{2} \cdot \frac{(3-\sqrt{5})}{4} \cdot (3+\sqrt{5}) &= \frac{\sqrt{2}}{4} [3(3+\sqrt{5}) - \sqrt{5}(3+\sqrt{5})] \\
 &= \frac{\sqrt{2}}{4} [9 + 3\sqrt{5} - 3\sqrt{5} - 5] \\
 &= \frac{\sqrt{2}}{4} [4] = \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 12. \quad \frac{-1 + \sqrt{3}}{1 + \sqrt{3}} &= \frac{-1 + \sqrt{3}}{1 + \sqrt{3}} \times \frac{1 - \sqrt{3}}{1 - \sqrt{3}} \\
 &= \frac{-1 + \sqrt{3} + \sqrt{3} - 3}{(1)^2 - (\sqrt{3})^2} \\
 &= \frac{-4 + 2\sqrt{3}}{1 - 3} \\
 &= \frac{-4 + 2\sqrt{3}}{-2} \\
 &= 2 - \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 13. \quad \frac{\sqrt{3}}{1 - \sqrt{3}} - \frac{1}{\sqrt{3}} &= \frac{(\sqrt{3})^2 - (1 - \sqrt{3})}{\sqrt{3}(1 - \sqrt{3})} \\
 &= \frac{3 - 1 + \sqrt{3}}{\sqrt{3} - 3} \\
 &= \frac{2 + \sqrt{3}}{\sqrt{3} - 3} \times \frac{\sqrt{3} + 3}{\sqrt{3} + 3} \\
 &= \frac{2\sqrt{3} + 6 + 3 + 3\sqrt{3}}{(\sqrt{3})^2 - 9} \\
 &= \frac{9 + 5\sqrt{3}}{3 - 9} \\
 &= \frac{9 + 5\sqrt{3}}{-6} \\
 &= \frac{-9 - 5\sqrt{3}}{6}
 \end{aligned}$$

Exercise 2.8

$$\begin{aligned}
 1. \quad d^2 &= (x + 2)^2 + (x - 2)^2 \\
 &= x^2 + 4x + 4 + x^2 - 4x + 4 \\
 &= 2x^2 + 8 \\
 \Rightarrow d &= \sqrt{2x^2 + 8}
 \end{aligned}$$



$$\begin{aligned}
 2. \quad (a) \quad |AC|^2 &= (2 + \sqrt{3})^2 + (2 - \sqrt{3})^2 \\
 &= 4 + 4\sqrt{3} + 3 + 4 - 4\sqrt{3} - 3 \\
 &= 14 \\
 \Rightarrow |AC| &= \sqrt{14}
 \end{aligned}$$

$$(b) \quad (i) \quad \text{Distance, 1st Runner} = 2 - \sqrt{3} + 2 + \sqrt{3} + 2 - \sqrt{3} + 2 + \sqrt{3} = 8$$

$$\text{Distance, 2nd Runner} = 2\sqrt{14}$$

$$\Rightarrow \text{Difference} = 8 - 2\sqrt{14} = 2(4 - \sqrt{14}) \text{ km}$$

$$(ii) \quad \text{Time, 1st Runner} = \frac{8000}{1.5} = 5333.33$$

$$\text{Time, 2nd Runner} = \frac{2\sqrt{14} \times 1000}{1.4} = 5345.22$$

$$\Rightarrow \text{Difference} = 5345.22 - 5333.33 = 11.89 = 12 \text{ secs}$$

$$3. \quad |EG|^2 = (4)^2 + (2)^2 = 16 + 4 = 20$$

$$\Rightarrow |EG| = \sqrt{20}$$

$$|KG|^2 = (\sqrt{20})^2 + (2)^2 = 20 + 4 = 24$$

$$\Rightarrow |KG| = \sqrt{24} = \sqrt{4}\sqrt{6} = 2\sqrt{6}$$

$$\text{Hence: Distance} = 4 + 2 + 2 + 2\sqrt{6} = 8 + 2\sqrt{6}$$

4. (i) $x + y = \sqrt{a} + \frac{1}{\sqrt{a}} + \sqrt{a} - \frac{1}{\sqrt{a}} = 2\sqrt{a}$

(ii) $x - y = \sqrt{a} + \frac{1}{\sqrt{a}} - \left(\sqrt{a} - \frac{1}{\sqrt{a}} \right)$
 $= \sqrt{a} + \frac{1}{\sqrt{a}} - \sqrt{a} + \frac{1}{\sqrt{a}} = \frac{2}{\sqrt{a}}$

Hence, $\sqrt{x^2 - y^2} = \sqrt{(x + y)(x - y)}$
 $= \sqrt{2\sqrt{a} \cdot \frac{2}{\sqrt{a}}} = \sqrt{4} = 2$

5. (i) $\sqrt{2x + 1} = 3$
 $\Rightarrow 2x + 1 = 9$
 $\Rightarrow 2x = 8$
 $\Rightarrow x = 4$

Check $x = 4 \Rightarrow \sqrt{2(4) + 1} = \sqrt{9} = 3$ True

(ii) $\sqrt{3x + 10} = x$
 $\Rightarrow 3x + 10 = x^2$
 $\Rightarrow x^2 - 3x - 10 = 0$
 $\Rightarrow (x + 2)(x - 5) = 0$
 $\Rightarrow x = -2, x = 5$

Check $x = -2 \Rightarrow \sqrt{3(-2) + 10} = \sqrt{4} = 2 \neq -2$
 False

Check $x = 5 \Rightarrow \sqrt{3(5) + 10} = \sqrt{25} = 5$
 True

(iii) $\sqrt{2x - 1} = \sqrt{x + 8}$
 $\Rightarrow 2x - 1 = x + 8$
 $\Rightarrow x = 9$

Check $x = 9 \Rightarrow \sqrt{2(9) - 1} = \sqrt{9 + 8}$
 $\Rightarrow \sqrt{17} = \sqrt{17}$ True

(iv) $\sqrt{3x - 5} = x - 1$
 $\Rightarrow (\sqrt{3x - 5})^2 = (x - 1)^2$
 $\Rightarrow 3x - 5 = x^2 - 2x + 1$
 $\Rightarrow x^2 - 5x + 6 = 0$
 $\Rightarrow (x - 2)(x - 3) = 0$
 $\Rightarrow x = 2, x = 3$

Check $x = 2 \Rightarrow \sqrt{3(2) - 5} = 2 - 1$
 $\Rightarrow 1 = 1$ True

Check $x = 3 \Rightarrow \sqrt{3(3) - 5} = 3 - 1$
 $\Rightarrow 2 = 2$ True

(v) $\sqrt{2x + 5} = x + 1$
 $\Rightarrow (\sqrt{2x + 5})^2 = (x + 1)^2$
 $\Rightarrow 2x + 5 = x^2 + 2x + 1$
 $\Rightarrow x^2 = 4$
 $\Rightarrow x = -2, x = 2$

Check $x = -2 \Rightarrow \sqrt{2(-2) + 5} = -2 + 1$
 $\Rightarrow 1 = -1$ False

Check $x = 2 \Rightarrow \sqrt{2(2) + 5} = 2 + 1$
 $\Rightarrow 3 = 3$ True

(vi) $\sqrt{2x^2 - 7} = x + 3$
 $\Rightarrow (\sqrt{2x^2 - 7})^2 = (x + 3)^2$
 $\Rightarrow 2x^2 - 7 = x^2 + 6x + 9$
 $\Rightarrow x^2 - 6x - 16 = 0$
 $\Rightarrow (x + 2)(x - 8) = 0$
 $\Rightarrow x = -2, x = 8$

Check $x = -2 \Rightarrow \sqrt{2(-2)^2 - 7} = -2 + 3$
 $\Rightarrow 1 = 1$ True

Check $x = 8 \Rightarrow \sqrt{2(8)^2 - 7} = 8 + 3$
 $\Rightarrow 11 = 11$ True

6. (i) $\sqrt{x + 5} = 5 - \sqrt{x}$
 $\Rightarrow (\sqrt{x + 5})^2 = (5 - \sqrt{x})^2$
 $\Rightarrow x + 5 = 25 - 10\sqrt{x} + x$
 $\Rightarrow 10\sqrt{x} = 20$
 $\Rightarrow \sqrt{x} = 2$
 $\Rightarrow x = 4$

Check $x = 4 \Rightarrow \sqrt{4 + 5} = 5 - \sqrt{4}$
 $\Rightarrow 3 = 3$ True

$$(ii) \sqrt{5x+6} = \sqrt{2x} + 2$$

$$\Rightarrow (\sqrt{5x+6})^2 = (\sqrt{2x} + 2)^2$$

$$\Rightarrow 5x + 6 = 2x + 4\sqrt{2x} + 4$$

$$\Rightarrow 3x + 2 = 4\sqrt{2x}$$

$$\Rightarrow (3x + 2)^2 = (4\sqrt{2x})^2$$

$$\Rightarrow 9x^2 + 12x + 4 = 16(2x)$$

$$\Rightarrow 9x^2 - 20x + 4 = 0$$

$$\Rightarrow (9x - 2)(x - 2) = 0$$

$$\Rightarrow x = \frac{2}{9}, x = 2$$

$$(iii) \sqrt{x+7} = 7 - \sqrt{x}$$

$$\Rightarrow (\sqrt{x+7})^2 = (7 - \sqrt{x})^2$$

$$\Rightarrow x + 7 = 49 - 14\sqrt{x} + x$$

$$\Rightarrow 14\sqrt{x} = 42$$

$$\Rightarrow \sqrt{x} = 3$$

$$\Rightarrow x = 9$$

$$(iv) \sqrt{3x-2} = \sqrt{x-2} + 2$$

$$\Rightarrow (\sqrt{3x-2})^2 = (\sqrt{x-2} + 2)^2$$

$$\Rightarrow 3x - 2 = x - 2 + 4\sqrt{x-2} + 4$$

$$\Rightarrow 2x - 4 = 4\sqrt{x-2}$$

$$\Rightarrow x - 2 = 2\sqrt{x-2}$$

$$\Rightarrow (x - 2)^2 = (2\sqrt{x-2})^2$$

$$\Rightarrow x^2 - 4x + 4 = 4(x - 2) = 4x - 8$$

$$\Rightarrow x^2 - 8x + 12 = 0$$

$$\Rightarrow (x - 2)(x - 6) = 0$$

$$\Rightarrow x = 2, x = 6$$

$$\begin{aligned} 7. \frac{1}{\sqrt{x+2}} - \frac{1}{\sqrt{4x+8}} &= \frac{1}{\sqrt{x+2}} - \frac{1}{\sqrt{4(x+2)}} \\ &= \frac{1}{\sqrt{x+2}} - \frac{1}{2\sqrt{x+2}} \\ &= \frac{2-1}{2\sqrt{x+2}} \\ &= \frac{1}{2\sqrt{x+2}} \end{aligned}$$

Thus

$$\frac{1}{\sqrt{x+2}} - \frac{1}{\sqrt{4x+8}} = 2$$

$$\frac{1}{2\sqrt{x+2}} = 2$$

$$1 = 4\sqrt{x+2}$$

$$1 = 16(x+2)$$

$$1 = 16x + 32$$

$$-31 = 16x$$

$$x = \frac{-31}{16}$$

$$\text{Check } x = \frac{2}{9} \Rightarrow \sqrt{5\left(\frac{2}{9}\right) + 6} = \sqrt{\frac{4}{9}} + 2$$

$$\Rightarrow \sqrt{\frac{64}{9}} = \frac{2}{3} + 2$$

$$\Rightarrow \frac{8}{3} = 2\frac{2}{3} \quad \text{True}$$

$$\text{Check } x = 2 \Rightarrow \sqrt{5(2) + 6} = \sqrt{2(2)} + 2$$

$$\Rightarrow \sqrt{16} = 2 + 2$$

$$\Rightarrow 4 = 4 \quad \text{True}$$

$$\text{Check } x = 9 \Rightarrow \sqrt{9+7} + \sqrt{9} = 7$$

$$\Rightarrow 4 + 3 = 7 \quad \text{True}$$

$$\text{Check } x = 2 \Rightarrow \sqrt{3(2) - 2} = \sqrt{2 - 2} + 2$$

$$\Rightarrow \sqrt{4} = 0 + 2$$

$$\Rightarrow 2 = 2 \quad \text{True}$$

$$\text{Check } x = 6 \Rightarrow \sqrt{3(6) - 2} = \sqrt{6 - 2} + 2$$

$$\Rightarrow \sqrt{16} = \sqrt{4} + 2$$

$$\Rightarrow 4 = 2 + 2 \quad \text{True}$$

8. $x = \sqrt{4x + 5}$

$$x^2 = 4x + 5$$

$$x^2 - 4x - 5 = 0$$

$$(x - 5)(x + 1) = 0$$

$$x = 5 \quad \text{or} \quad x = -1$$

check: $x = 5: \quad 5 = \sqrt{25}$
 $5 = 5 \quad \text{True.}$
 $x = -1: -1 = \sqrt{1}$
 $-1 = 1 \quad \text{False.}$

Thus the only solution is $x = 5$.

9. $x = \sqrt{a} + \frac{1}{\sqrt{a}} + 1$

$$\Rightarrow x - 2 = \sqrt{a} + \frac{1}{\sqrt{a}} + 1 - 2 = \sqrt{a} + \frac{1}{\sqrt{a}} - 1$$

Hence, $x^2 - 2x = x(x - 2)$

$$\begin{aligned} &= \left(\sqrt{a} + \frac{1}{\sqrt{a}} + 1 \right) \left(\sqrt{a} + \frac{1}{\sqrt{a}} - 1 \right) \\ &= \sqrt{a} \left(\sqrt{a} + \frac{1}{\sqrt{a}} - 1 \right) + \frac{1}{\sqrt{a}} \left(\sqrt{a} + \frac{1}{\sqrt{a}} - 1 \right) + 1 \left(\sqrt{a} + \frac{1}{\sqrt{a}} - 1 \right) \\ &= a + 1 - \sqrt{a} + 1 + \frac{1}{a} - \frac{1}{\sqrt{a}} + \sqrt{a} + \frac{1}{\sqrt{a}} - 1 \\ &= a + \frac{1}{a} + 1 \end{aligned}$$

10. $(a + \sqrt{3})(b - \sqrt{3}) = 7 + 3\sqrt{3}$

$$\Rightarrow ab - \sqrt{3}a + \sqrt{3}b - 3 = 7 + 3\sqrt{3}$$

$$\Rightarrow (ab - 3) + (-a + b)\sqrt{3} = 7 + 3\sqrt{3}$$

$$\Rightarrow ab - 3 = 7 \text{ and } -a + b = 3$$

$$\Rightarrow b = a + 3$$

$$\Rightarrow a(a + 3) - 3 - 7 = 0$$

$$\Rightarrow a^2 + 3a - 10 = 0$$

$$\Rightarrow (a + 5)(a - 2) = 0$$

$$\Rightarrow a = -5 \text{ (Not valid), } a = 2 \text{ (Valid)}$$

$$\Rightarrow b = 2 + 3 = 5$$

11. (i) $|IC|^2 = (x + 2)^2 + (x - 2)^2$
 $= x^2 + 4x + 4 + x^2 - 4x + 4$
 $= 2x^2 + 8$
 $\Rightarrow |IC| = \sqrt{2x^2 + 8}$

(ii) $|ID|^2 = (\sqrt{2x^2 + 8})^2 + x^2$
 $= 2x^2 + 8 + x^2$
 $= 3x^2 + 8$

$$\Rightarrow |ID| = \sqrt{3x^2 + 8}$$


Hence, $\sqrt{3x^2 + 8} = \sqrt{56}$

$$\Rightarrow 3x^2 + 8 = 56$$

$$\Rightarrow 3x^2 = 48$$

$$\Rightarrow x^2 = 16$$

$$\Rightarrow x = -4 \text{ (Not valid), } x = 4 \text{ (Valid)}$$



Exercises 2.9

1. $f(x) = x^2 - 8x + 15$ If $(x - 3)$ is a factor, then $x = 3$ is a root.

$$\begin{aligned}\Rightarrow f(3) &= (3)^2 - 8(3) + 15 \\ &= 9 - 24 + 15 \\ &= 24 - 24 = 0 \quad \Rightarrow (x - 3) \text{ is a factor}\end{aligned}$$

2. If $(x - 1)$ is a factor, then $x = 1$ is a root.

$$\begin{aligned}f(x) &= x^3 - x^2 - 9x + 9 \\ \Rightarrow f(1) &= (1)^3 - (1)^2 - 9(1) + 9 \\ &= 1 - 1 - 9 + 9 = 10 - 10 = 0 \\ \Rightarrow (x - 1) &\text{ is a factor}\end{aligned}$$

3. If $(x + 2)$ is a factor, then $x = -2$ is a root.

$$\begin{aligned}f(x) &= x^3 + 6x^2 + 11x + 6 \\ \Rightarrow f(-2) &= (-2)^3 + 6(-2)^2 + 11(-2) + 6 \\ &= -8 + 24 - 22 + 6 \\ &= 30 - 30 = 0 \\ \Rightarrow (x + 2) &\text{ is a factor}\end{aligned}$$

4. If $(x - 2)$ is a factor, then $x = 2$ is a root.

$$\begin{aligned}f(x) &= 2x^3 - 3x^2 - 12x + 20 \\ \Rightarrow f(2) &= 2(2)^3 - 3(2)^2 - 12(2) + 20 \\ &= 16 - 12 - 24 + 20 \\ &= 36 - 36 = 0 \quad \Rightarrow x - 2 \text{ is a factor}\end{aligned}$$

5. If $(x - 2)$ is a factor, then $x = 2$ is a root.

$$\begin{aligned}f(x) &= x^3 - 5x^2 + 8x - 4 \\ \Rightarrow f(2) &= (2)^3 - 5(2)^2 + 8(2) - 4 \\ &= 8 - 20 + 16 - 4 \\ &= 24 - 24 = 0 \quad \Rightarrow (x - 2) \text{ is a factor}\end{aligned}$$

6. If $(2x - 1)$ is a factor $\Rightarrow x = \frac{1}{2}$ is a root.

$$\begin{aligned}f(x) &= 2x^3 + 7x^2 + 2x - 3 \\ \Rightarrow f\left(\frac{1}{2}\right) &= 2\left(\frac{1}{2}\right)^3 + 7\left(\frac{1}{2}\right)^2 + 2\left(\frac{1}{2}\right) - 3 \\ &= \frac{1}{4} + \frac{7}{4} + 1 - 3 \\ &= 3 - 3 = 0 \quad \Rightarrow (2x - 1) \text{ is a factor}\end{aligned}$$

7. If $(2x + 1)$ is a factor $\Rightarrow x = -\frac{1}{2}$ is a root.

$$\begin{aligned}f(x) &= 2x^3 - x^2 - 5x - 2 \\ \Rightarrow f\left(-\frac{1}{2}\right) &= 2\left(-\frac{1}{2}\right)^3 - \left(-\frac{1}{2}\right)^2 - 5\left(-\frac{1}{2}\right) - 2 \\ &= -\frac{1}{4} - \frac{1}{4} + \frac{5}{2} - 2 \\ &= \frac{5}{2} - \frac{5}{2} = 0 \quad \Rightarrow (2x + 1) \text{ is a factor}\end{aligned}$$

8. If $(x - 1)$ is a factor $\Rightarrow x = 1$ is a root.

$$\begin{aligned}f(x) &= x^3 + kx^2 - x - 8 \\ \Rightarrow f(1) &= (1)^3 + k(1)^2 - (1) - 8 = 0 \\ \Rightarrow 1 + k - 1 - 8 &= 0 \\ \Rightarrow k &= 8\end{aligned}$$

9. If $(x + 2)$ is a factor $\Rightarrow x = -2$ is a root.

$$\begin{aligned} f(x) &= x^3 + 6x^2 + px + 6 \\ \Rightarrow f(-2) &= (-2)^3 + 6(-2)^2 + p(-2) + 6 = 0 \\ &\Rightarrow -8 + 24 - 2p + 6 = 0 \\ &\quad -2p = -22 \\ &\quad \Rightarrow p = 11 \end{aligned}$$

10. If $(x - 3)$ is a factor $\Rightarrow x = 3$ is a root.

$$\begin{aligned} f(x) &= x^3 - 2x^2 - 5x + 6 \\ \Rightarrow f(3) &= (3)^3 - 2(3)^2 - 5(3) + 6 \\ &= 27 - 18 - 15 + 6 = 33 - 33 = 0 \Rightarrow x = 3 \text{ is a root.} \end{aligned}$$

$$\begin{array}{r} x^2 + x - 2 \\ x - 3 \overline{) x^3 - 2x^2 - 5x + 6} \\ \underline{x^3 - 3x^2} \\ x^2 - 5x \\ \underline{x^2 - 3x} \\ -2x + 6 \\ \underline{-2x + 6} \\ 0 \end{array} \quad \begin{aligned} &\Rightarrow x^2 + x - 2 \\ &= (x + 2)(x - 1) \\ &\therefore \text{the other two factors are } (x + 2) \text{ and } (x - 1) \end{aligned}$$

11. If $(x + 3)$ is a factor $\Rightarrow x = -3$ is a root.

$$\begin{aligned} f(x) &= x^3 - 2x^2 - 9x + 18 \\ \Rightarrow f(-3) &= (-3)^3 - 2(-3)^2 - 9(-3) + 18 \\ &= -27 - 18 + 27 + 18 = 45 - 45 = 0 \Rightarrow x = -3 \text{ is a root.} \end{aligned}$$

$$\begin{array}{r} x^2 - 5x + 6 \\ x + 3 \overline{) x^3 - 2x^2 - 9x + 18} \\ \underline{x^3 + 3x^2} \\ -5x^2 - 9x \\ \underline{-5x^2 - 15x} \\ 6x + 18 \\ \underline{6x + 18} \\ 0 \end{array} \quad \begin{aligned} &\Rightarrow x^2 - 5x + 6 \\ &= (x - 2)(x - 3) \\ &\therefore \text{the other two factors are } (x - 2) \text{ and } (x - 3) \end{aligned}$$

12. (i) $f(x) = x^3 - 4x^2 - x + 4$
 $\Rightarrow f(1) = (1)^3 - 4(1)^2 - (1) + 4$
 $= 1 - 4 - 1 + 4 = 5 - 5 = 0$

$$\begin{aligned} &\Rightarrow x = 1 \text{ is a root} \\ &\Rightarrow (x - 1) \text{ is a factor.} \end{aligned}$$

$$\begin{array}{r} x^2 - 3x - 4 \\ x - 1 \overline{) x^3 - 4x^2 - x + 4} \\ \underline{x^3 - x^2} \\ -3x^2 - x \\ \underline{-3x^2 + 3x} \\ -4x + 4 \\ \underline{-4x + 4} \\ 0 \end{array} \quad \Rightarrow x^2 - 3x - 4 = (x + 1)(x - 4)$$

$$(ii) f(x) = x^3 - 8x^2 + 19x - 12$$

$$\Rightarrow f(1) = (1)^3 - 8(1)^2 + 19(1) - 12 \\ = 1 - 8 + 19 - 12 = 20 - 20 = 0$$

$$\Rightarrow x = 1 \text{ is a root}$$

$$\Rightarrow (x - 1) \text{ is a factor.}$$

$$\begin{array}{r} x^2 - 7x + 12 \\ x - 1 \overline{) x^3 - 8x^2 + 19x - 12} \\ \underline{x^3 - x^2} \\ -7x^2 + 19x \\ \underline{-7x^2 + 7x} \\ 12x - 12 \\ \underline{12x - 12} \\ 0 \end{array}$$

$$\Rightarrow x^2 - 7x + 12$$

$$= (x - 3)(x - 4)$$

$$(iii) f(x) = x^3 + 6x^2 - x - 30$$

$$\Rightarrow f(2) = (2)^3 + 6(2)^2 - (2) - 30 \\ = 8 + 24 - 2 - 32 = 32 - 32 = 0$$

$$\Rightarrow x = 2 \text{ is a root}$$

$$\Rightarrow (x - 2) \text{ is a factor.}$$

$$\begin{array}{r} x^2 + 8x + 15 \\ x - 2 \overline{) x^3 + 6x^2 - x - 30} \\ \underline{x^3 - 2x^2} \\ 8x^2 - x \\ \underline{8x^2 - 16x} \\ 15x - 30 \\ \underline{15x - 30} \\ 0 \end{array}$$

$$\Rightarrow x^2 + 8x + 15$$

$$= (x + 3)(x + 5)$$

$$(iv) f(x) = 3x^3 - 4x^2 - 3x + 4$$

$$\Rightarrow f(1) = 3(1)^3 - 4(1)^2 - 3(1) + 4 \\ = 3 - 4 - 3 + 4 = 7 - 7 = 0$$

$$\Rightarrow x = 1 \text{ is a root}$$

$$\Rightarrow (x - 1) \text{ is a factor.}$$

$$\begin{array}{r} 3x^2 - x - 4 \\ x - 1 \overline{) 3x^3 - 4x^2 - 3x + 4} \\ \underline{3x^3 - 3x^2} \\ -x^2 - 3x \\ \underline{-x^2 + x} \\ -4x + 4 \\ \underline{-4x + 4} \\ 0 \end{array}$$

$$\Rightarrow 3x^2 - x - 4$$

$$= (3x - 4)(x + 1)$$

$$(v) f(x) = 2x^3 - 3x^2 - 8x - 3$$

$$f(-1) = 2(-1)^3 - 3(-1)^2 - 8(-1) - 3 \\ = -2 - 3 + 8 - 3 = 8 - 8 = 0$$

$$\Rightarrow x = -1 \text{ is a root}$$

$$\Rightarrow (x + 1) \text{ is a factor.}$$

$$\begin{array}{r} 2x^2 - 5x - 3 \\ x + 1 \overline{) 2x^3 - 3x^2 - 8x - 3} \\ \underline{2x^3 + 2x^2} \\ -5x^2 - 8x \\ \underline{-5x^2 - 5x} \\ -3x - 3 \\ \underline{-3x - 3} \\ 0 \end{array}$$

$$\Rightarrow 2x^2 - 5x - 3$$

$$= (2x + 1)(x - 3)$$

$$(vi) f(x) = 2x^3 - 3x^2 - 12x + 20$$

$$f(2) = 2(2)^3 - 3(2)^2 - 12(2) + 20 \\ = 16 - 12 - 24 + 20 = 36 - 36 = 0$$

$\Rightarrow x = 2$ is a root
 $\Rightarrow (x - 2)$ is a factor.

$$\begin{array}{r} 2x^2 + x - 10 \\ x - 2 \overline{) 2x^3 - 3x^2 - 12x + 20} \\ \underline{2x^3 - 4x^2} \\ x^2 - 12x \\ \underline{x^2 - 2x} \\ -10x + 20 \\ \underline{-10x + 20} \\ 0 \end{array}$$

$\Rightarrow 2x^2 + x - 10$
 $\Rightarrow (2x + 5)(x - 2)$

$$13. f(x) = 2x^3 + 13x^2 + 13x - 10$$

$$f(-2) = 2(-2)^3 + 13(-2)^2 + 13(-2) - 10 \\ = -16 + 52 - 26 - 10 = 52 - 52 = 0$$

$\Rightarrow x = -2$ is a root
 $\Rightarrow (x + 2)$ is a factor.

$$\begin{array}{r} 2x^2 + 9x - 5 \\ x + 2 \overline{) 2x^3 + 13x^2 + 13x - 10} \\ \underline{2x^3 + 4x^2} \\ 9x^2 + 13x \\ \underline{9x^2 + 18x} \\ -5x - 10 \\ \underline{-5x - 10} \\ 0 \end{array}$$

$\Rightarrow 2x^2 + 9x - 5$
 $\Rightarrow (2x - 1)(x + 5)$

$$14. (x + 2) \text{ is a factor } \Rightarrow x = -2 \text{ is a root}$$

$$f(x) = x^3 + ax^2 - x - 2 \\ \Rightarrow f(-2) = (-2)^3 + a(-2)^2 - (-2) - 2 = 0 \\ \Rightarrow -8 + 4a + 2 - 2 = 0 \\ \Rightarrow 4a = 8 \\ \Rightarrow a = 2$$

$$\begin{array}{r} x^2 - 1 \\ x + 2 \overline{) x^3 + 2x^2 - x - 2} \\ \underline{x^3 + 2x^2} \\ 0 - x - 2 \\ \underline{-x - 2} \\ 0 \end{array}$$

$$\Rightarrow x^2 - 1 = (x + 1)(x - 1)$$

$$15. f(x) = x^3 - x^2 - 14x + 24$$

$$f(2) = (2)^3 - (2)^2 - 14(2) + 24 \\ = 8 - 4 - 28 + 24 = 32 - 32 = 0$$

$\Rightarrow x = 2$ is a root
 $\Rightarrow (x - 2)$ is a factor.

$$\begin{array}{r} x^2 + x - 12 \\ x - 2 \overline{) x^3 - x^2 - 14x + 24} \\ \underline{x^3 - 2x^2} \\ x^2 - 14x \\ \underline{x^2 - 2x} \\ -12x + 24 \\ \underline{-12x + 24} \\ 0 \end{array}$$

$$\Rightarrow x^2 + x - 12 \\ = (x - 3)(x + 4)$$

Hence, roots $x = 2, 3, -4$

16. $f(x) = x^3 + 5x^2 + 2x - 8$

$$\Rightarrow f(1) = (1)^3 + 5(1)^2 + 2(1) - 8 \\ = 1 + 5 + 2 - 8 = 8 - 8 = 0$$

$$\begin{array}{r} x^2 + 6x + 8 \\ x - 1 \overline{) x^3 + 5x^2 + 2x - 8} \\ \underline{x^3 - x^2} \\ 6x^2 + 2x \\ \underline{6x^2 - 6x} \\ 8x - 8 \\ \underline{8x - 8} \\ 0 \end{array}$$

$$\Rightarrow x = 1 \text{ is a root} \\ \Rightarrow (x - 1) \text{ is a factor.}$$

$$\Rightarrow x^2 + 6x + 8 = 0 \\ \Rightarrow (x + 2)(x + 4) = 0 \\ \Rightarrow \text{roots } x = -2, -4$$

17. (i) $f(x) = x^3 - 4x^2 - x + 4$

$$f(1) = (1)^3 - 4(1)^2 - (1) + 4 \\ = 1 - 4 - 1 + 4 = 5 - 5 = 0$$

$$\begin{array}{r} x^2 - 3x - 4 \\ x - 1 \overline{) x^3 - 4x^2 - x + 4} \\ \underline{x^3 - x^2} \\ -3x^2 - x \\ \underline{-3x^2 + 3x} \\ -4x + 4 \\ \underline{-4x + 4} \\ 0 \end{array}$$

$$\Rightarrow x = 1 \text{ is a root} \\ \Rightarrow (x - 1) \text{ is a factor.}$$

$$\Rightarrow x^2 - 3x - 4 = 0 \\ \Rightarrow (x + 1)(x - 4) = 0 \\ \Rightarrow \text{roots } x = -1, 4$$

(ii) $f(x) = x^3 + 2x^2 - 11x - 12$

$$f(-1) = (-1)^3 + 2(-1)^2 - 11(-1) - 12 \\ = -1 + 2 + 11 - 12 = 13 - 13 = 0$$

$$\begin{array}{r} x^2 + x - 12 \\ x + 1 \overline{) x^3 + 2x^2 - 11x - 12} \\ \underline{x^3 + x^2} \\ x^2 - 11x \\ \underline{x^2 + x} \\ -12x - 12 \\ \underline{-12x - 12} \\ 0 \end{array}$$

$$\Rightarrow x = -1 \text{ is a root} \\ \Rightarrow (x + 1) \text{ is a factor.}$$

$$\Rightarrow x^2 + x - 12 = 0 \\ \Rightarrow (x + 4)(x - 3) = 0 \\ \Rightarrow \text{roots } x = -4, 3$$

(iii) $f(x) = 3x^3 - 4x^2 - 3x + 4$

$$f(1) = 3(1)^3 - 4(1)^2 - 3(1) + 4 \\ = 3 - 4 - 3 + 4 = 7 - 7 = 0$$

$$\begin{array}{r} 3x^2 - x - 4 \\ x - 1 \overline{) 3x^3 - 4x^2 - 3x + 4} \\ \underline{3x^3 - 3x^2} \\ -x^2 - 3x \\ \underline{-x^2 + x} \\ -4x + 4 \\ \underline{-4x + 4} \\ 0 \end{array}$$

$$\Rightarrow x = 1 \text{ is a root} \\ \Rightarrow (x - 1) \text{ is a factor.}$$

$$\Rightarrow 3x^2 - x - 4 = 0 \\ \Rightarrow (3x - 4)(x + 1) = 0 \\ 3x = 4 \text{ or } x = -1 \\ \Rightarrow \text{roots } x = \frac{4}{3}, -1$$

$$(iv) f(x) = x^3 - 7x - 6$$

$$\begin{aligned} f(-1) &= (-1)^3 - 7(-1) - 6 \\ &= -1 + 7 - 6 = 7 - 7 = 0 \end{aligned}$$

$$\begin{aligned} \Rightarrow x = -1 \text{ is a root} \\ \Rightarrow (x + 1) \text{ is a factor.} \end{aligned}$$

$$\begin{array}{r} x^2 - x - 6 \\ x + 1 \overline{) x^3 - 7x - 6} \\ \underline{x^3 + x^2} \\ -x^2 - 7x \\ \underline{-x^2 - x} \\ -6x - 6 \\ \underline{-6x - 6} \\ 0 \end{array}$$

$$\begin{aligned} \Rightarrow x^2 - x - 6 &= 0 \\ \Rightarrow (x + 2)(x - 3) &= 0 \\ \Rightarrow \text{roots } x &= -2, 3 \end{aligned}$$

18. If $(x + 1)$ is a factor $\Rightarrow x = -1$ is a root

$$f(x) = 2x^3 + ax^2 + bx - 3$$

$$\begin{aligned} f(-1) &= 2(-1)^3 + a(-1)^2 + b(-1) - 3 = 0 \\ \Rightarrow -2 + a - b - 3 &= 0 \Rightarrow a - b = 5 \end{aligned}$$

If $(x + 3)$ is a factor $\Rightarrow x = -3$ is a root

$$\begin{aligned} \Rightarrow f(-3) &= 2(-3)^3 + a(-3)^2 + b(-3) - 3 = 0 \\ \Rightarrow -54 + 9a - 3b - 3 &= 0 \\ \Rightarrow 9a - 3b &= 57 \\ \Rightarrow 3a - b &= 19 \end{aligned}$$

$$\begin{array}{r} \text{and} \quad a - b = 5 \\ \Rightarrow 2a = 14 \\ \Rightarrow a = 7 \\ \Rightarrow 7 - b = 5 \\ \Rightarrow b = 2 \end{array}$$

$$\text{Hence, } (x + 1)(x + 3) = x^2 + 4x + 3$$

$$\begin{array}{r} 2x - 1 \\ \Rightarrow x^2 + 4x + 3 \overline{) 2x^3 + 7x^2 + 2x - 3} \\ \underline{2x^3 + 8x^2 + 6x} \\ -x^2 - 4x - 3 \\ \underline{-x^2 - 4x - 3} \\ 0 \end{array} \quad \begin{aligned} 2x - 1 &= 0 \\ \Rightarrow 3^{\text{rd}} \text{ root } x &= \frac{1}{2} \end{aligned}$$

19. If $(x + 1)$ is a factor $\Rightarrow x = -1$ is a root

$$f(x) = x^3 + 5x^2 + kx - 12$$

$$\begin{aligned} f(-1) &= (-1)^3 + 5(-1)^2 + k(-1) - 12 = 0 \\ \Rightarrow -1 + 5 - k - 12 &= 0 \\ -k - 8 &= 0 \\ \Rightarrow k &= -8 \end{aligned}$$

$$\begin{array}{r} x^2 + 4x - 12 \\ \Rightarrow x + 1 \overline{) x^3 + 5x^2 - 8x - 12} \\ \underline{x^3 + x^2} \\ 4x^2 - 8x \\ \underline{4x^2 + 4x} \\ -12x - 12 \\ \underline{-12x - 12} \\ 0 \end{array}$$

$$\begin{aligned} \Rightarrow x^2 + 4x - 12 \\ \Rightarrow (x + 6)(x - 2) \text{ other 2 factors} \end{aligned}$$

- 20.** If $(x + 2)$ is a factor $\Rightarrow x = -2$ is a root

$$f(x) = 2x^3 + ax^2 - 17x + b$$

$$f(-2) = 2(-2)^3 + a(-2)^2 - 17(-2) + b = 0$$

$$\Rightarrow -16 + 4a + 34 + b = 0 \Rightarrow 4a + b = -18$$

If $(x - 3)$ is a factor $\Rightarrow x = 3$ is a root

$$\Rightarrow f(3) = 2(3)^3 + a(3)^2 - 17(3) + b = 0$$

$$\Rightarrow 54 + 9a - 51 + b = 0$$

$$\Rightarrow 9a + b = -3$$

$$\text{and } 4a + b = -18$$

$$\Rightarrow \quad \quad 5a = 15$$

$$\Rightarrow \quad \quad a = 3$$

$$\Rightarrow 4(3) + b = -18$$

$$b = -30$$

Hence, $(x + 2)(x - 3) = x^2 - x - 6$

$$\begin{array}{r} 2x + 5 \\ \Rightarrow x^2 - x - 6 \overline{) 2x^3 + 3x^2 - 17x - 30} \\ \underline{2x^3 - 2x^2 - 12x} \\ 5x^2 - 5x - 30 \\ \underline{5x^2 - 5x - 30} \\ 0 \end{array}$$

$\Rightarrow 3^{\text{rd}}$ factor $(2x + 5)$

- 21.** $x^2 - 2x - 3 = (x + 1)(x - 3) = 0$

= roots $x = -1, x = 3$

$$f(x) = ax^3 + 8x^2 + bx + 6$$

$$\Rightarrow f(-1) = a(-1)^3 + 8(-1)^2 + b(-1) + 6 = 0$$

$$\Rightarrow -a + 8 - b + 6 = 0$$

$$\Rightarrow a + b = 14$$

$$f(3) = a(3)^3 + 8(3)^2 + b(3) + 6 = 0$$

$$\Rightarrow 27a + 72 + 3b + 6 = 0$$

$$\Rightarrow \quad \quad 27a + 3b = -78$$

$$\Rightarrow \quad \quad 9a + b = -26$$

$$\text{and} \quad \quad \quad \underline{a + b = 14}$$

$$8a = -40$$

$$\Rightarrow \quad \quad a = -5$$

$$\Rightarrow -5 + b = 14$$

$$\Rightarrow \quad \quad b = 19$$

$$\begin{array}{r} -5x - 2 \\ \text{Hence, } x^2 - 2x - 3 \overline{) -5x^3 + 8x^2 + 19x + 6} \\ \underline{-5x^3 + 10x^2 + 15x} \\ -2x^2 + 4x + 6 \\ \underline{-2x^2 + 4x + 6} \\ 0 \end{array}$$

$\Rightarrow 3^{\text{rd}}$ factor $= -5x - 2$

$\Rightarrow 3^{\text{rd}}$ root: $-5x = 2$

$$\Rightarrow x = -\frac{2}{5}$$

- 22.** (i) $ax^3 - b = c$

$$\Rightarrow ax^3 = b + c$$

$$\Rightarrow x^3 = \frac{b + c}{a}$$

$$\Rightarrow x = \sqrt[3]{\frac{b + c}{a}}$$

- (ii) $a(x + b)^3 = c$

$$\Rightarrow (x + b)^3 = \frac{c}{a}$$

$$\Rightarrow x + b = \left(\frac{c}{a}\right)^{\frac{1}{3}}$$

$$\Rightarrow x = \left(\frac{c}{a}\right)^{\frac{1}{3}} - b$$

Exercise 2.10

1. (i) Graph crosses the x-axis at $x = -1, 1, 3$
 $\Rightarrow f(x) = a(x+1)(x-1)(x-3)$
 Point $(0, 3) \Rightarrow f(0) = a(0+1)(0-1)(0-3) = 3$
 $\Rightarrow 3a = 3$
 $\Rightarrow a = 1$
 $\therefore f(x) = 1(x+1)(x-1)(x-3) = x^3 - 3x^2 - x + 3$
- (ii) Graph crosses the x-axis at $x = -4, 1, 2$
 $\Rightarrow f(x) = a(x+4)(x-1)(x-2)$
 Point $(0, 8) \Rightarrow f(0) = a(0+4)(0-1)(0-2) = 8$
 $\Rightarrow 8a = 8$
 $\Rightarrow a = 1$
 $\therefore f(x) = 1(x+4)(x-1)(x-2) = x^3 + x^2 - 10x + 8$
2. (i) Graph crosses the x-axis at $x = -3, 0, 2$
 $\Rightarrow f(x) = a(x+3)(x)(x-2)$
 Point $(1, -4) \Rightarrow f(1) = a(1+3)(1)(1-2) = -4$
 $\Rightarrow -4a = -4$
 $\Rightarrow a = 1$
 $\therefore f(x) = 1(x+3)(x)(x-2) = x^3 + x^2 - 6x$ (Green Graph)
 Point $(1, -12) \Rightarrow f(1) = a(1+3)(1)(1-2) = -12$
 $\Rightarrow -4a = -12$
 $\Rightarrow a = 3$
 $\therefore f(x) = 3(x+3)(x)(x-2) = 3x^3 + 3x^2 - 18x$ (Blue Graph)
- (ii) Graph crosses the x-axis at $x = 1, 2, 3$
 $\Rightarrow f(x) = a(x-1)(x-2)(x-3)$
 Point $(0, 6) \Rightarrow f(0) = a(0-1)(0-2)(0-3) = 6$
 $\Rightarrow -6a = 6 \Rightarrow a = -1$
 $\therefore f(x) = -1(x-1)(x-2)(x-3) = -x^3 + 6x^2 - 11x + 6$ (Red Graph)
 Point $(0, 12) \Rightarrow f(0) = a(0-1)(0-2)(0-3) = 12$
 $\Rightarrow -6a = 12$
 $\Rightarrow a = -2$
 $\therefore f(x) = -2(x-1)(x-2)(x-3) = -2x^3 + 12x^2 - 22x + 12$ (Blue Graph)
3. Graph crosses the x-axis at $x = 1, -2, \frac{1}{2}$
 $\Rightarrow f(x) = a(x-1)(x+2)(2x-1)$
 Point $(0, 6) \Rightarrow f(0) = a(0-1)(0+2)(0-1) = 6$
 $\Rightarrow 2a = 6 \Rightarrow a = 3$
 $\therefore f(x) = 3(x-1)(x+2)(2x-1)$
 $= 6x^3 + 3x^2 - 15x + 6$
4. $f(x) = (x-3)(x+1)(x+2)$
 $= (x-3)(x^2 + 3x + 2)$
 $= x^3 - 7x - 6 \Rightarrow a = 0, b = -7, c = -6$
5. (i) $f(x) = x^3 + 2$ Point $(0, 2) \Rightarrow f(0) = 0^3 + 2 = 2$ True
 (ii) $g(x) = x^3$ Point $(1, 1) \Rightarrow f(1) = (1)^3 = 1$ True
 (iii) $h(x) = 2x^3$ Point $(1, 2) \Rightarrow f(1) = 2(1)^3 = 2$ True
 $A(2^{\frac{1}{3}}, 4)$

6. $f(x) = x(x - 4)(x - 6)$


$$f(2) = 2(2 - 4)(2 - 6)$$

$$= 2(-2)(-4) = 16$$

$$f(5) = 5(5 - 4)(5 - 6)$$

$$= 5(1)(-1) = -5$$

Points: (0, 0) (4, 0) (6, 0) (2, 16) (5, -5)




7. $f(x) = (x + 2)(x - 1)(x - 3)$

$$f(0) = (0 + 2)(0 - 1)(0 - 3) = 6$$

$$f\left(\frac{1}{2}\right) = \left(\frac{1}{2} + 2\right)\left(\frac{1}{2} - 1\right)\left(\frac{1}{2} - 3\right) = 3\frac{1}{8}$$

$$f(2) = (2 + 2)(2 - 1)(2 - 3) = -4$$

Points: (-2, 0) (1, 0) (3, 0) (0, 6) $\left(\frac{1}{2}, 3\frac{1}{8}\right)$ (2, -4)



8. (i) Roots are $x = -1, 1, 2$

$$\Rightarrow f(x) = a(x + 1)(x - 1)(x - 2)^2$$

(ii) Point (0, 4) $\Rightarrow f(0) = a(0 + 1)(0 - 1)(0 - 2)^2 = 4$

$$\Rightarrow -4a = 4$$

$$a = -1$$

$$\text{Hence, } f(x) = -1(x^2 - 1)(x^2 - 4x + 4)$$

$$= -1x^4 + 4x^3 - 3x^2 - 4x + 4 = ax^4 + bx^3 + cx^2 + dx + e$$

$$\Rightarrow a = -1, b = 4, c = -3, d = -4, e = 4$$

9. (i) $(0, 4) \in f(x) \Rightarrow f(0) = 4$

$$(0, -2) \in g(x) \Rightarrow g(0) = -2$$

$$\text{Hence, } 4 = a(-2)$$

$$\Rightarrow a = -2$$

(ii) $f(x)$ has roots $x = -2, 1$ (double roots)

$$\Rightarrow f(x) = a(x + 2)^2(x - 1)^2$$

$$\text{Point } (0, 4) \Rightarrow f(0) = a(0 + 2)^2(0 - 1)^2 = 4$$

$$\Rightarrow 4a = 4$$

$$\Rightarrow a = 1$$

$$\text{Hence, } f(x) = (x + 2)^2(x - 1)^2$$

$$g(x) \text{ has roots } x = -2, 1 \text{ (double roots)}$$

$$\Rightarrow g(x) = a(x + 2)^2(x - 1)^2$$

$$\text{Point } (0, -2) \Rightarrow g(0) = a(0 + 2)^2(0 - 1)^2 = -2$$

$$\Rightarrow 4a = -2$$

$$\Rightarrow a = -\frac{1}{2}$$

$$\text{Hence, } g(x) = -\frac{1}{2}(x + 2)^2(x - 1)^2$$

13. (i) Graph $f(x) = 0$ meets x -axis at $x = -1, 0, 2$
 (ii) $f(x) = g(x) \Rightarrow$ Curve meets line at $x = -1.3, 0, 2.3$
 (iii) $f(x) = g(x) \Rightarrow x^3 - x^2 - 2x = x$
 $\Rightarrow x^3 - x^2 - 3x = 0$
 $\Rightarrow x(x^2 - x - 3) = 0$

$$\begin{aligned}\Rightarrow x = 0 \quad \text{OR} \quad x &= \frac{1 \pm \sqrt{(-1)^2 - 4(1)(-3)}}{2(1)} \\ x &= \frac{1 \pm \sqrt{13}}{2} \\ x &= \frac{1 + 3.60555}{2}, \frac{1 - 3.60555}{2} \\ &= 2.303, -1.303 \\ &= 2.3, -1.3 \\ \Rightarrow x &= 0, 2.3 \text{ or } -1.3\end{aligned}$$

14. (i) Volume $= x(x-1)(x+1) = (x^3 - x) \text{ cm}^3$
 (ii) Volume $= 24 = x^3 - x$
 $\Rightarrow f(x) = x^3 - x - 24$
 $f(3) = (3)^3 - (3) - 24 = 27 - 27 = 0$
 $\Rightarrow x = 3$ is a root

15. $h = 2r \Rightarrow r = \frac{h}{2}$

$$V = \pi \left(\frac{h}{2}\right)^2 \cdot h = \frac{\pi}{4} h^3$$

- (i) Hence, $ah^3 = \frac{\pi}{4} h^3 \Rightarrow a = \frac{3.14}{4} = 0.785 = 0.79$
 (ii) Volume cylinder $= ah^3 = (0.79)(11)^3 = 1051.49 \text{ cm}^3$
 (iii) Volume $= 215.58 = ah^3$
 $\Rightarrow (0.79)h^3 = 215.58$
 $\Rightarrow h^3 = \frac{215.58}{0.79} = 272.886$
 $\Rightarrow h = \sqrt[3]{272.886} = 6.486 = 6.5 \text{ cm}$

16. Volume of cube $= (3)^3 = 27$
 Volume of sphere $= (4.19)(27) = 113.13$
 Volume of cube $= x^3 = 113.13$
 $\Rightarrow x = \sqrt[3]{113.13} = 4.836 = 4.8 \text{ cm}$
 Hence, $4.19x^3 - 150 = x^3$
 $\Rightarrow 3.19x^3 = 150$
 $\Rightarrow x^3 = \frac{150}{3.19} = 47.0219$
 $\Rightarrow x = \sqrt[3]{47.0219} = 3.609 = 3.6 \text{ cm}$

Revision Exercise 2 (Core)

1. $x^2 - 6x + 5 = 0$
 $\Rightarrow (x-1)(x-5) = 0$
 $\Rightarrow x = 1, x = 5$

$$\text{Hence, } t - \frac{6}{t} = 1 \quad \text{OR} \quad t - \frac{6}{t} = 5$$

$$\Rightarrow t^2 - 6 = t \quad \text{OR} \quad t^2 - 6 = 5t$$

$$\Rightarrow t^2 - t - 6 = 0 \quad \text{OR} \quad t^2 - 5t - 6 = 0$$

$$\Rightarrow (t+2)(t-3) = 0 \quad \text{OR} \quad (t+1)(t-6) = 0$$

$$\Rightarrow t = -2, 3, -1, 6$$

$$2. \quad 2(x+1)(x-4) - (x-2)^2 = 0$$

$$\Rightarrow 2x^2 - 6x - 8 - (x^2 - 4x + 4) = 0$$

$$\Rightarrow x^2 - 2x - 12 = 0$$

$$\Rightarrow x = \frac{2 \pm \sqrt{(2)^2 - 4(1)(-12)}}{2(1)}$$

$$= \frac{2 \pm \sqrt{52}}{2}$$

$$= \frac{2 \pm 2\sqrt{13}}{2} = 1 \pm \sqrt{13}$$

$$3. \quad px^2 + 2x + 1 = 0$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{(2)^2 - 4(p)(1)}}{2(p)}$$

$$= \frac{-2 \pm \sqrt{4 - 4p}}{2p}$$

$$\text{No solution} \Rightarrow 4 - 4p < 0$$

$$\Rightarrow -4p < -4$$

$$\Rightarrow 4p > 4$$

$$\Rightarrow p > 1$$

$$4. \quad \text{Real roots} \Rightarrow b^2 - 4ac \geq 0$$

$$\Rightarrow (-a - d)^2 - 4(1)(ad - b^2) \geq 0$$

$$\Rightarrow a^2 + 2ad + d^2 - 4ad + 4b^2 \geq 0$$

$$\Rightarrow a^2 - 2ad + d^2 + 4b^2 \geq 0$$

$$\Rightarrow (a - d)^2 + (2b)^2 \geq 0 \quad \text{True}$$

$$5. \quad (x+1) \text{ and } (x-2) \text{ are factors} \Rightarrow x = -1 \text{ and } x = 2 \text{ are roots}$$

$$f(x) = 6x^4 - x^3 + ax^2 - 6x + b$$

$$\Rightarrow f(-1) = 6(-1)^4 - (-1)^3 + a(-1)^2 - 6(-1) + b = 0$$

$$\Rightarrow 6 + 1 + a + 6 + b = 0$$

$$\Rightarrow a + b = -13$$

$$f(2) = 6(2)^4 - (2)^3 + a(2)^2 - 6(2) + b = 0$$

$$\Rightarrow 96 - 8 + 4a - 12 + b = 0$$

$$\Rightarrow 4a + b = -76$$

$$\text{and } \underline{a + b = -13}$$

$$\Rightarrow \underline{3a = -63}$$

$$\Rightarrow \underline{a = -21}$$

$$\Rightarrow -21 + b = -13 \Rightarrow b = 8$$

$$6. \quad (i) \quad f(x) = x^3 - 4x^2 - 11x + 30$$

$$f(2) = (2)^3 - 4(2)^2 - 11(2) + 30$$

$$= 8 - 16 - 22 + 30 = 38 - 38 = 0$$

$$\Rightarrow x = 2 \text{ is a root} \quad \Rightarrow (x - 2) \text{ is a factor}$$

$$\begin{array}{r}
 \text{(ii)} \quad \frac{x^2 - 2x - 15}{x - 2} \overline{) x^3 - 4x^2 - 11x + 30} \\
 \underline{x^3 - 2x^2} \\
 -2x^2 - 11x \\
 \underline{-2x^2 + 4x} \\
 -15x + 30 \\
 \underline{-15x + 30} \\
 0
 \end{array}
 \Rightarrow x^2 - 2x - 15 = (x + 3)(x - 5)$$

$$\begin{aligned}
 \text{(iii)} \quad f(x) &= (x - 2)(x + 3)(x - 5) = 0 \\
 \Rightarrow \text{Roots: } x &= 2, -3, 5
 \end{aligned}$$

7. (i) $b^2 - 4ac = (-2)^2 - 4(1)(-5) = 24 > 0 \Rightarrow 2 \text{ distinct real roots}$
- (ii) $b^2 - 4ac = (-4)^2 - 4(1)(6) = 16 - 24 = -8 < 0 \Rightarrow 2 \text{ imaginary roots}$
- (iii) $b^2 - 4ac = (4)^2 - 4(-1)(-6) = 16 - 24 = -8 < 0 \Rightarrow 2 \text{ imaginary roots}$

$$\begin{aligned}
 8. \quad 3^{2x} - 12(3^x) + 27 &= 0 \\
 \Rightarrow (3^x)^2 - 12(3^x) + 27 &= 0 \\
 \text{If } y = 3^x \Rightarrow y^2 - 12y + 27 &= 0 \\
 \Rightarrow (y - 3)(y - 9) &= 0 \\
 \Rightarrow y = 3 \quad \text{OR} \quad y = 9 \\
 \Rightarrow 3^x = 3^1 \quad \text{OR} \quad 3^x = 9 = 3^2 \\
 \Rightarrow x = 1 \quad \text{OR} \quad x = 2
 \end{aligned}$$

Revision Exercise 2 (Advanced)

1. $2x^2 - 4x - 5$
 $= 2(x^2 - 2x) - 5$
 $= 2(x^2 - 2x + 1) - 2 - 5$
 $= 2(x - 1)^2 - 7$
- (i) $2x^2 - 4x - 5 = 0$
 $\Rightarrow 2(x - 1)^2 = 7$
 $\Rightarrow (x - 1)^2 = \frac{7}{2}$
 $\Rightarrow x - 1 = \pm \sqrt{\frac{7}{2}}$
 $\Rightarrow x = 1 \pm \sqrt{\frac{7}{2}}$
- (ii) Minimum Point = (1, -7)
2. $(2\sqrt{2} - \sqrt{3})^2 = (2\sqrt{2})^2 - 2(2\sqrt{2})(\sqrt{3}) + (-\sqrt{3})^2$
 $= 8 - 4\sqrt{6} + 3$
 $= 11 - 4\sqrt{6}$
3. $\frac{\sqrt{7} + \sqrt{5}}{\sqrt{80} + \sqrt{5}} = \frac{\sqrt{7} + \sqrt{5}}{4\sqrt{5} + \sqrt{5}} = \frac{\sqrt{7} + \sqrt{5}}{5\sqrt{5}}$
 $= \frac{\sqrt{7} + \sqrt{5}}{5\sqrt{5}} \cdot \frac{\sqrt{5}}{\sqrt{5}} = \frac{\sqrt{35} + 5}{25}$
4. $\sqrt{x + 2} = x - 4$
 $\Rightarrow x + 2 = (x - 4)^2 = x^2 - 8x + 16$
 $\Rightarrow x^2 - 9x + 14 = 0$
 $\Rightarrow (x - 2)(x - 7) = 0$

$$\Rightarrow x = 2, 7$$

$$\begin{aligned} \text{Test } x = 2 &\Rightarrow \sqrt{2+2} = 2-4 \\ &\Rightarrow \sqrt{4} = -2 \quad \text{False} \end{aligned}$$

$$\begin{aligned} \text{Test } x = 7 &\Rightarrow \sqrt{7+2} = 7-4 \\ &\Rightarrow \sqrt{9} = 3 \quad \text{True} \end{aligned}$$

5. (i) Test $t = 1 \Rightarrow s = 8(1)^2 + 4(1) = 8 + 4 = 12 \text{ m}$
 $\Rightarrow t < 1 \Rightarrow \text{Test } t = 0.9 \Rightarrow s = 8(0.9)^2 + 4(0.9) = 10.08 \text{ m}$

(ii) Solve $8t^2 + 4t = 10$
 $\Rightarrow 4t^2 + 2t - 5 = 0$
 $\Rightarrow t = \frac{-2 \pm \sqrt{(2)^2 - 4(4)(-5)}}{2(4)}$
 $= \frac{-2 \pm \sqrt{84}}{8}$
 $= \frac{-2 + 9.165}{8}, \frac{-2 - 9.165}{8}$
 $= 0.8956 \text{ (Valid)}, -1.3956 \text{ (Not valid)}$
 $= 0.90$

(iii) Answer = 0.8956
 Corrected answer = 0.90 $\Rightarrow \text{error} = 0.0044$
 $\Rightarrow \text{Percentage error} = \frac{0.0044}{0.8956} \times 100$
 $= 0.491\%$
 $= 0.49\%$

6. $\sigma = \frac{\sqrt{p(1+p)}}{n}$

$$\begin{aligned} \Rightarrow \sqrt{p+p^2} &= \sigma n \\ \Rightarrow p+p^2 &= \sigma^2 n^2 \\ \Rightarrow p^2 + p - \sigma^2 n^2 &= 0 \\ \Rightarrow p &= \frac{-1 \pm \sqrt{(1)^2 - 4(1)(-\sigma^2 n^2)}}{2(1)} \\ &= \frac{-1 \pm \sqrt{(1)^2 + 4\sigma^2 n^2}}{2} \\ &= -\frac{1}{2} \pm \frac{\sqrt{1+4\sigma^2 n^2}}{2} \end{aligned}$$

7.

	$k < 0$	$0 < k < \frac{1}{4}$	$k > \frac{1}{4}$
k	Negative	Positive	Positive
$4k$	Negative	Positive	Positive
$4k - 1$	Negative	Negative	Positive
$k(4k - 1)$	Positive	Negative	Positive

Solve $x^2 + 4kx + k = 0$.

$$\Rightarrow x = \frac{-4k \pm \sqrt{(4k)^2 - 4(1)(k)}}{2(1)}$$

$$\Rightarrow x = \frac{-4k \pm \sqrt{16k^2 - 4k}}{2}$$

$$\Rightarrow x = \frac{-4k \pm \sqrt{4} \cdot \sqrt{4k^2 - k}}{2}$$

$$\Rightarrow \text{Roots: } x = -2k \pm \sqrt{k(4k-1)}$$

Hence, $x^2 + 4kx + k > 0$ for $0 < k < \frac{1}{4}$

8. $ax^2 + 2bx + c = 0$

$$\Rightarrow \text{Roots: } x = \frac{-2b \pm \sqrt{(2b)^2 - 4(a)(c)}}{2a}$$

$$\Rightarrow x = \frac{-2b \pm \sqrt{4b^2 - 4ac}}{2a}$$

$$\Rightarrow x = \frac{-2b \pm \sqrt{4} \cdot \sqrt{b^2 - ac}}{2a}$$

$$\Rightarrow x = \frac{-b \pm \sqrt{b^2 - ac}}{a} \quad (\text{Real})$$

Similarly, $bx^2 + 2cx + a = 0 \Rightarrow \text{Roots: } x = \frac{-c \pm \sqrt{c^2 - ab}}{b} \quad (\text{Real})$

True if $c > a \Rightarrow bc > a^2$

and $b > a \Rightarrow a^2 - bc < 0$

Hence, $cx^2 + 2ax + b = 0 \Rightarrow \text{Roots: } x = \frac{-a \pm \sqrt{a^2 - bc}}{c}$

Since $a^2 - bc < 0 \Rightarrow \text{Roots are not real}$

9. Points of intersection at A and B occur when $g(x) = f(x)$.

Hence, $x^2 + 5x - 1 = -x^2 + 5x + 3$

$$\Rightarrow 2x^2 - 4 = 0$$

$$\Rightarrow x^2 - 2 = 0$$

$$\Rightarrow x^2 = 2$$

$$\Rightarrow x = -\sqrt{2}, x = \sqrt{2}$$

$$\Rightarrow g(-\sqrt{2}) = (-\sqrt{2})^2 + 5(-\sqrt{2}) - 1$$

$$= 2 - 5\sqrt{2} - 1 = 1 - 5\sqrt{2}$$

and $g(\sqrt{2}) = (\sqrt{2})^2 + 5(\sqrt{2}) - 1$

$$= 2 + 5\sqrt{2} - 1 = 1 + 5\sqrt{2}$$

Hence, $A = (\sqrt{2}, 1 - 5\sqrt{2}), B = (-\sqrt{2}, 1 + 5\sqrt{2})$

10. $kx^2 - 2kx - 3k - 12 = 0$

$[a = k, b = -2k, c = -3k - 12]$

For equal roots:

$$b^2 - 4ac = 0$$

$$(-2k)^2 - 4(k)(-3k - 12) = 0$$

$$4k^2 + 12k^2 + 48k = 0$$

$$16k^2 + 48k = 0$$

$$k^2 + 3k = 0$$

$$k(k + 3) = 0$$

$$k = 0 \quad \text{or} \quad k = -3$$

When $k = 0$, the equation is untrue.

Thus the only solution is $k = -3$.

11. $x^2 - \sqrt{3}x - 6 = 0$

$$\Rightarrow x = \frac{\sqrt{3} \pm \sqrt{(\sqrt{3})^2 - 4(1)(-6)}}{2(1)}$$

$$= \frac{\sqrt{3} \pm \sqrt{27}}{2} = \frac{\sqrt{3} + 3\sqrt{3}}{2}, \frac{\sqrt{3} - 3\sqrt{3}}{2}$$

$$= 2\sqrt{3}, -\sqrt{3}$$

Hence, $r_1 r_2 = (2\sqrt{3})(-\sqrt{3}) = -6$

12. $3x + y = -1 \cap x^2 + y^2 = 53$

$$\Rightarrow y = -3x - 1 \Rightarrow x^2 + (-3x - 1)^2 = 53$$

$$\Rightarrow x^2 + 9x^2 + 6x + 1 = 53$$

$$\Rightarrow 10x^2 + 6x - 52 = 0$$

$$\Rightarrow 5x^2 + 3x - 26 = 0$$

$$\begin{aligned}
&\Rightarrow (5x + 13)(x - 2) = 0 \\
&\Rightarrow 5x = -13, x = 2 \\
&\Rightarrow x = -\frac{13}{5} \quad \Rightarrow y = -3(2) - 1 \\
&\quad \quad \quad = -7 \\
&\Rightarrow y = -3\left(-\frac{13}{5}\right) - 1 \\
&\quad \quad \quad = \frac{34}{5}
\end{aligned}$$

\therefore the solutions are: $\left(-\frac{13}{5}, \frac{34}{5}\right)$ and $(2, -7)$

13. length = x and width = y .

$$\begin{aligned}
&\Rightarrow 2x + 2y = 48 \text{ and } x = \frac{1}{2}y^2 \\
&\Rightarrow x + y = 24 \\
&\Rightarrow x = -y + 24 \quad \Rightarrow -y + 24 = \frac{1}{2}y^2 \\
&\quad \quad \quad \Rightarrow -2y + 48 = y^2 \\
&\quad \quad \quad \Rightarrow y^2 + 2y - 48 = 0 \\
&\quad \quad \quad \Rightarrow (y + 8)(y - 6) = 0 \\
&\quad \quad \quad \Rightarrow y = -8 \text{ (Not valid), } y = 6 \text{ m (Valid)} \\
&\quad \quad \quad \Rightarrow x = -6 + 24 \\
&\quad \quad \quad \quad \quad = 18 \text{ m}
\end{aligned}$$

\Rightarrow Dimensions are 18 m and 6 m


$$\begin{aligned}
\text{14. } \frac{y}{1} &= \frac{x^2}{x+1} \Rightarrow x^2 = xy + y \\
&\Rightarrow x^2 - xy - y = 0 \\
&\Rightarrow x = \frac{y \pm \sqrt{(-y)^2 - 4(1)(-y)}}{2(1)} \\
&\quad \quad \quad = \frac{y \pm \sqrt{y^2 + 4y}}{2}
\end{aligned}$$

$$\text{Real roots } \Rightarrow b^2 - 4ac \geq 0 \quad \Rightarrow y^2 + 4y \geq 0$$

Factors: $y(y + 4)$

\Rightarrow Roots: $y = 0, -4$

$$\text{For } y^2 + 4y \geq 0 \Rightarrow y \leq -4 \text{ or } y \geq 0$$



15. $y = ax^2 + bx + c$

$$\text{Point } (-2, -1) \Rightarrow -1 = a(-2)^2 + b(-2) + c$$

$$\Rightarrow -1 = 4a - 2b + c$$

$$\text{Point } (1, 2) \Rightarrow 2 = a(1)^2 + b(1) + c$$

$$\Rightarrow 2 = a + b + c$$

$$\text{Point } (3, -16) \Rightarrow -16 = a(3)^2 + b(3) + c$$

$$\Rightarrow -16 = 9a + 3b + c$$

$$\Rightarrow 4a - 2b + c = -1 \quad 9a + 3b + c = -16$$

$$\text{and } a + b + c = 2 \quad a + b + c = 2$$

$$\Rightarrow 3a - 3b = -3 \quad 8a + 2b = -18$$

$$\Rightarrow a - b = -1 \quad \Rightarrow 4a + b = -9$$

$$\text{and } a - b = -1$$

$$5a = -10$$

$$\Rightarrow a = -2$$

$$\Rightarrow -2 - b = -1 \quad \text{and } a + b + c = 2$$


$$\Rightarrow b = -1 \quad \Rightarrow -2 - 1 + c = 2$$

$$\Rightarrow c = 5$$

Hence, $y = -2x^2 - x + 5$

16. (ii) $f(x) = x^3 - 6x^2 + 11x - 6$
 $\Rightarrow f(0) = (0)^3 - 6(0)^2 + 11(0) - 6 = -6$
 $\Rightarrow f(1) = (1)^3 - 6(1)^2 + 11(1) - 6 = 12 - 12 = 0$
 $\Rightarrow f(2) = (2)^3 - 6(2)^2 + 11(2) - 6 = 30 - 30 = 0$
 $\Rightarrow f(3) = (3)^3 - 6(3)^2 + 11(3) - 6 = 60 - 60 = 0$
 $\Rightarrow f(4) = (4)^3 - 6(4)^2 + 11(4) - 6 = 108 - 102 = 6$
 \Rightarrow Roots are $x = 1, 2, 3$

- (iii) Points are $(0, -6)(1, 0)$
 $(2, 0)(3, 0)$
 $(4, 6)$



17. (i) Roots $x = 2, 5$ (double root)

(ii) $f(x) = a(x - 2)(x - 5)^2$

(iii) $f(x) = a(x^3 - 12x^2 + 45x - 50)$

Point $(3, 8) \Rightarrow f(3) = a[(3)^3 - 12(3)^2 + 45(3) - 50] = 8$

$$\Rightarrow a[27 - 108 + 135 - 50] = 8$$

$$\Rightarrow 4a = 8 \Rightarrow a = 2$$

Hence, $f(x) = 2(x^3 - 12x^2 + 45x - 50)$

$$= 2x^3 - 24x^2 + 90x - 100 \quad \Rightarrow a = 2, b = -24$$

$$c = 90, d = -100$$

(iv) Point $(3, -8) \Rightarrow f(3) = a[(3)^3 - 12(3)^2 + 45(3) - 50] = -8$

$$\Rightarrow 4a = -8 \Rightarrow a = -2$$

$$\Rightarrow f(x) = -2(x^3 - 12x^2 + 45x - 50)$$

$$= -2x^3 + 24x^2 - 90x + 100$$

(v) Reflection in y -axis $\Rightarrow x$ is replaced by $(-x)$

$$f(x) = 2x^3 - 24x^2 + 90x - 100$$

$$\Rightarrow f(-x) = 2(-x)^3 - 24(-x)^2 + 90(-x) - 100$$

$$= -2x^3 - 24x^2 - 90x - 100$$

Revision Exercise 2 (Extended-Response)

1. (i) $C(t) = 0.02t - at^3$

$$t = 5 \Rightarrow C(5) = (0.02)(5) - a(5)^3 = 0.075$$

$$\Rightarrow 0.1 - 125a = 0.075$$

$$\Rightarrow -125a = -0.025$$

$$\Rightarrow a = \frac{-0.025}{-125} = 0.0002$$

(ii) $C(t) = 0 \Rightarrow 0.02t - 0.0002t^3 = 0$

$$\Rightarrow 200t - 2t^3 = 0$$

$$\Rightarrow 100t - t^3 = 0$$

$$\Rightarrow t(100 - t^2) = 0$$

$$\Rightarrow t = 0 \text{ (not valid), } t = 10 \text{ (valid)}$$

(iii) Beyond $t = 10 \Rightarrow C(t)$ is negative

2. (i) Area $= (2x + y)(3x + 2y)$

$$= 6x^2 + 7xy + 2y^2$$

(ii) Area dividing strips $=$ Area full poster $-$ Area 6 purple squares

$$= 6x^2 + 7xy + 2y^2 - 6x^2$$

$$= 7xy + 2y^2 \Rightarrow k = 7$$

(iii) Area 6 purple squares = $6x^2 = 1.5$

$$\Rightarrow x^2 = 0.25 \Rightarrow x = 0.5 \text{ m} = \frac{1}{2} \text{ m}$$

Hence, $7xy + 4y^2 = 3.5y + 2y^2 = 1$

$$\Rightarrow 4y^2 + 7y - 2 = 0$$

$$\Rightarrow (4y - 1)(y + 2) = 0$$

$$\Rightarrow y = \frac{1}{4} \text{ (valid)}, y = -2 \text{ (Not valid)}$$

3. (a) Volume = $(96 - 4x)(48 - 2x)(x)$

$$V = 4608x - 384x^2 + 8x^3$$

(b) (i) $0 < x < 24$

(ii) At C, $x = 0 \Rightarrow \text{Volume} = 0$

At A, $x = 8 \Rightarrow \text{Maximum volume occurs}$
(= 16 384 cm³)

At B, $x = 24 \Rightarrow \text{Minimum volume occurs}$
(= 0 cm³)

(iii) Maximum volume = 16 300 cm³ (using graph)

(iv) $x = 10 \Rightarrow V = [96 - 4(10)][48 - 2(10)][10]$

$$\Rightarrow (56)(28)(10) = 15\,680 \text{ cm}^3$$

(v) $x = 5 \Rightarrow \text{Volume} = 14\,440 \text{ cm}^3$

$$x < 5 \Rightarrow \text{Volume} < 14\,440$$

$$\Rightarrow \text{Volume} = 14\,439.9$$

(vi) If $5 \leq x \leq 15$, minimum volume occurs at $x = 15$ (using graph)

$$\Rightarrow x = 15 \Rightarrow \text{Volume} = [96 - 4(15)][48 - 2(15)][15]$$

$$\Rightarrow (36)(18)(15) = 9\,720 \text{ cm}^3$$

(c) Area = $2(x)(96 - 4x) + 2(x)(48 - 2x) + 2(96 - 4x)(48 - 2x)$

$$= 8x(24 - x) + 4x(24 - x) + 8(24 - x)(24 - x)$$

$$= 4(24 - x)(2x + x + 48 - 2x)$$

$$= 4(24 - x)(48 + x)$$

$$\Rightarrow a = 4, b = 24, c = 48$$

4. (i) $4l + 3w = 120$

$$\Rightarrow 4l = 120 - 3w$$

$$\Rightarrow l = 30 - \frac{3}{4}w$$

Hence, area = $2lw$

$$= 2w\left(30 - \frac{3}{4}w\right)$$

$$= 60w - \frac{3}{4}w^2$$

(ii) $60w - \frac{3}{4}w^2 = 0$




$$\Rightarrow w\left(60 - \frac{3}{4}w\right) = 0$$

$$\Rightarrow w = 0, 60 - \frac{3}{4}w = 0$$

$$120 - 3w = 0$$

$$\Rightarrow 3w = 120$$

$$\Rightarrow w = 40$$



w =	0	10	20	30	40
A =	0	450	600	450	0

$$\begin{aligned}
 \text{(iii)} \quad A &= -\frac{3}{2}w^2 + 60w \\
 &= -\frac{3}{2}(w^2 - 40w) \\
 &= -\frac{3}{2}(w^2 - 40w + 400) + 600 \\
 &= 600 - \frac{3}{2}(w - 20)^2
 \end{aligned}$$

Hence, maximum area = 600 m²

(iv) $w = 20$ m

(v) Dimensions are: $w = 20$ m, $l = 30 - \frac{3}{4}(20) = 15$ m.

5. (a) (i) $h = 5 \Rightarrow t = 1, 3$

(ii) $t = 4.5$ secs

(b) $2 + 4t - t^2 = 0$

$$\Rightarrow t = \frac{-4 \pm \sqrt{(4)^2 - 4(-1)(2)}}{2(-1)} = \frac{-4 \pm \sqrt{16 + 8}}{-2}$$

$$\Rightarrow t = \frac{-4 + \sqrt{24}}{-2} = 2 + \sqrt{6} \text{ (valid), } t = 2 - \sqrt{6} = -0.449 \text{ (Not valid)}$$

$$\Rightarrow t = 4.449$$

$$\Rightarrow t = 4.45$$

(c) $h = 2 + 4t - t^2$

$$= 2 - (t^2 - 4t)$$

$$= 2 - (t^2 - 4t + 4) + 4$$

$$= 6 - (t - 2)^2 \Rightarrow (p, q) = (2, 6)$$

6. No. of Price hikes	Price per Rental	Number of Rentals	Total Income (I)
	€12	36	€432
1 Price hike	€12.50	34	€425
2 Price hikes	€13	32	€416
3 Price hikes	€13.50	30	€405
x price hikes	€(12 + 0.5x)	36 - 2x	€(12 + 0.5x)(36 - 2x)

(i) $I = (12 + 0.5x)(36 - 2x) = 432 - 6x - x^2$

(ii) $I = 432 - 6x - x^2$

$$= 432 - (x^2 + 6x)$$

$$= 432 - (x^2 + 6x + 9) + 9$$

$$= 441 - (x + 3)^2$$

(iii) Maximum Income = €441

(iv) Price per rental

7. (a) Area of garden = 2 Area BAC + Area ACED

$$= 2\left(\frac{\pi}{4}x^2\right) + (x)(y)$$

$$= \frac{\pi}{4}x^2 + xy$$

(b) (i) Length = $2\left(\frac{2\pi x}{4}\right) + y = 100$

$$\Rightarrow \pi x + y = 100$$

$$\Rightarrow y = 100 - \pi x$$

(ii) Area = $\frac{\pi}{2}x^2 + x(100 - \pi x)$

$$= \frac{\pi}{2}x^2 + 100x - \pi x^2$$

$$A = 100x - \frac{\pi}{2}x^2$$

$$(iii) 100 - \pi x = 0$$

$$\Rightarrow \pi x = 100$$

$$\Rightarrow x = \frac{100}{\pi}$$

$$\text{Domain: } 0 < x < \frac{100}{\pi}$$

$$(c) \text{ Area} = 100x - \frac{\pi}{2}x^2 = 1000$$

$$\Rightarrow 200x - \pi x^2 = 2000$$

$$\Rightarrow \pi x^2 - 200x + 2000 = 0$$

$$\Rightarrow x = \frac{200 \pm \sqrt{(-200)^2 - 4(\pi)(2000)}}{2(\pi)}$$

$$= \frac{200 \pm \sqrt{40000 - 8000\pi}}{2\pi}$$

$$= \frac{200 \pm \sqrt{14867.25877}}{2\pi}$$

$$= \frac{200 + 121.9314}{2\pi}, \frac{200 - 121.9314}{2\pi}$$

$$= 51.236, 12.425$$

$$= 51.2, 12.4$$

$$(d) (i) \text{ Volume} = (\text{Area})(\text{height})$$

$$= \left(100x - \frac{\pi}{2}x^2\right)\left(\frac{x}{50}\right)$$

$$= 2x^2 - \frac{\pi}{100}x^3$$

$$(ii) x = 12.4 \Rightarrow \text{Volume} = 2(12.4)^2 - \frac{\pi}{100}(12.4)^3$$

$$= 307.52 - 59.898$$

$$= 247.62 = 247.6 \text{ m}^3$$

$$x = 51.2 \Rightarrow \text{Volume} = 2(51.2)^2 - \frac{\pi}{100}(51.2)^3$$

$$= 5242.88 - 4216.574$$

$$= 1026.306 = 1026.3 \text{ m}^3$$

$$(iii) \text{ Volume} = 2x^2 - \frac{\pi}{100}x^3 = 500$$

$$\Rightarrow 200x^2 - \pi x^3 = 50000$$

$$\Rightarrow f(x) = \pi x^3 - 200x^2 + 50000$$

$$\Rightarrow f(18) = 18321.77 - 64800 + 50000 = 3521.77$$

$$\Rightarrow f(18.8) = \pi(18.8)^3 - 200(18.8)^2 + 50000$$

$$= 20874.85 - 70688 + 50000 = 186$$

$$\Rightarrow f(18.9) = \pi(18.9)^3 - 200(18.9)^2 + 50000$$

$$= 21209.74 - 71442 + 50000 = -232.26$$

$$\text{Hence, } x = 18.8$$

$$8. (a) y = 2x \cap y = 6 + 5x - x^2$$

$$\Rightarrow 2x = 6 + 5x - x^2$$

$$\Rightarrow x^2 - 3x - 6 = 0$$

$$\Rightarrow x = \frac{3 \pm \sqrt{(3)^2 - 4(1)(-6)}}{2(1)}$$

$$= \frac{3 \pm \sqrt{33}}{2}$$

$$\Rightarrow x_1 = \frac{3 + \sqrt{33}}{2} \Rightarrow y_1 = 2\left(\frac{3 + \sqrt{33}}{2}\right) = 3 + \sqrt{33}$$

$$\text{Hence, } A = \left(\frac{3 + \sqrt{33}}{2}, 3 + \sqrt{33} \right)$$

$$\Rightarrow x_2 = \frac{3 - \sqrt{33}}{2} \Rightarrow y_2 = 2 \left(\frac{3 - \sqrt{33}}{2} \right) = 3 - \sqrt{33}$$


$$\text{Hence, } B = \left(\frac{3 - \sqrt{33}}{2}, 3 - \sqrt{33} \right)$$

(b) $(x, 6 + 5x - x^2)$ and $(x, 2x)$

$$\begin{aligned} d &= \sqrt{(x - x)^2 + (6 + 5x - x^2 - 2x)^2} \\ &= \sqrt{(6 + 3x - x^2)^2} \\ \Rightarrow d(x) &= 6 + 3x - x^2 \end{aligned}$$

(c)

$x =$	-1	0	1	2	3	4
$d(x) =$	2	6	8	8	6	2



(d) $d(x) = 6 + 3x - x^2$

$$\begin{aligned} &= 6 - \left(x^2 - 3x - \left(1\frac{1}{2} \right)^2 \right) + 2\frac{1}{4} \\ &= 8\frac{1}{4} - \left(x - 1\frac{1}{2} \right)^2 \end{aligned}$$

(e) Maximum point = $\left(1\frac{1}{2}, 8\frac{1}{4} \right)$

$$\Rightarrow \text{Maximum vertical distance} = 8\frac{1}{4} \text{ occurs at } x = 1\frac{1}{2}$$

(f) $0 \leq d(x) \leq 8\frac{1}{4}$

9. (i) $y = x \cap y = \sqrt{x - b} + c$

$$\Rightarrow x = \sqrt{x - b} + c$$

$$\Rightarrow x - c = \sqrt{x - b}$$

$$\Rightarrow (x - c)^2 = x - b$$

$$\Rightarrow x^2 - 2cx + c^2 - x + b = 0$$

$$\Rightarrow x^2 - x(2c + 1) + c^2 + b = 0$$

$$\text{At } (a, a) \Rightarrow x = a \Rightarrow a^2 - a(2c + 1) + c^2 + b = 0$$

(ii) Tangent $\Rightarrow b^2 - 4ac = 0$

$$\Rightarrow [-(2c + 1)]^2 - 4(1)(c^2 + b) = 0$$


$$\Rightarrow 4c^2 + 4c + 1 - 4c^2 - 4b = 0$$

$$\Rightarrow 4c = 4b - 1$$

$$\Rightarrow c = \frac{4b - 1}{4}$$

(iii)

$x =$	0	1	2	3	4
$y =$	-0.25	0.75	1.16	1.48	1.75



$$(iv) y = x \cap y = \sqrt{x} - \frac{1}{4}$$

$$\Rightarrow \sqrt{x} - \frac{1}{4} = x$$

$$\Rightarrow \sqrt{x} = x + \frac{1}{4}$$

$$\Rightarrow x = \left(x + \frac{1}{4}\right)^2 = x^2 + \frac{1}{2}x + \frac{1}{16}$$

$$\Rightarrow x^2 - \frac{1}{2}x + \frac{1}{16} = 0$$

$$\Rightarrow \left(x + \frac{1}{4}\right)^2 = 0 \Rightarrow x = -\frac{1}{4} \Rightarrow y = -\frac{1}{4} \Rightarrow \left(-\frac{1}{4}, -\frac{1}{4}\right)$$

$$(v) y = x + k \cap y = \sqrt{x} - \frac{1}{4}$$

$$\Rightarrow x + k = \sqrt{x} - \frac{1}{4}$$

$$\Rightarrow x + \left(k + \frac{1}{4}\right) = \sqrt{x}$$

$$\Rightarrow x^2 + 2\left(k + \frac{1}{4}\right)x + k^2 + \frac{1}{2}k + \frac{1}{16} = x$$

$$\Rightarrow x^2 + \left(2k - \frac{1}{2}\right)x + k^2 + \frac{1}{2}k + \frac{1}{16} = 0$$

$$(a) \text{ Twice } \Rightarrow b^2 - 4ac > 0$$

$$\Rightarrow \left(2k - \frac{1}{2}\right)^2 - 4(1)\left(k^2 + \frac{1}{2}k + \frac{1}{16}\right) > 0$$

$$\Rightarrow 4k^2 - 2k + \frac{1}{4} - 4k^2 - 2k - \frac{1}{4} > 0$$

$$\Rightarrow -4k^2 > 0$$

$$\Rightarrow k > 0$$

$$(b) \text{ Once } \Rightarrow b^2 - 4ac = 0$$

$$\Rightarrow k = 0$$

$$(c) \text{ At no point } \Rightarrow b^2 - 4ac < 0$$

$$\Rightarrow k > 0$$

Chapter 12

Exercise 12.1

1. (i) $3x - 5 > x + 3$

$$\Rightarrow 2x > 8$$

$$\Rightarrow x > 4$$

(ii) $6x - 5 \leq 2x - 1$

$$\Rightarrow 4x \leq 4$$

$$\Rightarrow x \leq 1$$

(iii) $1 - 3x > 10$

$$\Rightarrow -3x > 9$$


$$\Rightarrow 3x < -9$$

$$\Rightarrow x < -3$$

2. (i) $\frac{x}{2} + 2 < 7, x \in \mathbb{N}$

$$\Rightarrow \frac{x}{2} < 5$$

$$\Rightarrow x < 10$$




(ii) $\frac{1}{6}(x - 1) \geq \frac{1}{3}(x - 4), x \in \mathbb{Z}$

$$\Rightarrow 1(x - 1) \geq 2(x - 4)$$

$$\Rightarrow x - 1 \geq 2x - 8$$

$$\Rightarrow -x \geq -7$$

$$\Rightarrow x \leq 7$$




(iii) $\frac{4 - x}{2} > \frac{2 - x}{3}, x \in \mathbb{R}$

$$\Rightarrow 3(4 - x) > 2(2 - x)$$

$$\Rightarrow 12 - 3x > 4 - 2x$$

$$\Rightarrow -x > -8$$

$$\Rightarrow x < 8$$




3. (i) $12x - 3(x - 3) < 45, x \in \mathbb{R}$

$$\Rightarrow 12x - 3x + 9 < 45$$

$$\Rightarrow 9x < 36$$

$$\Rightarrow x < 4$$




(ii) $x(x - 4) \geq x^2 + 2, x \in \mathbb{R}$

$$\Rightarrow x^2 - 4x \geq x^2 + 2$$

$$\Rightarrow -4x \geq 2$$

$$\Rightarrow 4x \leq -2$$

$$\Rightarrow x \leq -\frac{1}{2}$$




(iii) $x - 2(5 + 2x) < 11$

$$\Rightarrow x - 10 - 4x < 11$$


$$\Rightarrow -3x < 21$$

$$\Rightarrow 3x > -21$$


$$\Rightarrow x > -7$$




4. (i) $-2 \leq x + 1 \leq 3, x \in R$
 $\Rightarrow -3 \leq x \leq 2$



(ii) $13 > 1 - 3x \geq 7, x \in R$
 $\Rightarrow 12 > -3x \geq 6$
 $\Rightarrow -12 < 3x \leq 6$
 $\Rightarrow -4 < x \leq 2$



(iii) $3 \geq 4x + 1 > -1$
 $\Rightarrow 2 \geq 4x > -2$
 $\Rightarrow \frac{1}{2} \geq x > -\frac{1}{2}$
 $\Rightarrow -\frac{1}{2} < x \leq \frac{1}{2}$



5. (i) $3 > \frac{3}{5}(x - 2) > 0$
 $\Rightarrow 15 > 3x - 6 > 0$
 $\Rightarrow 21 > 3x > 6$
 $\Rightarrow 7 > x > 2$
 $\Rightarrow 2 < x < 7$

(ii) $-4 \leq \frac{2}{5}(1 - 3x) \leq 1$
 $\Rightarrow -20 \leq 2 - 6x \leq 5$
 $\Rightarrow -22 \leq -6x \leq 3$
 $\Rightarrow 22 \geq 6x \geq -3$
 $\Rightarrow 3\frac{2}{3} \geq x \geq -\frac{1}{2}$
 $\Rightarrow -\frac{1}{2} \leq x \leq 3\frac{2}{3}$


(iii) $3 \leq 2 - \frac{x}{7} < 4$
 $\Rightarrow 21 \leq 14 - x \leq 28$
 $\Rightarrow 7 \leq -x \leq 14$
 $\Rightarrow -7 \geq x \geq -14$
 $\Rightarrow -14 \leq x \leq -7$

6. $3(x - 2) > x - 4$ and $4x + 12 > 2x + 17, x \in R$
 $\Rightarrow 3x - 6 > x - 4 \quad \Rightarrow 2x > 2$
 $\Rightarrow 2x > 2 \quad \Rightarrow x > 1$
 $\Rightarrow x > 1$
 $\Rightarrow \text{ANS: } x > 2\frac{1}{2}$

7. (i) $2x - 5 < x - 1, x \in R$
 $\Rightarrow x < 4$

(ii) $7(x + 1) > 23 - x, x \in R$
 $\Rightarrow 7x + 7 > 23 - x$
 $\Rightarrow 8x > 16$
 $\Rightarrow x > 2$


(iii) $2 < x < 4, x \in R$



8. (i) $2x - 3 > 2, x \in R$
 $\Rightarrow 2x > 5$
 $\Rightarrow x > 2\frac{1}{2}$

$$\begin{aligned}
 \text{(ii)} \quad & 3(x+2) < 12+x, x \in R \\
 \Rightarrow & 3x+6 < 12+x \\
 \Rightarrow & 2x < 6 \\
 \Rightarrow & x < 3
 \end{aligned}$$

$$\text{(iii)} \quad 2\frac{1}{2} < x < 3, x \in R$$



$$\begin{aligned}
 \text{9. (i)} \quad & 15-x < 2(11-x), x \in Z \\
 \Rightarrow & 15-x < 22-2x \\
 \Rightarrow & x < 7
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad & 5(3x-1) > 12x+19, x \in Z \\
 \Rightarrow & 15x-5 > 12x+19 \\
 \Rightarrow & 3x > 24 \\
 \Rightarrow & x > 8
 \end{aligned}$$

(iii) \therefore Null set

$$\begin{aligned}
 \text{10. (i)} \quad & 3x+8 \leq 20, x \in N \\
 \Rightarrow & 3x \leq 12 \\
 \Rightarrow & x \leq 4
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii)} \quad & 2(3x-7) \geq x+6, x \in N \\
 \Rightarrow & 6x-14 \geq x+6 \\
 \Rightarrow & 5x \geq 20 \\
 \Rightarrow & x \geq 4
 \end{aligned}$$

$$\text{(iii)} \quad x = 4$$

$$\begin{aligned}
 \text{11.} \quad & \text{Width} = x, \text{ Length} = x+1 \\
 \text{Perimeter} &= 2(x) + 2(x+1) \leq 38 \\
 \Rightarrow & 2x + 2x + 2 \leq 38 \\
 \Rightarrow & 4x \leq 36 \\
 \Rightarrow & x \leq 9
 \end{aligned}$$

Width = 9 m, length = 10 m

$$\begin{aligned}
 \text{12.} \quad & 100 < 2^n < 200 \\
 \Rightarrow & 2^{6.645} < 2^n < 2^{7.645} \\
 \Rightarrow & 6.645 < n < 7.645
 \end{aligned}$$

Hence $a = 6.645, b = 7.645, n = 7$

$$\begin{aligned}
 \text{13. Example 1: } & 4 > 3 > 0 \quad \text{Thus } 4^2 > 3^2 \\
 \text{Example 2: } & 4 > 3 > 0 \quad \text{Thus } 4^{-2} < 3^{-2}, \text{ as } \frac{1}{16} < \frac{1}{9}.
 \end{aligned}$$

- (i) if $a < b < 0$ and $n(\text{odd}) > 0$, then $a^n < b^n$.
- (ii) if $a < b < 0$ and $n(\text{even}) > 0$, then $a^n > b^n$.
- (iii) if $a < b < 0$ and $n(\text{odd}) < 0$, then $a^n > b^n$.
- (iv) if $a < b < 0$ and $n(\text{even}) < 0$, then $a^n < b^n$.

$$\begin{aligned}
 \text{14.} \quad & 5-3x < -10 \quad \cap \quad 4x+6 < 32, \quad x \in Z \\
 \Rightarrow & -3x < -15 \quad \cap \quad 4x < 26 \\
 \Rightarrow & 3x > 15 \quad \cap \quad x < 6\frac{1}{2} \\
 \Rightarrow & x > 5 \quad \cap \quad x < 6\frac{1}{2}, \quad x \in Z \\
 \Rightarrow & x = 6
 \end{aligned}$$


Exercise 12.2

1. (i) Let $x^2 - x - 6 = 0$

$$\Rightarrow (x + 2)(x - 3) = 0$$

$$\Rightarrow \text{Roots: } x = -2, 3$$

$$\text{Hence } x^2 - x - 6 \geq 0 \Rightarrow x \leq -2 \text{ or } x \geq 3$$




(ii) Let $x^2 + 3x - 10 = 0$

$$\Rightarrow (x + 5)(x - 2) = 0$$

$$\Rightarrow \text{Roots: } x = -5, 2$$

$$\text{Hence } x^2 + 3x - 10 \leq 0 \Rightarrow -5 \leq x \leq 2$$




(iii) Let $2x^2 - 5x + 2 = 0$

$$\Rightarrow (2x - 1)(x - 2) = 0$$

$$\Rightarrow \text{Roots: } x = \frac{1}{2}, 2$$

$$\text{Hence } 2x^2 - 5x + 2 < 0 \Rightarrow \frac{1}{2} < x < 2$$




2. (i) Let $6 - x - x^2 = 0$

$$\Rightarrow (3 + x)(2 - x) = 0$$

$$\Rightarrow \text{Roots: } x = -3, 2$$

$$\text{Hence } 6 - x - x^2 \geq 0 \Rightarrow -3 \leq x \leq 2$$




(ii) Let $12 - 5x - 2x^2 = 0$

$$\Rightarrow (4 + x)(3 - 2x) = 0$$

$$\Rightarrow \text{Roots: } x = -4, 1\frac{1}{2}$$

$$\text{Hence } 12 - 5x - 2x^2 > 0 \Rightarrow -4 < x < 1\frac{1}{2}$$




(iii) Let $-2x^2 - 7x = 0$

$$\Rightarrow 2x^2 + 7x = 0$$

$$\Rightarrow x(2x + 7) = 0$$

$$\Rightarrow \text{Roots: } x = 0, -3\frac{1}{2}$$

$$\text{Hence } -2x^2 - 7x \geq 0 \Rightarrow -3\frac{1}{2} \leq x \leq 0$$




3. (i) Let $6x^2 - x - 15 = 0$

$$\Rightarrow (2x + 3)(3x - 5) = 0$$

$$\Rightarrow \text{Roots: } x = -1\frac{1}{2}, 1\frac{2}{3}$$

$$\text{Hence } 6x^2 - x - 15 > 0 \Rightarrow x < -1\frac{1}{2} \text{ or } x > 1\frac{2}{3}$$




(ii) Let $16 - x^2 = 0$

$$\Rightarrow (4 + x)(4 - x) = 0$$

$$\Rightarrow \text{Roots: } x = -4, 4$$

$$\text{Hence } 16 - x^2 \leq 0 \Rightarrow x \leq -4 \text{ or } x \geq 4$$



(iii) Let $2(x^2 - 6) = 5x$


$$\Rightarrow 2x^2 - 12 = 5x$$

$$\Rightarrow 2x^2 - 5x - 12 = 0$$

$$\Rightarrow (2x + 3)(x - 4) = 0$$

$$\Rightarrow \text{Roots: } x = -1\frac{1}{2}, 4$$

$$\text{Hence } 2(x^2 - 6) \geq 5x \Rightarrow x \leq -1\frac{1}{2} \text{ or } x \geq 4$$



4. Let $(4 - x)(1 - x) = x + 11$


$$\Rightarrow 4 - 5x + x^2 - x - 11 = 0$$

$$\Rightarrow x^2 - 6x - 7 = 0$$

$$\Rightarrow (x + 1)(x - 7) = 0$$

$$\Rightarrow \text{Roots: } x = -1, 7$$

$$\text{Hence } (4 - x)(1 - x) < x + 11 \Rightarrow -1 < x < 7$$



5. Let $x^2 - 6x + 2 = 0$

$$\Rightarrow \text{Roots: } x = \frac{6 \pm \sqrt{(-6)^2 - 4(1)(2)}}{2(1)}$$


$$= \frac{6 \pm \sqrt{36 - 8}}{2}$$

$$= \frac{6 \pm \sqrt{28}}{2}$$

$$= \frac{6 \pm 2\sqrt{7}}{2}$$

$$= 3 - \sqrt{7}, 3 + \sqrt{7}$$

$$\text{Hence } x^2 - 6x + 2 \leq 0 \Rightarrow 3 - \sqrt{7} \leq x \leq 3 + \sqrt{7}$$



6. $x^2 + (k + 1)x + 1 = 0$

$$\text{Real Roots } \Rightarrow b^2 - 4ac \geq 0$$

$$\Rightarrow (k + 1)^2 - 4(1)(1) \geq 0$$

$$\Rightarrow k^2 + 2k + 1 - 4 \geq 0$$


$$\Rightarrow k^2 + 2k + 1 - 3 \geq 0$$

$$\text{Solve } k^2 + 2k - 3 = 0$$

$$\Rightarrow (k + 3)(k - 1) = 0$$

$$\Rightarrow \text{Roots} = -3, 1$$

$$\text{Hence } k^2 + 2k - 3 \geq 0 \Rightarrow k \leq -3 \text{ or } k \geq 1$$



7. $kx^2 + 4x + 3 + k = 0$

$$\text{Real Roots } \Rightarrow b^2 - 4ac \geq 0$$

$$\Rightarrow (4)^2 - 4(k)(3 + k) \geq 0$$

$$\Rightarrow 16 - 12k - 4k^2 \geq 0$$


$$\Rightarrow k^2 + 3k - 4 \leq 0$$

$$\text{solve } k^2 + 3k - 4 = 0$$

$$\Rightarrow (k + 4)(k - 1) = 0$$

$$\Rightarrow \text{Roots: } k = -4, 1$$

$$\text{Hence } k^2 + 3k - 4 \leq 0 \Rightarrow -4 \leq k \leq 1$$



8. $px^2 + (p+3)x + p = 0$


$$\begin{aligned}\text{Real Roots} &\Rightarrow b^2 - 4ac \geq 0 \\ &\Rightarrow (p+3)^2 - 4(p)(p) \geq 0 \\ &\Rightarrow p^2 + 6p + 9 - 4p^2 \geq 0 \\ &\Rightarrow -3p^2 + 6p + 9 \geq 0 \\ &\Rightarrow p^2 - 2p - 3 \leq 0\end{aligned}$$

Solve $p^2 - 2p - 3 = 0$

$$\begin{aligned}&\Rightarrow (p+1)(p-3) = 0 \\ &\Rightarrow \text{Roots: } p = -1, 3\end{aligned}$$

Hence $p^2 - 2p - 3 \leq 0 \Rightarrow -1 \leq p \leq 3$

$$\begin{aligned}x = -2 &\Rightarrow p(-2)^2 + (p+3)(-2) + p = 0 \\ &\Rightarrow 4p - 2p - 6 + p = 0 \\ &\Rightarrow 3p = 6 \\ &\Rightarrow p = 2\end{aligned}$$




9. (i) $\frac{x+3}{x+2} < 2, x \neq -2$

$$\begin{aligned}&\Rightarrow \frac{x+3}{x+2}(x+2)^2 < 2(x+2)^2 \\ &\Rightarrow (x+3)(x+2) < 2(x^2 + 4x + 4) \\ &\Rightarrow x^2 + 5x + 6 < 2x^2 + 8x + 8 \\ &\Rightarrow x^2 + 3x + 2 > 0\end{aligned}$$

Solve $x^2 + 3x + 2 = 0$

$$\begin{aligned}&\Rightarrow (x+2)(x+1) = 0 \\ &\Rightarrow \text{Roots } x = -2, -1\end{aligned}$$

Hence $x^2 + 3x + 2 > 0 \Rightarrow x < -2 \text{ or } x > -1$



(ii) $\frac{x+5}{x-3} > 1, x \neq 3$

$$\begin{aligned}&\Rightarrow \frac{x+5}{x-3}(x-3)^2 > 1(x-3)^2 \\ &\Rightarrow (x+5)(x-3) > 1(x^2 - 6x + 9) \\ &\Rightarrow x^2 + 2x - 15 > x^2 - 6x + 9 \\ &\Rightarrow 8x > 24 \\ &\Rightarrow x > 3\end{aligned}$$


(iii) $\frac{2x-1}{x+3} > 3, x \neq -3$

$$\begin{aligned}&\Rightarrow \frac{2x-1}{x+3}(x+3)^2 > 3(x+3)^2 \\ &\Rightarrow (2x-1)(x+3) > 3(x^2 + 6x + 9) \\ &\Rightarrow 2x^2 + 5x - 3 > 3x^2 + 18x + 27 \\ &\Rightarrow x^2 + 13x + 30 < 0\end{aligned}$$

Solve $x^2 + 13x + 30 = 0$


$$\begin{aligned}&\Rightarrow (x+10)(x+3) = 0 \\ &\Rightarrow x = -10, -3\end{aligned}$$

Hence $x^2 + 13x + 30 < 0 \Rightarrow -10 < x < -3$



10. (i) $\frac{3x+4}{x-5} > 2, x \neq 5$

$$\begin{aligned}&\Rightarrow \frac{3x+4}{x-5}(x-5)^2 > 2(x-5)^2 \\ &\Rightarrow (3x+4)(x-5) > 2(x^2 - 10x + 25) \\ &\Rightarrow 3x^2 - 11x - 20 > 2x^2 - 20x + 50 \\ &\Rightarrow x^2 + 9x - 70 > 0\end{aligned}$$



Solve $x^2 + 9x - 70 = 0$

$\Rightarrow (x + 14)(x - 5) = 0$

$\Rightarrow x = -14, 5$

Hence $x^2 + 9x - 70 > 0 \Rightarrow x < -14 \text{ or } x > 5$

(ii) $\frac{1-2x}{4x+2} > 2, x \neq -\frac{1}{2}$

$\Rightarrow \frac{1-2x}{4x+2}(4x+2)^2 > 2(4x+2)^2$

$\Rightarrow (1-2x)(4x+2) > 2(16x^2 + 16x + 4)$

$\Rightarrow 4x + 2 - 8x^2 - 4x > 32x^2 + 32x + 8$

$\Rightarrow 40x^2 + 32x + 6 < 0$


$\Rightarrow 20x^2 + 16x + 3 < 0$

Solve $20x^2 + 16x + 3 = 0$

$\Rightarrow (2x + 1)(10x + 3) = 0$

$\Rightarrow \text{Roots: } x = -\frac{1}{2}, -\frac{3}{10}$

Hence $20x^2 + 16x + 3 < 0 \Rightarrow -\frac{1}{2} < x < -\frac{3}{10}$



(iii) $\frac{3+4x}{5x-1} > 3, x \neq \frac{1}{5}$

$\Rightarrow \frac{3+4x}{5x-1}(5x-1)^2 > 3(5x-1)^2$

$\Rightarrow (3+4x)(5x-1) > 3(25x^2 - 10x + 1)$

$\Rightarrow 15x - 3 + 20x^2 - 4x > 75x^2 - 30x + 3$


$\Rightarrow 55x^2 - 41x + 6 > 0$

Solve $55x^2 - 41x + 6 = 0$

$\Rightarrow (5x - 1)(11x - 6) = 0$

$\Rightarrow \text{Roots: } x = \frac{1}{5}, \frac{6}{11}$

Hence $55x^2 - 41x + 6 < 0 \Rightarrow \frac{1}{5} < x < \frac{6}{11}$



11. (i) $\frac{x}{2x-3} \leq 1, x \neq \frac{3}{2}$

$\Rightarrow \frac{x}{2x-3}(2x-3)^2 \leq 1(2x-3)^2$

$\Rightarrow x(2x-3) \leq 1(4x^2 - 12x + 9)$

$\Rightarrow 2x^2 - 3x \leq 4x^2 - 12x + 9$


$\Rightarrow 2x^2 - 9x + 9 \geq 0$

Solve $2x^2 - 9x + 9 = 0$

$\Rightarrow (2x - 3)(x - 3) = 0$

$\Rightarrow \text{Roots: } x = 1\frac{1}{2}, 3$

Hence $2x^2 - 9x + 9 \geq 0 \Rightarrow x \leq 1\frac{1}{2} \text{ or } x \geq 3$



(ii) $\frac{2x-4}{x-1} < 1, x \neq 1$

$\Rightarrow \frac{2x-4}{x-1}(x-1)^2 < 1(x-1)^2$

$\Rightarrow (2x-4)(x-1) < 1(x^2 - 2x + 1)$

$\Rightarrow 2x^2 - 6x + 4 < x^2 - 2x + 1$


$\Rightarrow x^2 - 4x + 3 < 0$

Solve $x^2 - 4x + 3 = 0$

$\Rightarrow (x - 1)(x - 3) = 0$

$\Rightarrow x = 1, 3$

Hence $x^2 - 4x + 3 < 0 \Rightarrow 1 < x < 3$



$$(iii) \quad \frac{x-5}{x-1} \leq 3, \quad x \neq 1$$

$$\Rightarrow \frac{x-5}{x-1}(x-1)^2 \leq 3(x-1)^2$$

$$\Rightarrow (x-5)(x-1) \leq 3(x^2-2x+1)$$

$$\Rightarrow x^2-6x+5 \leq 3x^2-6x+3$$

$$\Rightarrow 2x^2-2 \geq 0$$


$$\Rightarrow x^2-1 \geq 0$$

$$\text{Solve } x^2-1=0$$

$$\Rightarrow (x+1)(x-1)=0$$

$$\Rightarrow \text{Roots: } x = -1, 1$$

$$\text{Hence } x^2-1 \geq 0 \Rightarrow x \leq -1 \text{ or } x \geq 1$$



$$12. (i) \quad \frac{2x-7}{x+3} < 1, \quad x \neq -3$$

$$\Rightarrow \frac{2x-7}{x+3}(x+3)^2 < 1(x+3)^2$$

$$\Rightarrow (2x-7)(x+3) < 1(x^2+6x+9)$$

$$\Rightarrow 2x^2-x-21 < x^2+6x+9$$


$$\Rightarrow x^2-7x-30 < 0$$

$$\text{Solve } x^2-7x-30=0$$

$$\Rightarrow (x+3)(x-10)=0$$

$$\Rightarrow \text{Roots: } x = -3, 10$$

$$\text{Hence } x^2-7x-30 < 0 \Rightarrow -3 < x < 10$$



$$(ii) \quad \frac{2x-3}{x-5} < \frac{3}{2}, \quad x \neq 5$$

$$\Rightarrow \frac{2x-3}{x-5}(x-5)^2 < \frac{3}{2}(x-5)^2$$

$$\Rightarrow (2x-3)(x-5) < \frac{3}{2}(x^2-10x+25)$$

$$\Rightarrow 2x^2-13x+15 < \frac{3x^2-30x+75}{2}$$

$$\Rightarrow 4x^2-26x+30 < 3x^2-30x+75$$


$$\Rightarrow x^2+4x-45 < 0$$

$$\text{Solve } x^2+4x-45=0$$

$$\Rightarrow (x+9)(x-5)=0$$

$$\Rightarrow x = -9, 5$$

$$\text{Hence } x^2+4x-45 < 0 \Rightarrow -9 < x < 5$$



$$(iii) \quad \frac{x+2}{x-1} \leq 3, \quad x \neq 1$$

$$\Rightarrow \frac{x+2}{x-1}(x-1)^2 \leq 3(x-1)^2$$

$$\Rightarrow (x+2)(x-1) \leq 3(x^2-2x+1)$$

$$\Rightarrow x^2+x-2 \leq 3x^2-6x+3$$


$$\Rightarrow 2x^2-7x+5 \geq 0$$

$$\text{Solve } 2x^2-7x+5=0$$

$$\Rightarrow (x-1)(2x-5)=0$$


$$\Rightarrow \text{Roots: } x = 1, 2\frac{1}{2}$$

$$\text{Hence } 2x^2-7x+5 \geq 0 \Rightarrow x \leq 1 \text{ or } x \geq 2\frac{1}{2}$$





- 13.** From graph: $-3 > x > -2$
 Using inequality $\Rightarrow 2x^2 + 4x > x^2 - x - 6, \quad x \in R$
 $\Rightarrow x^2 + 5x + 6 > 0$

Solve $x^2 + 5x + 6 = 0$
 $\Rightarrow (x + 3)(x + 2) = 0$
 \Rightarrow Roots: $x = -3, -2$
 Hence $x^2 + 5x + 6 > 0 \Rightarrow x < -3$ or $x > -2$




- 14.** No real roots if $b^2 - 4ac < 0$
 $x^2 + x + 1 = 0 \Rightarrow b^2 - 4ac = (1)^2 - 4(1)(1) = -3 < 0$
 Hence $x^2 + x + 1 > 0$
 $\Rightarrow x^2 + x + \frac{1}{4} + \frac{3}{4} > 0$
 $\Rightarrow \left(x + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2 > 0$ True

- 15.** (i) $f(t) \leq 4 \Rightarrow -11 + 13t - 2t^2 \leq 4$
 $\Rightarrow 2t^2 - 13t + 15 \geq 0$
 Solve $2t^2 - 13t + 15 = 0$
 $\Rightarrow (2t - 3)(t - 5) = 0$
 $\Rightarrow t = 1\frac{1}{2}, 5$
 Hence $2t^2 - 13t + 15 \geq 0 \Rightarrow t \leq 1\frac{1}{2}$ or $t \geq 5$
 (ii) $f(t) \geq 7 \Rightarrow -11 + 13t - 2t^2 \geq 7$
 $\Rightarrow 2t^2 - 13t + 18 \leq 0$
 $\Rightarrow (t - 2)(2t - 9) = 0$
 \Rightarrow Roots: $t = 2, 4\frac{1}{2}$
 Hence $2t^2 - 13t + 18 \leq 0 \Rightarrow 2 \leq t \leq 4\frac{1}{2}$



- (iii) $4 < f(t) < 7 \Rightarrow$
 $1\frac{1}{2} < t < 2$ and $4\frac{1}{2} < t < 5$




- 16.** (i) $x < -3$ or $x > -\frac{1}{2}$
 (ii) $x \leq 1$ or $x \geq 3$
 (iii) $-1\frac{1}{2} \leq x \leq \frac{1}{2}$
 (iv) $-1 < x < 5$

- 17.** Length = x and Width = $x - 3$

Ratio $< 5 \Rightarrow \frac{x}{x-3} < 5$
 $\Rightarrow \frac{x}{x-3}(x-3)^2 < 5(x-3)^2$
 $\Rightarrow x(x-3) < 5(x^2 - 6x + 9)$
 $\Rightarrow x^2 - 3x < 5x^2 - 30x + 45$
 $\Rightarrow 4x^2 - 27x + 45 > 0$

Solve $4x^2 - 27x + 45 = 0$
 $\Rightarrow (x - 3)(4x - 15) = 0$
 \Rightarrow Roots: $x = 3, 3.75$
 Hence $4x^2 - 27x + 45 > 0 \Rightarrow x < 3$ or $x > 3.75$
 Since $x < 3$ is not valid, hence (i) Length > 3.75
 (ii) Width > 0.75




18. Positive graphs \Rightarrow No real roots
 $\Rightarrow b^2 - 4ac < 0$
 $x^2 - 2px + p + 6 = 0 \Rightarrow (-2p)^2 - 4(1)(p + 6) < 0$
 $\Rightarrow 4p^2 - 4p - 24 < 0$
 $\Rightarrow p^2 - p - 6 < 0$

Solve $p^2 - p - 6 = 0$

$\Rightarrow (p + 2)(p - 3) = 0$

\Rightarrow Roots: $p = -2, 3$

Hence $p^2 - p - 6 < 0 \Rightarrow -2 < p < 3$



19. (i) Perimeter $< 50 \Rightarrow 2(x + 3) + 2(x + 2) < 50$
 $\Rightarrow 2x + 6 + 2x + 4 < 50$
 $\Rightarrow 4x < 50 - 10$
 $\Rightarrow 4x < 40$
 $\Rightarrow x < 10$

- (ii) Area $> 12 \Rightarrow (x + 3)(x + 2) > 12$
 $\Rightarrow x^2 + 5x + 6 > 12$
 $\Rightarrow x^2 + 5x - 6 > 0$


Solve $x^2 + 5x - 6 = 0$

$\Rightarrow (x + 6)(x - 1) = 0$

$\Rightarrow x = -6$ (Not valid), $x = 1$ (valid)

Hence $x^2 + 5x - 6 > 0 \Rightarrow x > 1$

- (iii) $1 < x < 10$, by combining (i) and (ii) above



20. Use Pythagoras $\Rightarrow h^2 = x^2 + 3^2 = x^2 + 9$
 $\Rightarrow h = \sqrt{x^2 + 9}$
Perimeter $> 8 \Rightarrow \sqrt{x^2 + 9} + x + 3 > 8$
 $\Rightarrow \sqrt{x^2 + 9} > -x + 5$
 $\Rightarrow x^2 + 9 > (-x + 5)^2$
 $\Rightarrow x^2 + 9 > x^2 - 10x + 25$
 $\Rightarrow 10x > 16$
 $\Rightarrow x > 1.6$
Perimeter $< 12 \Rightarrow \sqrt{x^2 + 9} + x + 3 < 12$
 $\Rightarrow \sqrt{x^2 + 9} < -x + 9$
 $\Rightarrow x^2 + 9 < (-x + 9)^2$
 $\Rightarrow x^2 + 9 < x^2 - 18x + 81$
 $\Rightarrow 18x < 72$
 $\Rightarrow x < 4$

Hence $1.6 < x < 4$

Since $x \in \mathbb{Z} \Rightarrow x = 2 \text{ m or } 3 \text{ m}$

Exercise 12.3

1. (i) $x + 3 = -1$ OR $x + 3 = 1$
 $\Rightarrow x = -4$ OR $x = -2$
(ii) $x - 2 = -4$ OR $x - 2 = 4$
 $\Rightarrow x = -2$ OR $x = 6$
(iii) $2x - 1 = -5$ OR $2x - 1 = 5$
 $\Rightarrow 2x = -4$ OR $2x = 6$
 $\Rightarrow x = -2$ OR $x = 3$


$$\begin{aligned}
 \text{(iv)} \quad & 3x - 2 = -x \quad \text{OR} \quad 3x - 2 = x \\
 & \Rightarrow 4x = 2 \quad \text{OR} \quad 2x = 2 \\
 & \Rightarrow x = \frac{1}{2} \quad \text{OR} \quad x = 1 \\
 \text{(v)} \quad & 2(x - 3) = -2 \quad \text{OR} \quad 2(x - 3) = 2 \\
 & \Rightarrow x - 3 = -1 \quad \text{OR} \quad x - 3 = 1 \\
 & \Rightarrow x = 2 \quad \text{OR} \quad x = 4 \\
 \text{(vi)} \quad & x - 5 = -(x + 1) \quad \text{OR} \quad x - 5 = +(x + 1) \\
 & \Rightarrow x - 5 = -x - 1 \quad \text{OR} \quad x - 5 = x + 1 \\
 & \Rightarrow 2x = 4 \quad \text{OR} \quad -5 = +1 \text{ (Not valid)} \\
 & \Rightarrow x = 2
 \end{aligned}$$

2. Copy and complete the following table and hence sketch a graph of $f(x) = |3x - 2|$

x	-3	-2	-1	0	1	2	3
$f(x) = 3x - 2 $	11	8	5	2	1	4	7

Solve $|3x - 2| = 5$.

$$\Rightarrow x = -1, 2\frac{1}{3}$$



3. $f(x) = |x|$, $g(x) = |x - 4|$, $h(x) = |x + 3|$
 $f(-2) = |-2| = 2 \Rightarrow \text{point } (-2, 2) \in f(x)$
 $g(2) = |2 - 4| = |-2| = 2 \Rightarrow \text{point } (2, 2) \in g(x)$
 $h(-5) = |-5 + 3| = |-2| = 2 \Rightarrow \text{point } (-5, 2) \in h(x)$

4. 2 points on $f(x)$ are $(-1, 0)$ and $(0, 1)$
 $f(-1) = |a(-1) + b| = 0$ and $f(0) = |a(0) + b| = 1$
 $\Rightarrow -a + b = 0 \quad \Rightarrow b = 1$
Hence, $-a + 1 = 0$
 $\Rightarrow a = 1$

Hence, $f(x) = |x + 1|$.

2 points on $g(x)$ are $(-1, 0)$ and $(0, 2)$

$$\begin{aligned}
 g(-1) &= |a(-1) + b| = 0 \quad \text{and} \quad g(0) = |a(0) + b| = 2 \\
 \Rightarrow -a + b &= 0 \quad \Rightarrow b = 2 \\
 \text{Hence } -a + 2 &= 0 \\
 \Rightarrow a &= 2
 \end{aligned}$$

Hence, $g(x) = |2x + 2|$

2 points on $h(x)$ are $(-1, 0)$ and $(0, 3)$

$$\begin{aligned}
 h(-1) &= |a(-1) + b| = 0 \quad \text{and} \quad h(0) = |a(0) + b| = 3 \\
 \Rightarrow -a + b &= 0 \quad \Rightarrow b = 3 \\
 \text{Hence } -a + 3 &= 0 \\
 \Rightarrow a &= 3
 \end{aligned}$$

Hence, $h(x) = |3x + 3|$

$$x = -2 \Rightarrow f(-2) = |-2 + 1| = 1 \Rightarrow (-2, 1) \in f(x)$$

$$x = -2 \Rightarrow g(-2) = |2(-2) + 2| = 2 \Rightarrow (-2, 2) \in g(x)$$

$$x = -2 \Rightarrow h(-2) = |3(-2) + 3| = 3 \Rightarrow (-2, 3) \in h(x)$$

5. $f(x) = |x - 2|$

$$\Rightarrow f(0) = |0 - 2| = 2 \quad (0, 2) \in f(x)$$

$$\Rightarrow f(2) = |2 - 2| = 0 \quad (2, 0) \in f(x)$$

$$\Rightarrow f(4) = |4 - 2| = 2 \quad (4, 2) \in f(x)$$

$$g(x) = |x - 6|$$

$$\Rightarrow g(0) = |0 - 6| = 6 \quad (0, 6) \in g(x)$$

$$\Rightarrow g(6) = |6 - 6| = 0 \quad (6, 0) \in g(x)$$

$$\Rightarrow g(8) = |8 - 6| = 2 \quad (8, 2) \in g(x)$$

Hence, $|x - 2| = |x - 6|$ at $x = 4$.


Solve $|x - 2| = |x - 6|$

$$\Rightarrow x - 2 = -(x - 6) \quad \text{OR} \quad x - 2 = +(x - 6)$$

$$\Rightarrow x - 2 = -x + 6 \quad \text{OR} \quad x - 2 = x - 6$$

$$\Rightarrow 2x = 8 \quad \text{OR} \quad -2 = -6 \text{ (Not valid)}$$

$$\Rightarrow x = 4$$



6. (i) $|x - 6| < 2$

$$\Rightarrow -2 < x - 6 < 2$$

$$\Rightarrow 4 < x < 8$$

(ii) $|x + 2| \leq 4$

$$\Rightarrow -4 \leq x + 2 \leq 4$$

$$\Rightarrow -6 \leq x \leq 2$$

(iii) $|2x - 1| \geq 5$

$$\Rightarrow -5 \geq 2x - 1 \geq 5$$

$$\Rightarrow -4 \geq 2x \geq 6 \quad \Rightarrow -4 \geq 2x \quad \text{or} \quad 2x \geq 6$$

$$\Rightarrow 2x \leq -4 \quad \text{or} \quad x \geq 3$$

$$\Rightarrow x \leq -2 \quad \text{or} \quad x \geq 3$$

(iv) $|2x - 1| \geq 11$

$$\Rightarrow -11 \geq 2x - 1 \geq 11$$

$$\Rightarrow -10 \geq 2x \geq 12 \quad \Rightarrow -10 \geq 2x \quad \text{or} \quad 2x \geq 12$$

$$\Rightarrow 2x \leq -10 \quad \text{or} \quad x \geq 6$$

$$\Rightarrow x \leq -5 \quad \text{or} \quad x \geq 6$$

(v) $|3x + 5| < 4$

$$\Rightarrow -4 < 3x + 5 < 4$$

$$\Rightarrow -9 < 3x < -1$$

$$\Rightarrow -3 < x < -\frac{1}{3}$$

(vi) $|x - 4| < 3$

$$\Rightarrow -3 < x - 4 < 3$$

$$\Rightarrow 1 < x < 7$$

7. (i) $|2x - 1| \geq 7$

$$\Rightarrow -7 \geq 2x - 1 \geq 7$$

$$\Rightarrow -6 \geq 2x \geq 8$$

$$\Rightarrow -3 \geq x \geq 4$$

(ii) $|3x + 4| \leq |x + 2|$

$$\Rightarrow (3x + 4)^2 \leq (x + 2)^2$$

$$\Rightarrow 9x^2 + 24x + 16 \leq x^2 + 4x + 4$$

$$\Rightarrow 8x^2 + 20x + 12 \leq 0$$

$$\Rightarrow 2x^2 + 5x + 3 \leq 0$$

$$\text{Solve } 2x^2 + 5x + 3 = 0$$

$$\Rightarrow (2x + 3)(x + 1) = 0$$

$$\Rightarrow \text{Roots: } x = -1\frac{1}{2}, -1$$

$$\text{Hence, } 2x^2 + 5x + 3 \leq 0 \Rightarrow -1\frac{1}{2} \leq x \leq -1$$

$$(iii) \quad 2|x - 1| \leq |x + 3|$$

$$\Rightarrow [2(x - 1)]^2 \leq (x + 3)^2$$

$$\Rightarrow 4(x^2 - 2x + 1) \leq x^2 + 6x + 9$$

$$\Rightarrow 4x^2 - 8x + 4 \leq x^2 + 6x + 9$$

$$\Rightarrow 3x^2 - 14x - 5 \leq 0$$

$$\text{Solve } 3x^2 - 14x - 5 = 0$$

$$\Rightarrow (3x + 1)(x - 5) = 0$$

$$\Rightarrow \text{Roots: } x = -\frac{1}{3}, 5$$

$$\text{Hence, } 3x^2 - 14x - 5 \leq 0 \Rightarrow -\frac{1}{3} \leq x \leq 5$$

8. $f(x) = |x| - 4$

$$f(-4) = |-4| - 4 = 4 - 4 = 0 \quad (-4, 0) \in f(x)$$

$$f(0) = |0| - 4 = -4 \quad (0, -4) \in f(x)$$

$$f(4) = |4| - 4 = 4 - 4 = 0 \quad (4, 0) \in f(x)$$

$$g(x) = \frac{1}{2}x$$

$$g(-4) = \frac{1}{2}(-4) = -2 \quad (-4, -2) \in g(x)$$

$$g(0) = \frac{1}{2}(0) = 0 \quad (0, 0) \in g(x)$$

$$g(4) = \frac{1}{2}(4) = 2 \quad (4, 2) \in g(x)$$

$$f(x) \leq g(x) \Rightarrow |x| - 4 \leq \frac{1}{2}x$$

$$\text{Solve } |x| = \frac{1}{2}x + 4$$

$$\Rightarrow x^2 = \frac{1}{4}x^2 + 4x + 16$$

$$\Rightarrow 4x^2 = x^2 + 16x + 64$$

$$\Rightarrow 3x^2 - 16x - 64 = 0$$

$$\Rightarrow (3x + 8)(x - 8) = 0$$

$$\Rightarrow \text{Roots } x = -2\frac{2}{3}, 8$$

$$\text{Hence, } 3x^2 - 16x - 64 \leq 0 \Rightarrow -2\frac{2}{3} \leq x \leq 8$$



9. $f(x) = \left| \frac{1}{4}x + 3 \right|$

$$f(-24) = \left| \frac{1}{4}(-24) + 3 \right| = |-6 + 3| = |-3| = 3 \Rightarrow (-24, 3) \in f(x)$$

$$f(-12) = \left| \frac{1}{4}(-12) + 3 \right| = |-3 + 3| = 0 \Rightarrow (-12, 0) \in f(x)$$

$$f(0) = \left| \frac{1}{4}(0) + 3 \right| = |0 + 3| = 3 \Rightarrow (0, 3) \in f(x)$$

$$\text{Hence, } \left| \frac{1}{4}x + 3 \right| \geq 3 \Rightarrow x \leq -24 \text{ or } x \geq 0$$



$$\begin{aligned}
 10. \quad & |1 + 2x| < |x + 2| \\
 & \Rightarrow (1 + 2x)^2 < (x + 2)^2 \\
 & \Rightarrow 1 + 4x + 4x^2 < x^2 + 4x + 4 \\
 & \Rightarrow 3x^2 - 3 < 0 \\
 & \Rightarrow x^2 - 1 < 0 \\
 & \text{Solve } x^2 - 1 = 0 \\
 & \Rightarrow (x + 1)(x - 1) = 0 \\
 & \Rightarrow \text{Roots: } x = -1, 1 \\
 & \text{Hence, } x^2 - 1 < 0 \Rightarrow -1 < x < 1
 \end{aligned}$$

$$\begin{aligned}
 11. \quad & \left| \frac{1}{1 + 2x} \right| = 1 \Rightarrow \left(\frac{1}{1 + 2x} \right)^2 = (1)^2 \\
 & \Rightarrow \frac{1}{1 + 4x + 4x^2} = \frac{1}{1} \\
 & \Rightarrow 1 + 4x + 4x^2 = 1 \\
 & \Rightarrow 4x^2 + 4x = 0 \\
 & \Rightarrow x^2 + x = 0 \\
 & \Rightarrow x(x + 1) = 0 \\
 & \Rightarrow \text{Roots: } x = 0, -1 \\
 & \text{Solve } \left| \frac{1}{1 + 2x} \right| < 1 \\
 & \text{Hence, } \frac{1}{1 + 4x + 4x^2} < 1 \\
 & \Rightarrow 1 + 4x + 4x^2 > 1 \\
 & \Rightarrow 4x^2 + 4x > 0 \\
 & \Rightarrow x^2 + x > 0 \\
 & \Rightarrow -1 > x > 0
 \end{aligned}$$

$$\begin{aligned}
 12. \quad & \text{(i) } -4 < x < 2 \\
 & \text{(ii) } x < -4 \text{ or } x > 2 \\
 & \text{(iii) } 1\frac{1}{4} < x < 3\frac{1}{2} \\
 & \text{(iv) } 2 < x < 3 \\
 & \text{(v) } 1\frac{1}{4} < x < 2 \\
 & \text{(vi) } 2 < x < 3 \\
 & \text{(vii) } 3 < x < 3\frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 13. \quad & \text{(i) } \frac{x}{2x - 1} < -2 \\
 & \Rightarrow + \left(\frac{x}{2x - 1} \right) < -2 \\
 & \Rightarrow \frac{x}{2x - 1} (2x - 1)^2 < -2(2x - 1)^2 \\
 & \Rightarrow x(2x - 1) < -2(4x^2 - 4x + 1) \\
 & \Rightarrow 2x^2 - x < -8x^2 + 8x - 2 \\
 & \Rightarrow 10x^2 - 9x + 2 < 0 \\
 & \text{Solve } 10x^2 - 9x + 2 = 0 \\
 & \Rightarrow (5x - 2)(2x - 1) = 0 \\
 & \Rightarrow \text{Roots: } x = \frac{2}{5}, \frac{1}{2} \\
 & \text{Hence, } 10x^2 - 9x + 2 < 0 \Rightarrow \frac{2}{5} < x < \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
\text{(ii)} \quad & |x - 3| = 2|x - 1| \\
& \Rightarrow (x - 3)^2 = [2(x - 1)]^2 \\
& \Rightarrow x^2 - 6x + 9 = 4(x^2 - 2x + 1) \\
& \Rightarrow x^2 - 6x + 9 = 4x^2 - 8x + 4 \\
& \Rightarrow 3x^2 - 2x - 5 = 0 \\
& \Rightarrow (x + 1)(3x - 5) = 0 \\
& \Rightarrow \text{Roots: } x = -1, 1\frac{2}{3} \\
\text{(iii)} \quad & |x - 1| - |2x + 1| > 0 \\
& \Rightarrow |x - 1| > |2x + 1| \\
& \Rightarrow (x - 1)^2 > (2x + 1)^2 \\
& \Rightarrow x^2 - 2x + 1 > 4x^2 + 4x + 1 \\
& \Rightarrow 3x^2 + 6x < 0 \\
& \Rightarrow x^2 + 2x < 0 \\
& \text{Solve } x^2 + 2x = 0 \\
& \Rightarrow x(x + 2) = 0 \\
& \Rightarrow \text{Roots: } x = -2, 0 \\
& \text{Hence, } x^2 + 2x < 0 \Rightarrow -2 < x < 0
\end{aligned}$$

Exercise 12.4

1. Proof: Assume x and y are both positive integers
and $x^2 - y^2 = (x + y)(x - y)$.

If $x > y$, then $(x + y)$ and $(x - y)$ are positive integers;
hence, $(x + y)(x - y)$ is a positive integer $\neq 1$.

If $x < y$, then $(x + y)$ is a positive integer
and $(x - y)$ is a negative integer;
hence, $(x + y)(x - y)$ is a negative integer $\neq 1$.

Hence, there is a contradiction in both cases.

\therefore There are no positive integer solutions to $x^2 - y^2 = 1$.

2. Proof: Assume $(a + b)$ is a rational number.

Hence, $(a + b)$ can be written as $\frac{p}{q}$ where $p, q \in \mathbb{Z}, q \neq 0$.

Since " a " is a rational number $= \frac{m}{n}$ where $m, n \in \mathbb{Z}, n \neq 0$

$$\text{Then, } a + b = \frac{p}{q}$$

$$\Rightarrow \frac{m}{n} + b = \frac{p}{q}$$

$$\Rightarrow b = \frac{p}{q} - \frac{m}{n}$$

$$= \frac{pn - mq}{qn} \text{ which is rational.}$$

This is a contradiction as b is an irrational number;
hence, $a + b$ is an irrational number.

- 3. Proof:** Assume x and y are both positive integers

$$\text{and } x^2 - y^2 = (x + y)(x - y).$$

If $x > y$, then $(x + y)$ and $(x - y)$ they are positive integers;

hence, $(x + y)(x - y)$ is a positive integer $\neq 10$.

Note: $x + y = 5$ and $x + y = 10$

$$\frac{x - y = 2}{2x = 7} \quad \frac{x - y = 1}{2x = 11}$$

$$\Rightarrow x = 3\frac{1}{2} \notin Z \quad \Rightarrow x = 5\frac{1}{2} \in Z$$

If $x < y$, then $(x + y)$ is a positive integer

and $(x - y)$ is a negative integer;

hence, $(x + y)(x - y)$ is a negative integer $\neq 10$.

Hence, there is a contradiction in both cases.

\therefore There are no positive integer solutions to $x^2 - y^2 = 10$.

- 4. Proof:** a divides $b \Rightarrow b = k(a), k \in N$

$$b \text{ divides } c \Rightarrow c = m(b), m \in N$$

$$\Rightarrow c = m[k(a)]$$

$$\Rightarrow c = mk(a), m, k \in N$$

Then a divides c .

- 5. Proof:** a divides $b \Rightarrow b = k(a), k \in N$

$$a \text{ divides } c \Rightarrow c = m(a), m \in N$$

$$\text{hence, } (b + c) = k(a) + m(a)$$

$$= (k + m)(a), k, m \in N$$

Then a divides $(b + c)$.

- 6. Proof:** $a^2 + b^2 \geq 2ab$

$$\Rightarrow a^2 - 2ab + b^2 \geq 0$$

$$\Rightarrow (a - b)(a - b) \geq 0$$

$$\Rightarrow (a - b)^2 \geq 0, \text{ true for } a, b \in R$$

Hence, $a^2 + b^2 \geq 2ab$.

- 7. Proof:** a is a rational number

Hence, a can be written as $\frac{p}{q}$ where $p, q \in Z, q \neq 0$.

b is a rational number.

Hence, b can be written as $\frac{m}{n}$ where $m, n \in Z, n \neq 0$.

$$\text{Hence, } a + b = \frac{p}{q} + \frac{m}{n} = \frac{pn + mq}{qn}$$

= a rational number, as $p, q, m, n \in Z$
 $qn \neq 0$.

- 8. Proof:** x is an odd number $= 2a + 1$, where $a \in N$.

y is an odd number $= 2b + 1$, where $b \in N$.

$$\text{Hence, } x + y = 2a + 1 + 2b + 1$$

$$= 2a + 2b + 2$$

$$= 2(a + b + 1)$$

Hence, $x + y$ has a factor of 2.

$\Rightarrow x + y$ must be even.

Hence, the sum of 2 odd numbers is always even.

Exercise 12.5

1. (i) $a^2 + 2ab + b^2 \geq 0$
 $\Rightarrow (a + b)(a + b) \geq 0$
 $\Rightarrow (a + b)^2 \geq 0$, true for $a, b \in R$
- (ii) $a^2 + 2ab + b^2 \geq 0$
 $\Rightarrow a^2 + 2ab + b^2 + b^2 \geq 0$
 $\Rightarrow (a + b)^2 + (b)^2 \geq 0$, true for $a, b \in R$
2. $(a + b)^2 \geq 4ab$
 $\Rightarrow a^2 + 2ab + b^2 - 4ab \geq 0$
 $\Rightarrow a^2 - 2ab + b^2 \geq 0$
 $\Rightarrow (a - b)^2 \geq 0$, true for $a, b \in R$
3. $-(a^2 + 2ab + b^2) \leq 0$
 $\Rightarrow a^2 + 2ab + b^2 \geq 0$
 $\Rightarrow (a + b)^2 \geq 0$, true for $a, b \in R$
4. (i) $a + \frac{1}{a} \geq 2$
 $\Rightarrow a^2 + 1 \geq 2a$
 $\Rightarrow a^2 - 2a + 1 \geq 0$
 $\Rightarrow (a - 1)^2 \geq 0$, true for $a > 0$ and $a \in R$
- (ii) $\frac{1}{a} + \frac{1}{b} \geq \frac{2}{a + b}$
 $(a)(b)(a + b) \cdot \frac{1}{a} + (a)(b)(a + b) \cdot \frac{1}{b} \geq (a)(b)(a + b) \cdot \frac{2}{a + b}$
 $\Rightarrow b(a + b) + a(a + b) \geq 2ab$
 $\Rightarrow ab + b^2 + a^2 + ab \geq 2ab$
 $\Rightarrow a^2 + b^2 \geq 0$, true for $a > 0, b > 0$ and $a, b \in R$
5. $a^2 - 6a + 9 + b^2 \geq 0$
 $\Rightarrow (a - 3)(a - 3) + b^2 \geq 0$
 $\Rightarrow (a - 3)^2 + b^2 \geq 0$, true for $a, b \in R$
6. (i) $x^2 + 6x + 9 \geq 0$
 $\Rightarrow (x + 3)^2 \geq 0$, true for $x \in R$
- (ii) $x^2 - 10x + 25 \geq 0$
 $\Rightarrow (x - 5)^2 \geq 0$, true for $x \in R$
- (iii) $x^2 + 4x + 6 \geq 0$
 $\Rightarrow x^2 + 4x + 4 + 2 \geq 0$
 $\Rightarrow (x + 2)^2 + 2 \geq 0$, true for $x \in R$
- (iv) $x^2 - 6x + 10 \geq 0$
 $\Rightarrow x^2 - 6x + 9 + 1 \geq 0$
 $\Rightarrow (x - 3)^2 + 1 \geq 0$, true for $x \in R$
- (v) $4x^2 + 12x + 11 \geq 0$
 $\Rightarrow x^2 + 3x + \frac{11}{4} \geq 0$
 $\Rightarrow x^2 + 3x + \frac{9}{4} + \frac{2}{4} \geq 0$
 $\Rightarrow \left(x + 1\frac{1}{2}\right)^2 + \frac{1}{2} \geq 0$, true for $x \in R$

$$\begin{aligned}
 \text{(vi)} \quad & 4x^2 - 4x + 2 \geq 0 \\
 \Rightarrow & x^2 - x + \frac{1}{2} \geq 0 \\
 \Rightarrow & x^2 - x + \frac{1}{4} + \frac{1}{4} \geq 0 \\
 \Rightarrow & \left(x - \frac{1}{2}\right)^2 + \frac{1}{4} \geq 0, \text{ true for } x \in R
 \end{aligned}$$

$$\begin{aligned}
 \text{7. (i)} \quad & -x^2 + 10x - 25 \geq 0 \\
 \Rightarrow & x^2 - 10x + 25 \geq 0 \\
 \Rightarrow & (x - 5)^2 \geq 0, \text{ true for } x \in R \\
 \text{(ii)} \quad & -x^2 - 4x - 7 \leq 0 \\
 \Rightarrow & x^2 + 4x + 7 \geq 0 \\
 \Rightarrow & x^2 + 4x + 4 + 3 \geq 0 \\
 \Rightarrow & (x + 2)^2 + 3 \geq 0, \text{ true for } x \in R
 \end{aligned}$$

$$\begin{aligned}
 \text{8. (i)} \quad & p^2 + 4q^2 \geq 4pq \\
 \Rightarrow & p^2 - 4pq + 4q^2 \geq 0 \\
 \Rightarrow & (p - 2q)(p - 2q) \geq 0 \\
 \Rightarrow & (p - 2q)^2 \geq 0, \text{ true for } p, q \in R \\
 \text{(ii)} \quad & (p + q)^2 \leq 2(p^2 + q^2) \\
 \Rightarrow & p^2 + 2pq + q^2 \leq 2p^2 + 2q^2 \\
 \Rightarrow & -p^2 + 2pq - q^2 \leq 0 \\
 \Rightarrow & p^2 - 2pq + q^2 \geq 0 \\
 \Rightarrow & (p - q)^2 \geq 0, \text{ true for } p, q \in R
 \end{aligned}$$

$$\begin{aligned}
 \text{9. } a^3 + b^3 &= (a + b)(a^2 - ab + b^2) \\
 \text{Proof: } a^3 + b^3 &> a^2b + ab^2 \\
 \Rightarrow a^3 + b^3 - a^2b - ab^2 &> 0 \\
 \Rightarrow (a + b)(a^2 - ab + b^2) - ab(a + b) &> 0 \\
 \Rightarrow (a + b)(a^2 - ab + b^2 - ab) &> 0 \\
 \Rightarrow (a + b)(a - b)^2 > 0, &\text{ true for } a > 0, b > 0
 \end{aligned}$$

$$\text{10. Given } a^2 + b^2 \geq 2ab.$$

$$\text{(i) } a^2 + c^2 \geq 2ac$$

$$\text{(ii) } b^2 + c^2 \geq 2bc$$

$$\begin{aligned}
 & a^2 + b^2 \geq 2ab \\
 & a^2 + c^2 \geq 2ac \\
 & b^2 + c^2 \geq 2bc \\
 \text{Add: } \Rightarrow & \frac{2a^2 + 2b^2 + 2c^2}{2} \geq 2ab + 2bc + 2ac \\
 \Rightarrow & a^2 + b^2 + c^2 \geq ab + bc + ac
 \end{aligned}$$

$$\begin{aligned}
 \text{11. } & \frac{p + q}{2} > \sqrt{pq} \\
 \Rightarrow & p + q > 2\sqrt{pq} \\
 \Rightarrow & (p + q)^2 > (2\sqrt{pq})^2 \\
 \Rightarrow & p^2 + 2pq + q^2 > 4pq \\
 \Rightarrow & p^2 - 2pq + q^2 > 0 \\
 \Rightarrow & (p - q)^2 > 0 \dots \text{true}
 \end{aligned}$$

$$\begin{aligned}
 \text{12. } & (ax + by)^2 \leq (a^2 + b^2)(x^2 + y^2) \\
 \Rightarrow & a^2x^2 + 2axby + b^2y^2 \leq a^2x^2 + a^2y^2 + b^2x^2 + b^2y^2 \\
 \Rightarrow & -a^2y^2 + 2axby - b^2x^2 \leq 0 \\
 \Rightarrow & a^2y^2 - 2axby + b^2x^2 \geq 0 \\
 \Rightarrow & (ay - bx)^2 \geq 0, \text{ true for } a, b, x, y \in R
 \end{aligned}$$

$$\begin{aligned}
 13. \quad & a^4 + b^4 \geq 2a^2b^2 \\
 \Rightarrow & a^4 - 2a^2b^2 + b^4 \geq 0 \\
 \Rightarrow & (a^2 - b^2)(a^2 - b^2) \geq 0 \\
 \Rightarrow & (a^2 - b^2)^2 \geq 0, \text{ true for } a, b \in R
 \end{aligned}$$

$$\begin{aligned}
 14. \quad & (a + 2b)\left(\frac{1}{a} + \frac{1}{2b}\right) \geq 4 \\
 \Rightarrow & a \cdot \frac{1}{a} + \frac{a}{2a} + \frac{2b}{a} + 2b \cdot \frac{1}{2b} \geq 4 \\
 \Rightarrow & 1 + \frac{a}{2b} + \frac{2b}{a} + 1 \geq 4 \\
 \Rightarrow & \frac{a}{2b} + \frac{2b}{a} \geq 2 \\
 \Rightarrow & 2ab \cdot \frac{a}{2b} + 2ab \cdot \frac{2b}{a} \geq 2ab \cdot 2 \dots \text{ since } a > 0, b > 0 \\
 \Rightarrow & a^2 + 4b^2 \geq 4ab \\
 \Rightarrow & a^2 - 4ab + 4b^2 \geq 0 \\
 \Rightarrow & (a - 2b)^2 \geq 0, \text{ true for } a, b \text{ positive}
 \end{aligned}$$

$$\begin{aligned}
 15. \quad & \frac{a}{(a+1)^2} \leq \frac{1}{4} \\
 \Rightarrow & \frac{a}{(a+1)^2} \cdot (a+1)^2 \leq \frac{1}{4}(a+1)^2 \\
 \Rightarrow & a \leq \frac{a^2 + 2a + 1}{4} \\
 \Rightarrow & 4a \leq a^2 + 2a + 1 \\
 \Rightarrow & -a^2 + 2a - 1 \leq 0 \\
 \Rightarrow & a^2 - 2a + 1 \geq 0 \\
 \Rightarrow & (a - 1)^2 \geq 0, \text{ true for } a \in R
 \end{aligned}$$

$$\begin{aligned}
 16. \quad (i) \quad & a^4 - b^4 = (a^2 + b^2)(a^2 - b^2) \\
 & = (a^2 + b^2)(a + b)(a - b) \\
 (ii) \quad & a^5 - a^4b - ab^4 + b^5 \\
 & = a^4(a - b) - b^4(a - b) \\
 & = (a - b)(a^4 - b^4) \\
 & = (a - b)(a^2 + b^2)(a + b)(a - b) \\
 & = (a^2 + b^2)(a + b)(a - b)^2 \\
 (iii) \quad & a^5 + b^5 > a^4b + ab^4 \\
 \Rightarrow & a^5 - a^4b - ab^4 + b^5 > 0 \\
 \Rightarrow & (a^2 + b^2)(a + b)(a - b)^2 > 0, \text{ true if } a \text{ and } b \\
 & \text{are positive unequal real numbers.}
 \end{aligned}$$

$$\begin{aligned}
 17. \text{ Given: } & a^2 + b^2 = 1 \quad \text{and} \quad c^2 + d^2 = 1. \\
 & a^2 + b^2 \geq 2ab \Rightarrow a^2 - 2ab + b^2 \geq 0 \\
 & \Rightarrow (a - b)^2 \geq 0
 \end{aligned}$$

$$\begin{aligned}
 \text{Hence,} \quad & (a - b)^2 + (c - d)^2 \geq 0 \\
 \Rightarrow & a^2 - 2ab + b^2 + c^2 - 2cd + d^2 \geq 0 \\
 \Rightarrow & 1 - 2ab + 1 - 2cd \geq 0 \\
 \Rightarrow & -2ab - 2cd \geq -2 \\
 \Rightarrow & 2ab + 2cd \leq 2 \\
 \Rightarrow & ab + cd \leq 1
 \end{aligned}$$

$$\begin{aligned}
 18. \quad & \sqrt{ab} > \frac{2ab}{a+b} \\
 \Rightarrow & (a+b) \cdot \sqrt{ab} > (a+b) \frac{2ab}{a+b} \text{ if } a \text{ and } b \text{ are positive and unequal} \\
 \Rightarrow & (a+b)\sqrt{ab} > 2ab \\
 \Rightarrow & (a+b) > \frac{2ab}{\sqrt{ab}} \\
 \Rightarrow & a+b > 2\sqrt{ab} \\
 \Rightarrow & (a+b)^2 > (2\sqrt{ab})^2 \\
 \Rightarrow & a^2 + 2ab + b^2 > 4ab \\
 \Rightarrow & a^2 - 2ab + b^2 > 0 \\
 \Rightarrow & (a-b)^2 > 0, \text{ true if } a \text{ and } b \text{ are positive and unequal.}
 \end{aligned}$$

$$\begin{aligned}
 19. \quad & a + \frac{9}{a+2} \geq 4 \\
 \Rightarrow & a(a+2) + \frac{9}{a+2}(a+2) \geq 4(a+2), \text{ where } (a+2) > 0 \\
 \Rightarrow & a^2 + 2a + 9 \geq 4a + 8 \\
 \Rightarrow & a^2 - 2a + 1 \geq 0 \\
 \Rightarrow & (a-1)^2 \geq 0, \text{ true where } (a+2) > 0
 \end{aligned}$$

$$\begin{aligned}
 20. \quad & \text{If } \frac{a}{b} > \frac{c}{d}, \\
 \Rightarrow & \frac{a}{b}(bd) > \frac{c}{d}(bd) \text{ if } a, b, c, d \text{ are positive numbers} \\
 \Rightarrow & ad > cd \\
 \text{Hence,} \quad & \frac{a+c}{b+d} > \frac{c}{d} \\
 \Rightarrow & \frac{(a+c)}{(b+d)} \cdot (d)(b+d) > \frac{c}{d}(d)(b+d) \text{ if } a, b, c, d \text{ are positive numbers} \\
 \Rightarrow & ad + cd > cb + cd \\
 \Rightarrow & ad > cb, \text{ true}
 \end{aligned}$$

$$\begin{aligned}
 21. \quad & (a^3 - b^3)(a - b) = (a - b)(a^2 + ab + b^2)(a - b) \\
 & = (a - b)^2(a^2 + ab + b^2) \\
 & \text{If } a > b \Rightarrow (a - b) > 0. \\
 & \text{Hence, } (a - b)^2 \text{ is positive,} \\
 & \text{and } a^2 + ab + b^2 > a^2 - 2ab + b^2 = (a - b)^2 > 0 \\
 & \text{Hence, } a^2 + ab + b^2 \text{ is a positive, true if } a, b, c, d \text{ are positive numbers.} \\
 & \text{Hence,} \quad a^4 + b^4 \geq a^3b + ab^3 \\
 \Rightarrow & a^4 - a^3b - ab^3 + b^4 \geq 0 \\
 \Rightarrow & a^3(a - b) - b^3(a - b) \geq 0 \\
 \Rightarrow & (a - b)(a^3 - b^3) \geq 0 \\
 \Rightarrow & (a - b)(a - b)(a^2 + ab + b^2) \geq 0 \\
 \Rightarrow & (a - b)^2(a^2 + ab + b^2) \geq 0, \text{ true for } a, b \in R \text{ and } a > b
 \end{aligned}$$

Exercise 12.6

1.
 - (i) $a^2 \times a^3 = a^{2+3} = a^5$
 - (ii) $x \cdot x \cdot x^2 = x^{1+1+2} = x^4$
 - (iii) $2x^3 \times 3x^3 = 6x^{3+3} = 6x^6$
 - (iv) $\frac{x^5}{x^2} = x^{5-2} = x^3$
 - (v) $\frac{x^4}{x^5} = x^{4-5} = x^{-1}$
 - (vi) $a^0 = 1$
 - (vii) $\sqrt[3]{27} = 3$
 - (viii) $(a^3)^2 = a^6$
 - (ix) $\frac{(x^3)^2}{x^3} = \frac{x^6}{x^3} = x^{6-3} = x^3$
 - (x) $(3ab)^2 = 9a^2b^2$
2.
 - (i) $\sqrt[3]{64} = 4$
 - (ii) $3^{-2} = \frac{1}{3^2} = \frac{1}{9}$
 - (iii) $\frac{1}{2^{-3}} = 2^3 = 8$
 - (iv) $\frac{2^{-2}}{3^{-2}} = \frac{3^2}{2^2} = \frac{9}{4}$
 - (v) $\frac{1}{4^{-\frac{1}{2}}} = 4^{\frac{1}{2}} = 2$
3.
 - (i) $8^{\frac{2}{3}} = (2^3)^{\frac{2}{3}} = 2^2 = 4$
 - (ii) $16^{\frac{3}{4}} = (2^4)^{\frac{3}{4}} = 2^3 = 8$
 - (iii) $27^{\frac{2}{3}} = (3^3)^{\frac{2}{3}} = 3^2 = 9$
 - (iv) $81^{\frac{3}{4}} = (3^4)^{\frac{3}{4}} = 3^3 = 27$
 - (v) $125^{\frac{2}{3}} = (5^3)^{\frac{2}{3}} = 5^2 = 25$
4.
 - (i) $\left(\frac{2}{3}\right)^{-2} = \frac{1}{\left(\frac{2}{3}\right)^2} = \frac{1}{\frac{4}{9}} = \frac{9}{4}$
 - (ii) $\left(\frac{4}{9}\right)^{-\frac{1}{2}} = \frac{1}{\left(\frac{4}{9}\right)^{\frac{1}{2}}} = \frac{1}{\frac{2}{3}} = \frac{3}{2}$
 - (iii) $\left(\frac{9}{25}\right)^{-\frac{3}{2}} = \frac{1}{\left(\frac{9}{25}\right)^{\frac{3}{2}}} = \frac{1}{\left(\left(\frac{3}{5}\right)^2\right)^{\frac{3}{2}}} = \frac{1}{\left(\frac{3}{5}\right)^3} = \frac{1}{\frac{27}{125}} = \frac{125}{27}$
 - (iv) $\left(\frac{27}{125}\right)^{-\frac{2}{3}} = \left(\left(\frac{3}{5}\right)^3\right)^{-\frac{2}{3}} = \left(\frac{3}{5}\right)^{-2} = \frac{1}{\left(\frac{3}{5}\right)^2} = \frac{1}{\frac{9}{25}} = \frac{25}{9}$
 - (v) $\left(3\frac{3}{8}\right)^{\frac{1}{3}} = \left(\frac{27}{8}\right)^{\frac{1}{3}} = \frac{(27)^{\frac{1}{3}}}{(8)^{\frac{1}{3}}} = \frac{3}{2}$
5. $\frac{4^2 \times 16^{\frac{1}{2}}}{64^{\frac{2}{3}} \times 4^3} = \frac{4^2 \times 4}{(4^3)^{\frac{2}{3}} \times 4^3} = \frac{4^3}{4^2 \times 4^3} = \frac{4^3}{4^5} = 4^{3-5} = 4^{-2}$

$$6. \frac{3^{\frac{1}{4}} \times 3 \times 3^{\frac{1}{6}}}{\sqrt{3}} = \frac{3^{\frac{1}{4}+1+\frac{1}{6}}}{3^{\frac{1}{2}}} = \frac{3^{\frac{5}{12}}}{3^{\frac{1}{2}}} = 3^{\frac{5}{12}-\frac{1}{2}} = 3^{\frac{11}{12}}$$

$$3^p = 3^{\frac{11}{12}} \Rightarrow p = \frac{11}{12}$$

$$7. (i) \frac{(xy^2)^3 \times (x^2y)^{-2}}{xy} = \frac{x^3y^6 \cdot x^{-4}y^{-2}}{xy} = \frac{x^{-1}y^4}{x^1y^1}$$

$$= \frac{y^{4-1}}{x^{1+1}} = \frac{y^3}{x^2}$$

$$(ii) \left(\frac{p^2q}{p^{-1}q^3} \right)^4 = \frac{p^8q^4}{p^{-4}q^{12}} = \frac{p^{8+4}}{q^{12-4}} = \frac{p^{12}}{q^8}$$

$$(iii) a^{\frac{1}{4}} \times a^{-\frac{5}{4}} = a^{\frac{1}{4}-\frac{5}{4}} = a^{-1} = \frac{1}{a}$$

$$(iv) \left(\frac{y^{-2}}{y^{-3}} \right)^{\frac{2}{3}} = (y^{-2+3})^{\frac{2}{3}} = (y^1)^{\frac{2}{3}} = y^{\frac{2}{3}}$$

$$(v) \frac{(a\sqrt{b})^{-3}}{\sqrt{a^3b}} = \frac{(a^1b^{\frac{1}{2}})^{-3}}{(a^3b^1)^{\frac{1}{2}}} = \frac{a^{-3}b^{-\frac{3}{2}}}{a^{\frac{3}{2}}b^{\frac{1}{2}}} = \frac{1}{a^{3+\frac{3}{2}}b^{\frac{1}{2}+\frac{3}{2}}} = \frac{1}{a^{\frac{9}{2}}b^2}$$

$$(vi) \frac{\sqrt[4]{x^7}}{\sqrt{x^3}} = \frac{x^{\frac{7}{4}}}{x^{\frac{3}{2}}} = x^{\frac{7}{4}-\frac{3}{2}} = x^{\frac{1}{4}}$$

$$8. (i) \frac{x^{\frac{1}{2}} + x^{-\frac{1}{2}}}{x^{\frac{1}{2}}} = \frac{x^{-\frac{1}{2}}(x + 1)}{x^{\frac{1}{2}}} = \frac{x + 1}{x^{\frac{1}{2}+\frac{1}{2}}} = \frac{x + 1}{x}$$

$$(ii) (x + x^{\frac{1}{2}})(x - x^{\frac{1}{2}}) = x^2 - x \cdot x^{\frac{1}{2}} + x \cdot x^{\frac{1}{2}} - x^{\frac{1}{2}+\frac{1}{2}}$$

$$= x^2 - x$$

$$(iii) \frac{\sqrt{x} + \sqrt{x^3}}{\sqrt{x}} = \frac{x^{\frac{1}{2}} + x^{\frac{3}{2}}}{x^{\frac{1}{2}}} = \frac{x^{\frac{1}{2}}(1 + x)}{x^{\frac{1}{2}}} = 1 + x$$

$$9. \frac{(x-1)^{\frac{1}{2}} + (x-1)^{-\frac{1}{2}}}{(x-1)^{\frac{1}{2}}} = \frac{(x-1)^{\frac{1}{2}}(x-1)^{\frac{1}{2}} + (x-1)^{\frac{1}{2}}(x-1)^{-\frac{1}{2}}}{(x-1)^{\frac{1}{2}}(x-1)^{\frac{1}{2}}}$$

$$= \frac{(x-1)^1 + (x-1)^0}{(x-1)^1} = \frac{x-1+1}{x-1} = \frac{x}{x-1}$$

$$10. \sqrt{3^{2n+1}} \times \sqrt[3]{3^{-3n}} = (3^{2n+1})^{\frac{1}{2}} \times (3^{-3n})^{\frac{1}{3}}$$

$$= 3^{n+\frac{1}{2}} \times 3^{-n} = 3^{n+\frac{1}{2}-n} = 3^{\frac{1}{2}}$$

$$\Rightarrow 3^k = 3^{\frac{1}{2}} \Rightarrow k = \frac{1}{2}$$

$$11. 220 \cdot 2^{\frac{1}{12}} \cdot 2^{\frac{1}{12}} \cdot 2^{\frac{1}{12}} = 220 \cdot (2^{\frac{1}{4}})$$

$$= 261.626$$

$$= 262 \text{ Hz}$$

$$12. \frac{A_1}{A_2} = \frac{4\pi R_1^2}{4\pi R_2^2} = \frac{R_1^2}{R_2^2}$$

$$\left(\frac{V_1}{V_2} \right)^{\frac{2}{3}} = \left(\frac{\frac{4}{3}\pi R_1^3}{\frac{4}{3}\pi R_2^3} \right)^{\frac{2}{3}} = \left(\frac{\frac{4}{3}}{\frac{4}{3}} \right)^{\frac{2}{3}} \left(\frac{\pi}{\pi} \right)^{\frac{2}{3}} \frac{R_1^{3(\frac{2}{3})}}{R_2^{3(\frac{2}{3})}} = \frac{R_1^2}{R_2^2}$$

$$\text{Hence, } \frac{A_1}{A_2} = \left(\frac{V_1}{V_2} \right)^{\frac{2}{3}}$$

$$\text{Hence, } \frac{A_1}{A_2} = \left(\frac{162}{384} \right)^{\frac{2}{3}} = \left(\frac{27}{64} \right)^{\frac{2}{3}} = \frac{9}{16}$$

$$\begin{aligned}
 13. \quad f(n) = 3^n &\Rightarrow \quad (i) \quad f(n+3) = 3^{n+3} \\
 &\quad \text{and} \quad (ii) \quad f(n+1) = 3^{n+1} \\
 &\Rightarrow f(n+3) - f(n+1) = 3^{n+3} - 3^{n+1} \\
 &\quad = 3^n \cdot 3^3 - 3^n \cdot 3^1 \\
 &\quad = 3^n(27 - 3) = 24(3^n) \\
 &\Rightarrow k f(n) = k(3^n) = 24(3^n) \\
 &\Rightarrow k = 24
 \end{aligned}$$

$$\begin{aligned}
 14. \quad f(n) = 3^{n-1} &\Rightarrow f(n+3) + f(n) = k f(n) \\
 &\Rightarrow 3^{n+3-1} + 3^{n-1} = k(3^{n-1}) \\
 &\Rightarrow 3^{n-1} \cdot 3^3 + 3^{n-1} = k(3^{n-1}) \\
 &\Rightarrow 3^{n-1}(27 + 1) = k(3^{n-1}) \\
 &\Rightarrow 28 = k
 \end{aligned}$$

Exercise 12.7

$$\begin{aligned}
 1. \quad (i) \quad 2^x &= 32 \\
 &\Rightarrow 2^x = 2^5 \Rightarrow x = 5 \\
 (ii) \quad 16^x &= 64 \\
 &\Rightarrow (4^2)^x = 4^3 \\
 &\Rightarrow 4^{2x} = 4^3 \Rightarrow 2x = 3 \Rightarrow x = \frac{3}{2} \\
 (iii) \quad 25^x &= 125 \\
 2. \quad (i) \quad 9^x &= \frac{1}{27} \\
 &\Rightarrow (3^2)^x = \frac{1}{3^3} \\
 &\Rightarrow 3^{2x} = 3^{-3} \Rightarrow 2x = -3 \Rightarrow x = -\frac{3}{2} \\
 (ii) \quad 4^x &= \frac{1}{32} \\
 &\Rightarrow (2^2)^x = \frac{1}{2^5} \\
 &\Rightarrow 2^{2x} = 2^{-5} \Rightarrow 2x = -5 \Rightarrow x = -\frac{5}{2} \\
 (iii) \quad 4^{x-1} &= 2^{x+1} \\
 3. \quad (i) \quad 2^x &= \frac{\sqrt{2}}{2} \\
 &\Rightarrow 2^x = \frac{2^{\frac{1}{2}}}{2^1} \\
 &\Rightarrow 2^x = 2^{\frac{1}{2}-1} = 2^{-\frac{1}{2}} \\
 &\Rightarrow x = -\frac{1}{2} \\
 (ii) \quad 25^x &= \frac{125}{\sqrt{5}} \\
 &\Rightarrow (5^2)^x = \frac{5^3}{5^{\frac{1}{2}}} \\
 &\Rightarrow 5^{2x} = 5^{3-\frac{1}{2}} = 5^{\frac{5}{2}} \\
 &\Rightarrow 2x = \frac{5}{2} \\
 &\Rightarrow x = \frac{5}{4}
 \end{aligned}$$

$$\begin{aligned}
 &\Rightarrow (5^2)^x = 5^3 \\
 &\Rightarrow 5^{2x} = 5^3 \Rightarrow 2x = 3 \Rightarrow x = \frac{3}{2} \\
 (iv) \quad 3^x &= \frac{1}{27} \\
 &\Rightarrow 3^x = \frac{1}{3^3} \\
 &\Rightarrow 3^x = 3^{-3} \Rightarrow x = -3 \\
 &\Rightarrow (2^2)^{x-1} = 2^{x+1} \\
 &\Rightarrow 2^{2x-2} = 2^{x+1} \Rightarrow 2x - 2 = x + 1 \\
 &\quad \Rightarrow x = 3 \\
 (iv) \quad \frac{1}{9^x} &= 27 \\
 &\Rightarrow \frac{1}{(3^2)^x} = 3^3 \\
 &\Rightarrow \frac{1}{3^{2x}} = 3^3 \\
 &\Rightarrow 3^{-2x} = 3^3 \Rightarrow -2x = 3 \\
 &\quad \Rightarrow x = -\frac{3}{2} \\
 (iii) \quad \frac{1}{8^x} &= \sqrt{2} \\
 &\Rightarrow \frac{1}{(2^3)^x} = 2^{\frac{1}{2}} \\
 &\Rightarrow \frac{1}{2^{3x}} = 2^{\frac{1}{2}} \\
 &\Rightarrow 2^{-3x} = 2^{\frac{1}{2}} \\
 &\Rightarrow -3x = \frac{1}{2} \\
 &\Rightarrow x = -\frac{1}{6} \\
 (iv) \quad 7^x &= \frac{1}{\sqrt[3]{7}} \\
 &\Rightarrow 7^x = \frac{1}{7^{\frac{1}{3}}} \\
 &\Rightarrow 7^x = 7^{-\frac{1}{3}} \\
 &\Rightarrow x = -\frac{1}{3}
 \end{aligned}$$

$$4. \quad \sqrt{32} = (2^5)^{\frac{1}{2}} = 2^{\frac{5}{2}}$$

$$\Rightarrow 16^{x-1} = 2\sqrt{32}$$

$$\Rightarrow (2^4)^{x-1} = 2^1 \cdot 2^{\frac{5}{2}}$$

$$\Rightarrow 2^{4x-4} = 2^{1+\frac{5}{2}} = 2^{\frac{7}{2}}$$

$$\Rightarrow 4x - 4 = \frac{7}{2}$$

$$\Rightarrow 4x = \frac{7}{2} + 4 = \frac{15}{2}$$

$$\Rightarrow x = \frac{15}{8}$$

$$5. \quad 27^x = 9 \quad \text{and} \quad 2^{x-y} = 64$$

$$\Rightarrow (3^3)^x = 3^2 \quad \Rightarrow 2^{x-y} = 2^6$$

$$\Rightarrow 3^{3x} = 3^2 \quad \Rightarrow x - y = 6$$

$$\Rightarrow 3x = 2$$

$$\Rightarrow x = \frac{2}{3} \quad \Rightarrow \frac{2}{3} - y = 6$$

$$-y = 6 - \frac{2}{3} = \frac{16}{3}$$

$$\Rightarrow y = -\frac{16}{3}$$

$$6. \quad (i) \quad 2^{x+2} = 2^x \cdot 2^2 = 4 \cdot 2^x$$

$$(ii) \quad 2^x + 2^x = 2 \cdot 2^x$$

$$\text{Hence, } 2^x + 2^x = 2^{x+2}(c - 2)$$

$$\Rightarrow 2 \cdot 2^x = 4 \cdot 2^x(c - 2)$$

$$\Rightarrow 2 = 4c - 8$$

$$\Rightarrow 4c = 2 + 8 = 10$$

$$\Rightarrow c = \frac{10}{4} = \frac{5}{2}$$

$$7. \quad 3^x = y \Rightarrow 3^{2x} = (3^x)^2 = y^2$$

$$\text{Solve } 3^{2x} - 12(3^x) + 27 = 0$$

$$\text{Let } y = 3^x \Rightarrow y^2 - 12y + 27 = 0$$

$$\Rightarrow (y - 3)(y - 9) = 0$$

$$\Rightarrow y = 3, y = 9$$

$$\text{Hence, } 3^x = 3, 3^x = 9$$

$$\Rightarrow 3^x = 3^1, 3^x = 3^2$$

$$\Rightarrow x = 1, x = 2$$

$$8. \quad \text{Solve } 2^{2x} - 3(2^x) - 4 = 0$$

$$\text{Let } y = 2^x \Rightarrow y^2 - 3y - 4 = 0$$

$$\Rightarrow (y - 4)(y + 1) = 0$$

$$\Rightarrow y = 4, y = -1$$

$$\text{Hence, } 2^x = 4 = 2^2, 2^x = -1 \text{ (Not valid)}$$

$$\Rightarrow x = 2$$

$$9. \quad (i) \quad \text{Solve } 2^{2x} - 9(2^x) + 8 = 0$$

$$\text{Let } y = 2^x \Rightarrow (y^2 - 9y + 8 = 0$$

$$\Rightarrow (y - 1)(y - 8) = 0$$

$$\Rightarrow y = 1, y = 8$$

$$\text{Hence, } 2^x = 1 = 2^0, 2^x = 8 = 2^3$$

$$\Rightarrow x = 0, x = 3$$

(ii) Solve $3^{2x} - 10(3^x) + 9 = 0$

Let $y = 3^x \Rightarrow y^2 - 10y + 9 = 0$

$$\Rightarrow (y - 1)(y - 9) = 0$$

$$\Rightarrow y = 1, y = 9$$

Hence, $3^x = 1 = 3^0, 3^x = 9 = 3^2$

$$\Rightarrow x = 0, x = 2$$

10. $y = 2^x \Rightarrow$ (i) $2^{2x} = (2^x)^2 = y^2$

(ii) $2^{2x+1} = (2^x)^2 \cdot 2^1 = 2y^2$

(iii) $2^{2x+3} = 2^x \cdot 2^3 = 8y$

Hence, solve $2^{2x+1} - 2^{x+3} - 2^x + 4 = 0$

Let $y = 2^x \Rightarrow 2y^2 - 8y - y + 4 = 0$

$$\Rightarrow 2y^2 - 9y + 4 = 0$$

$$\Rightarrow (2y - 1)(y - 4) = 0$$

$$\Rightarrow 2y = 1, y = 4$$

$$\Rightarrow y = \frac{1}{2}$$

Hence $2^x = \frac{1}{2} = 2^{-1}, 2^x = 4 = 2^2$

$$\Rightarrow x = -1, x = 2$$

11. Solve $3 \cdot 3^x + 3^{-x} = 4$

Let $y = 3^x \Rightarrow 3y + \frac{1}{y} = 4$

$$\Rightarrow 3y^2 + 1 = 4y$$

$$\Rightarrow 3y^2 - 4y + 1 = 0$$

$$\Rightarrow (3y - 1)(y - 1) = 0$$

$$\Rightarrow y = \frac{1}{3}, y = 1$$

Hence, $3^x = \frac{1}{3} = 3^{-1}, 3^x = 1 = 3^0$

$$\Rightarrow x = -1, x = 0$$

12. Solve $2(4^x) + 4^{-x} = 3$

Let $y = 4^x \Rightarrow 2y + \frac{1}{y} = 3$

$$\Rightarrow 2y^2 + 1 = 3y$$

$$\Rightarrow 2y^2 - 3y + 1 = 0$$

$$\Rightarrow (2y - 1)(y - 1) = 0$$

$$\Rightarrow y = \frac{1}{2}, y = 1$$

Hence, $4^x = \frac{1}{2}, 4^x = 1$

$$\Rightarrow (2^2)^x = 2^{-1}, (2^2)^x = 1 = 2^0$$

$$\Rightarrow 2^{2x} = 2^{-1} \quad 2^{2x} = 2^0$$

$$\Rightarrow 2x = -1 \Rightarrow 2x = 0$$

$$\Rightarrow x = -\frac{1}{2} \Rightarrow x = 0$$

13. Solve $3^x - 28 + 27(3^{-x}) = 0$

Let $y = 3^x \Rightarrow y - 28 + 27\left(\frac{1}{y}\right) = 0$

$$\Rightarrow y^2 - 28y + 27 = 0$$

$$\Rightarrow (y - 1)(y - 27) = 0$$

$$\Rightarrow y = 1, y = 27$$

Hence, $3^x = 1 = 3^0, 3^x = 27 = 3^3$

$$\Rightarrow x = 0, x = 3$$

14. Solve $2^{x+1} + 2(2^{-x}) - 5 = 0$

$$\begin{aligned}\text{Let } 2^x = y &\Rightarrow 2y + 2\left(\frac{1}{y}\right) - 5 = 0 \\ &\Rightarrow 2y^2 + 2 - 5y = 0 \\ &\Rightarrow 2y^2 - 5y + 2 = 0 \\ &\Rightarrow (2y - 1)(y - 2) = 0 \\ &\Rightarrow y = \frac{1}{2}, \quad y = 2\end{aligned}$$

$$\begin{aligned}\text{Hence, } 2^x = \frac{1}{2} = 2^{-1}, \quad 2^x = 2^1 \\ \Rightarrow x = -1, \quad x = 1\end{aligned}$$

15. Solve $3^x + 81(3^{-x}) - 30 = 0$

$$\begin{aligned}\text{Let } y = 3^x &\Rightarrow y + 81\left(\frac{1}{y}\right) - 30 = 0 \\ &\Rightarrow y^2 + 81 - 30y = 0 \\ &\Rightarrow y^2 - 30y + 81 = 0 \\ &\Rightarrow (y - 3)(y - 27) = 0 \\ &\Rightarrow y = 3, \quad y = 27\end{aligned}$$

$$\begin{aligned}\text{Hence, } 3^x = 3^1, \quad 3^x = 27 = 3^3 \\ \Rightarrow x = 1, \quad x = 3\end{aligned}$$

Exercise 12.8

- 1.** (i) B (ii) A (iii) D (iv) C

2. $A(n) = 1000 \times 2^{0.2n}$

$$\begin{aligned}\text{(i) } A(0) &= 1000 \times 2^{0.2(0)} \\ &= 1000 \times 2^0 \\ &= 1000 \times 1 = 1000 \text{ ha}\end{aligned}$$


$$\begin{aligned}\text{(ii) (a) } A(10) &= 1000 \times 2^{0.2(10)} \\ &= 1000 \times 2^2 \\ &= 1000 \times 4 = 4000 \text{ ha}\end{aligned}$$

$$\begin{aligned}\text{(b) } A(12) &= 1000 \times 2^{0.2(12)} \\ &= 1000 \times 2^{2.4} \\ &= 1000(5.278) = 5278 \text{ ha}\end{aligned}$$

(iii)

$n =$	0	2	4	6	8	10
$A(n) =$	1000	1320	1741	2297	3031	4000


- (iv) 5 weeks



3. (i) Decreasing
 (ii) Decreasing
 (iii) Increasing
 (iv) Decreasing
4. (i) $x = 0 \Rightarrow y = (0.6)2^0 = (0.6)(1) = 0.6$
 (ii) $x = 0 \Rightarrow y = 3(2^{-0}) = 3(1) = 3$
 (iii) $x = 0 \Rightarrow y = 8(2^0) = 8(1) = 8$
 (iv) $x = 0 \Rightarrow y = 6(4^{-0}) = 6(1) = 6$


5. (i)

$x =$	-2	-1	0	1	2	3	4
$2^x =$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8	16
$3^x =$	$\frac{1}{9}$	$\frac{1}{3}$	1	3	9	27	81




(ii)

$x =$	-2	-1	0	1	2	3	4
$2^{-x} =$	4	2	1	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$
$3^{-x} =$	9	3	1	$\frac{1}{3}$	$\frac{1}{9}$	$\frac{1}{27}$	$\frac{1}{81}$



(iii)

$x =$	-2	-1	0	1	2	3	4
$3.2^x =$	$\frac{3}{4}$	$1\frac{1}{2}$	3	6	12	24	48
$2.3^{-x} =$	18	6	2	$\frac{2}{3}$	$\frac{2}{9}$	$\frac{2}{27}$	$\frac{2}{81}$

(iv) $-2 \leq x < 0$ (v) $0 < x \leq 4$ (vi) $x = 0$ (vii) $0 < x \leq 4$ 6. $D = 18(0.72)^T$

(i) Decay: Graph is decreasing

(ii) (a) $T = 5^\circ\text{C} \Rightarrow D = 18(0.72)^5 = 3.48 = 3 \text{ days}$ (b) $T = 2^\circ\text{C} \Rightarrow D = 18(0.72)^2 = 9.33 = 9 \text{ days}$ (c) $T = 0^\circ\text{C} \Rightarrow D = 18(0.72)^0 = 18 \text{ days}$ (iii) $18(0.72)^T \geq 5$

$$\Rightarrow (0.72)^T \geq \frac{5}{18}$$

$$\Rightarrow \log(0.72)^T \geq \log \frac{5}{18}$$

$$\Rightarrow T(\log 0.72) \geq \log \frac{5}{18}$$

$$\Rightarrow T \geq \frac{\log \frac{5}{18}}{\log(0.72)}$$

$$\Rightarrow T \geq 3.899$$

$$\Rightarrow T = 3.9^\circ\text{C}$$

7. $P = 100(0.99988)^n$ (a) (i) $n = 200 \Rightarrow P = 100(0.99988)^{200} = 97.628 = 97.6\%$ (ii) $n = 500 \Rightarrow P = 100(0.99988)^{500} = 94.176 = 94.2\%$ (b) $n = 5000 \Rightarrow P = 100(0.99988)^{5000} = 54.879 = 54.9\%$ $n = 6000 \Rightarrow P = 100(0.99988)^{6000} = 48.673 = 48.7\%$ Hence, $100(0.99988)^n = 50$

$$\Rightarrow (0.99988)^n = 0.5$$

$$\Rightarrow \log(0.99988)^n = \log(0.5)$$

$$\Rightarrow n \log(0.99988) = \log(0.5)$$

$$\Rightarrow n = \frac{\log(0.5)}{\log(0.99988)} = 5775.88 = 5780 \text{ years}$$

$$\begin{aligned}
 \text{(c)} \quad & 100(0.99988)^n = 79 \\
 \Rightarrow & (0.99988)^n = 0.79 \\
 \Rightarrow & \log(0.99988)^n = \log(0.79) \\
 \Rightarrow & n \log(0.99988) = \log(0.79) \\
 \Rightarrow & n = \frac{\log(0.79)}{\log(0.99988)} = 1964.2 = 1964 \text{ years}
 \end{aligned}$$

8. $A(n) = 1000 \times 2^{0.2n}$

(i) $A(0) = 1000 \times 2^{0.2(0)}$

$$= 1000 \times 2^0$$

$$= 1000 \times 1 = 1000 \text{ ha}$$

(ii) (a) $A(5) = 1000 \times 2^{0.2(5)}$

$$= 1000 \times 2^1 = 2000 \text{ ha}$$


(b) $A(10) = 1000 \times 2^{0.2(10)}$

$$= 1000 \times 2^2 = 4000 \text{ ha}$$

(c) $A(12) = 1000 \times 2^{0.2(12)}$

$$= 1000 \times 2^{2.4}$$

$$= 1000(5.278) = 5278 \text{ ha}$$



9. $P(t) = 90 \times 3^{-0.25t} + 50$

(i)

$t =$	0	2	4	6	8	10
$90 \times 3^{-0.25t} + 50$	140	102	80	67	60	56

(ii) $P(0) = 90 \times 3^{-0.25(0)} + 50$

$$= 90 \times 3^0 + 50$$

$$= 90 + 50 = 140 \text{ beats/min}$$

(iii) (a) $90 \times 3^{-0.25t} + 50 = 70$

$$\Rightarrow 90 \times 3^{-0.25t} = 20$$


$$\Rightarrow 3^{-0.25t} = \frac{20}{90} = \frac{2}{9}$$

$$\Rightarrow \log 3^{-0.25t} = \log \frac{2}{9}$$

$$\Rightarrow -0.25t(\log 3) = \log \frac{2}{9}$$

$$\Rightarrow -0.25t = \frac{\log \frac{2}{9}}{\log 3} = -1.369$$

$$\Rightarrow t = \frac{-1.369}{-0.25} = 5.476 = 5.5 \text{ minutes}$$



(b) $90 \times 3^{-0.25t} + 50 = 55$

$$\Rightarrow 90 \times 3^{-0.25t} = 5$$

$$\Rightarrow 3^{-0.25t} = \frac{5}{90} = \frac{1}{18}$$

$$\Rightarrow \log 3^{-0.25t} = \log \frac{1}{18}$$

$$\Rightarrow -0.25t(\log 3) = \log \frac{1}{18}$$

$$\Rightarrow -0.25t = \frac{\log \frac{1}{18}}{\log 3} = -2.63$$

$$\Rightarrow t = \frac{-2.63}{-0.25} = 10.52 = 10.5 \text{ minutes}$$

(iv) 50 beats/min because $P(t) = 50$ as t gets larger using graph of $P(t)$.

10. $E \cong 10^{1.5M+4.8}$

$$M = 7 \Rightarrow E = 10^{1.5(7)+4.8} = 10^{15.3}$$

$$M = 5 \Rightarrow E = 10^{1.5(5)+4.8} = 10^{12.3}$$

$$\Rightarrow \text{No. of times} = \frac{10^{15.3}}{10^{12.3}} = 10^{15.3-12.3} = 10^3 = 1000$$

11. $P(t) = 40b^t$

(i) $t = 0 \Rightarrow P(0) = 40b^0 = 40$


(ii) $t = 1 \Rightarrow P(1) = 40b^1 = 48$

$$\Rightarrow b = \frac{48}{40} = 1.2 > 1$$

\Rightarrow No. of flies is increasing

(iii)

$t =$	0	1	2	3	4	5
$P(t) = 40(1.2)^t$	40	48	57.6	69.1	82.9	99.5



Exercise 12.9

1. (i) $\log_2 4 = 2$

(ii) $\log_3 81 = 4$

(iii) $\log_{10} 1000 = 3$

(iv) $\log_2 64 = 6$

2. (i) $\log_8 16 = x \Rightarrow 8^x = 16$

$$\Rightarrow (2^3)^x = 2^4$$

$$\Rightarrow 2^{3x} = 2^4$$

$$\Rightarrow 3x = 4 \Rightarrow x = \frac{4}{3}$$

(ii) $\log_9 27 = x \Rightarrow 9^x = 27$

$$\Rightarrow (3^2)^x = 3^3$$

$$\Rightarrow 3^{2x} = 3^3$$

$$\Rightarrow 2x = 3 \Rightarrow x = \frac{3}{2}$$

(iii) $\log_{16} 32 = x \Rightarrow 16^x = 32$

$$\Rightarrow (2^4)^x = 2^5$$

$$\Rightarrow 2^{4x} = 2^5$$

$$\Rightarrow 4x = 5 \Rightarrow x = \frac{5}{4}$$

$$\begin{aligned}
 \text{(iv) } \log_{\frac{1}{2}} 8 = x &\Rightarrow \left(\frac{1}{2}\right)^x = 8 \\
 &\Rightarrow (2^{-1})^x = 2^3 \\
 &\Rightarrow 2^{-x} = 2^3 \\
 &\Rightarrow -x = 3 \Rightarrow x = -3
 \end{aligned}$$

$$\begin{aligned}
 \text{(v) } \log_{\frac{1}{3}} 81 = x &\Rightarrow \left(\frac{1}{3}\right)^x = 81 \\
 &\Rightarrow (3^{-1})^x = 3^4 \\
 &\Rightarrow 3^{-x} = 3^4 \\
 &\Rightarrow -x = 4 \Rightarrow x = -4
 \end{aligned}$$

$$\begin{aligned}
 \text{3. (i) } \log_{\frac{1}{3}} 27 = x &\Rightarrow \left(\frac{1}{3}\right)^x = 27 \\
 &\Rightarrow (3^{-1})^x = 3^3 \\
 &\Rightarrow 3^{-x} = 3^3 \\
 &\Rightarrow -x = 3 \Rightarrow x = -3
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \log_{\sqrt{2}} 4 = x &\Rightarrow (\sqrt{2})^x = 4 \\
 &\Rightarrow (2^{\frac{1}{2}})^x = 2^2 \\
 &\Rightarrow 2^{\frac{1}{2}x} = 2^2 \\
 &\Rightarrow \frac{1}{2}x = 2 \\
 &\Rightarrow x = 4
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } \log_8 x = 2 \\
 &\Rightarrow 8^2 = x \\
 &\Rightarrow x = 64
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) } \log_{64} x = \frac{1}{2} \\
 &\Rightarrow 64^{\frac{1}{2}} = x \\
 &\Rightarrow x = 8
 \end{aligned}$$

$$\begin{aligned}
 \text{4. (i) } \log_2 x = -1 \\
 &\Rightarrow 2^{-1} = x \\
 &\Rightarrow x = \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \log_3 \sqrt{27} = x \\
 &\Rightarrow 3^x = \sqrt{27} = (3^3)^{\frac{1}{2}} = 3^{\frac{3}{2}} \\
 &\Rightarrow x = \frac{3}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } \log_x 2 = 2 &\Rightarrow x^2 = 2 \\
 &\Rightarrow x = 2^{\frac{1}{2}} = \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) } \log_2 (0.5) = x &\Rightarrow 2^x = \frac{1}{2} = 2^{-1} \\
 &\Rightarrow x = -1
 \end{aligned}$$

$$\begin{aligned}
 \text{5. (i) } \log_4 2 + \log_4 32 &= \log_4 (2) \cdot (32) \\
 &= \log_4 64 = x \\
 &\Rightarrow 4^x = 64 = 4^3 \\
 &\Rightarrow x = 3
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } \log_6 9 + \log_6 8 - \log_6 2 &= \log_6 \frac{(9)(8)}{(2)} \\
 &= \log_6 36 = x \\
 &\Rightarrow 6^x = 36 = 6^2 \\
 &\Rightarrow x = 2
 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad \log_6 4 + 2 \log_6 3 &= \log_6 4 + \log_6 3^2 \\ &= \log_6 4 + \log_6 9 = \log_6 36 = 2 \end{aligned}$$

$$\begin{aligned} \text{6. (i)} \quad \log_3 2 + 2 \log_3 3 - \log_3 18 \\ &= \log_3 2 + \log_3 3^2 - \log_3 18 \\ &= \log_3 2 + \log_3 9 - \log_3 18 \\ &= \log_3 \frac{(2)(9)}{18} = \log_3 1 = 0 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \log_8 72 - \log_8 \left(\frac{9}{8} \right) \\ &= \log_8 72 - (\log_8 9 - \log_8 8) \\ &= \log_8 72 - \log_8 9 + \log_8 8 \\ &= \log_8 \frac{(72)(8)}{(9)} = \log_8 64 = 2 \end{aligned}$$

$$\text{7. } \log_3 5 = a$$

$$\text{(i)} \quad \log_3 15 = \log_3 5 \cdot 3 = \log_3 5 + \log_3 3 = a + 1$$

$$\text{(ii)} \quad \log_3 \left(\frac{5}{3} \right) = \log_3 5 - \log_3 3 = a - 1$$

$$\begin{aligned} \text{(iii)} \quad \log_3 \left(8\frac{1}{3} \right) &= \log_3 \left(\frac{25}{3} \right) = \log_3 25 - \log_3 3 \\ &= \log_3 5^2 - 1 \\ &= 2 \log_3 5 - 1 = 2a - 1 \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad \log_3 \left(\frac{25}{27} \right) &= \log_3 25 - \log_3 27 \\ &= \log_3 (5)^2 - \log_3 (3)^3 \\ &= 2 \log_3 5 - 3 \log_3 3 = 2a - 3 \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad \log_3 75 &= \log_3 (25) \cdot (3) = \log_3 25 + \log_3 3 \\ &= \log_3 5^2 + 1 \\ &= 2 \log_3 5 + 1 \\ &= 2a + 1 \end{aligned}$$

$$\begin{aligned} \text{8. (i)} \quad 2^x &= 200 \\ \Rightarrow \log 2^x &= \log 200 \\ \Rightarrow x \log 2 &= \log 200 \\ \Rightarrow x &= \frac{\log 200}{\log 2} = 7.643 = 7.64 \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad 5^x &= 500 \\ \Rightarrow \log 5^x &= \log 500 \\ \Rightarrow x \log 5 &= \log 500 \\ \Rightarrow x &= \frac{\log 500}{\log 5} = 3.861 = 3.86 \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad 3^{x+1} &= 25 \\ \Rightarrow \log 3^{(x+1)} &= \log 25 \\ \Rightarrow (x+1) \log 3 &= \log 25 \\ \Rightarrow x+1 &= \frac{\log 25}{\log 3} = 2.929 \\ \Rightarrow x &= 1.929 = 1.93 \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad 5^{2x+3} &= 51 \\ \Rightarrow \log 5^{(2x+3)} &= \log 51 \\ \Rightarrow (2x+3) \log 5 &= \log 51 \\ \Rightarrow 2x+3 &= \frac{\log 51}{\log 5} = 2.4429 \\ \Rightarrow 2x &= -0.5571 \\ \Rightarrow x &= -0.2785 = -0.279 \end{aligned}$$

$$\begin{aligned}
 \text{9. (i)} \quad & y = 2^{x+1} + 3 \\
 & \Rightarrow 2^{x-1} = y - 3 \\
 & \Rightarrow \log(2^{x-1}) = \log(y - 3) \\
 & \Rightarrow (x - 1) \log 2 = \log(y - 3) \\
 & \Rightarrow x - 1 = \frac{\log(y - 3)}{\log 2} \\
 & \Rightarrow x = \frac{\log(y - 3)}{\log 2} + 1 \\
 \text{(ii)} \quad & y = 8 \Rightarrow x = \frac{\log(8 - 3)}{\log 2} + 1 = \frac{\log 5}{\log 2} + 1 = 2.321928 + 1 \\
 & = 3.321928 \\
 & = 3.3219
 \end{aligned}$$

$$\begin{aligned}
 \text{10.} \quad & \log_{10} x = 1 + a \Rightarrow 10^{1+a} = x \\
 & \log_{10} y = 1 - a \Rightarrow 10^{1-a} = y \\
 & \Rightarrow xy = 10^{1+a} \cdot 10^{1-a} = 10^{1+a+1-a} \\
 & = 10^2 = 100
 \end{aligned}$$

$$\begin{aligned}
 \text{11.} \quad & p = \log_a \frac{21}{4} = \log_a 21 - \log_a 4 \\
 & = \log_a 7 \cdot 3 - \log_a 4 \\
 & = \log_a 7 + \log_a 3 - \log_a 4 \\
 & q = \log_a \frac{7}{3} = \log_a 7 - \log_a 3 \\
 \Rightarrow \quad & p + q = \log_a 7 + \log_a 3 - \log_a 4 + \log_a 7 - \log_a 3 \\
 & = 2 \log_a 7 - \log_a 4 \\
 & 2r = 2 \log_a \frac{7}{2} = 2[\log_a 7 - \log_a 2] \\
 & = 2 \log_a 7 - 2 \log_a 2 \\
 & = 2 \log_a 7 - \log_a 2^2 = 2 \log_a 7 - \log_a 4 \\
 & \text{Hence, } p + q = 2r
 \end{aligned}$$

$$\begin{aligned}
 \text{12.} \quad & \text{If } \log_a x = 4 \text{ and } \log_a y = 5 \\
 \text{(i)} \quad & \log_a x^2 y = \log_a x^2 + \log_a y \\
 & = 2 \log_a x + \log_a y = 2(4) + 5 = 13 \\
 \text{(ii)} \quad & \log_a axy = \log_a a + \log_a x + \log_a y \\
 & = 1 + 4 + 5 = 10 \\
 \text{(iii)} \quad & \log_a \frac{\sqrt{x}}{y} = \log_a \sqrt{x} - \log_a y \\
 & = \log_a x^{\frac{1}{2}} - \log_a y \\
 & = \frac{1}{2} \log_a x - \log_a y \\
 & = \frac{1}{2}(4) - 5 = -3
 \end{aligned}$$

$$\text{13. Prove } \log_{25} x = \frac{1}{2} \log_5 x$$

$$\begin{aligned}
 \text{Proof } \log_{25} x &= \frac{\log_5 x}{\log_5 25} \\
 &= \frac{\log_5 x}{2} = \frac{1}{2} \log_5 x
 \end{aligned}$$

$$\begin{aligned}
 \text{14.} \quad & \text{(i) } \log_{10} 4 = 0.60205999 = 0.602 \\
 & \text{(ii) } \log_{10} 27 = 1.43136 = 1.43 \\
 & \text{(iii) } \log_{10} 356 = 2.5514 = 2.55 \\
 & \text{(iv) } \log_{10} 5600 = 3.748 = 3.75
 \end{aligned}$$

- (v) $\log_{10} 29\,000 = 4.462 = 4.46$
 (vi) $\log_{10} 35\,0000 = 5.544 = 5.54$
 (vii) $\log_{10} 38\,70\,000 = 6.5877 = 6.59$

15. Minimum $= 10^3 = 1000$
 Maximum $= 10^4 = 10\,000$

16. $\log_3 15 = \frac{\log_{10} 15}{\log_{10} 3} = \frac{1.17609}{0.47712} = 2.464977$
 $\log_2 5 = \frac{\log_{10} 5}{\log_{10} 2} = \frac{0.69897}{0.30103} = 2.3219$
 Hence, $\log_3 15 - \log_2 5 = 2.464977 - 2.3219$
 $= 0.143077 = 0.143$

17. (i) $\log_{27} 81 = \frac{\log_3 81}{\log_3 27} = \frac{4}{3}$
 (ii) $\log_{32} 8 = \frac{\log_2 8}{\log_2 32} = \frac{3}{5}$

18. Show: $\log_b a = \frac{1}{\log_a b}$
 Proof: $\log_b a = \frac{\log_a a}{\log_a b} = \frac{1}{\log_a b}$

19. $\frac{1}{\log_2 x} + \frac{1}{\log_3 x} + \frac{1}{\log_5 x} = \log_x 2 + \log_x 3 + \log_x 5$
 $= \log_x (2)(3)(5)$
 $= \log_x 30$
 $= \frac{1}{\log_{30} x}$

20. $\log_r p = \log_r 2 + 3 \log_r q$
 $\Rightarrow \log_r p = \log_r 2 + \log_r q^3$
 $\Rightarrow \log_r p = \log_r 2q^3$
 $\Rightarrow p = 2q^3$

21. $\log_3 a + \log_9 a = \frac{3}{4}$
 $\Rightarrow \log_3 a + \frac{\log_3 a}{\log_3 9} = \frac{3}{4}$
 $\Rightarrow \log_3 a + \frac{\log_3 a}{2} = \frac{3}{4}$
 $\Rightarrow 2 \log_3 a + \log_3 a = \frac{3}{2}$
 $\Rightarrow 3 \log_3 a = \frac{3}{2}$
 $\Rightarrow \log_3 a = \frac{1}{2}$
 $\Rightarrow 3^{\frac{1}{2}} = a \Rightarrow a = \sqrt{3}$

22. $3 \ln 41.5 - \ln 250 = 3(3.7256934) - 5.52146$
 $= 11.17708 - 5.52146$
 $= 5.6556$
 $= 5.66$

23. Solve $\log_2(x - 2) + \log_2 x = 3$

$$\begin{aligned} \Rightarrow \log_2(x - 2)(x) &= 3 \\ \Rightarrow x^2 - 2x &= 2^3 - 8 \\ \Rightarrow x^2 - 2x - 8 &= 0 \\ \Rightarrow (x - 4)(x + 2) &= 0 \\ \Rightarrow x = 4, \quad x = -2 & \text{ (Not valid)} \end{aligned}$$

24. $\log_{10}(x^2 + 6) - \log_{10}(x^2 - 1) = 1$

$$\begin{aligned} \Rightarrow \log_{10} \frac{x^2 + 6}{x^2 - 1} &= 1 \\ \Rightarrow \frac{x^2 + 6}{x^2 - 1} &= 10^1 = 10 \\ \Rightarrow 10x^2 - 10 &= x^2 + 6 \\ \Rightarrow 9x^2 - 16 &= 0 \\ \Rightarrow (3x + 4)(3x - 4) &= 0 \\ \Rightarrow x = -\frac{4}{3}, x = \frac{4}{3} &\Rightarrow x = \pm \frac{4}{3} \end{aligned}$$

25. $\log 2x - \log(x - 7) = \log 3$

$$\begin{aligned} \Rightarrow \log \frac{2x}{x - 7} &= \log 3 \\ \Rightarrow \frac{2x}{x - 7} &= 3 \\ \Rightarrow 3x - 21 &= 2x \\ \Rightarrow x &= 21 \end{aligned}$$

26. $\log(2x + 3) + \log(x - 2) = 2 \log x$

$$\begin{aligned} \Rightarrow \log(2x + 3)(x - 2) &= \log x^2 \\ \Rightarrow 2x^2 - x - 6 &= x^2 \\ \Rightarrow x^2 - x - 6 &= 0 \\ \Rightarrow (x - 3)(x + 2) &= 0 \\ \Rightarrow x = 3, x = -2 & \text{ (Not valid)} \end{aligned}$$

27. $\log_{10}(17 - 3x) + \log_{10} x = 1$

$$\begin{aligned} \Rightarrow \log_{10}(17 - 3x)(x) &= 1 \\ \Rightarrow 10^1 &= 17x - 3x^2 \\ \Rightarrow 3x^2 - 17x + 10 &= 0 \\ \Rightarrow (3x - 2)(x - 5) &= 0 \Rightarrow x = \frac{2}{3}, 5 \end{aligned}$$

28. $\log_{10}(x^2 - 4x - 11) = 0$

$$\begin{aligned} \Rightarrow 10^0 &= x^2 - 4x - 11 \\ \Rightarrow x^2 - 4x - 11 &= 1 \\ \Rightarrow x^2 - 4x - 12 &= 0 \\ \Rightarrow (x - 6)(x + 2) &= 0 \\ \Rightarrow x &= 6, -2 \end{aligned}$$

29. $2 \log_2 x = y$ and $\log_2(2x) = y + 4$

$$\begin{aligned} \Rightarrow \log_2 x^2 &= y \Rightarrow 2^{y+4} = 2x \\ \Rightarrow 2^y &= x^2 \Rightarrow 2^y \cdot 2^4 = 2x \\ &\Rightarrow x^2 \cdot 16 = 2x \\ &\Rightarrow 16x^2 - 2x = 0 \\ &\Rightarrow 8x^2 - x = 0 \\ &\Rightarrow x(8x - 1) = 0 \\ &\Rightarrow x = \frac{1}{8}, x = 0 \text{ (Not valid)} \end{aligned}$$

30. $\log_6 x + \log_6 y = 1$

$$\Rightarrow \log_6 x \cdot y = 1 \quad \Rightarrow 6 = xy$$

$$\Rightarrow x = \frac{6}{y}$$

Solve $\log_6 x + \log_6 y = 1 \cap 5x + y = 17$

$$\Rightarrow x = \frac{6}{y} \quad \Rightarrow 5\left(\frac{6}{y}\right) + y = 17$$

$$\Rightarrow \frac{30}{y} + y = 17$$

$$\Rightarrow 30 + y^2 = 17y$$

$$\Rightarrow y^2 - 17y + 30 = 0$$

$$(y - 2)(y - 15) = 0$$

$$\Rightarrow y = 2, y = 15$$

$$\Rightarrow x = \frac{6}{2} = 3, x = \frac{6}{15} = \frac{2}{5}$$

31. (i) Solve $4 \log_x 2 - \log_2 x - 3 = 0$

$$\Rightarrow 4 \log_x 2 - \frac{\log_x x}{\log_x 2} - 3 = 0$$

$$\Rightarrow 4 \log_x 2 - \frac{1}{\log_x 2} - 3 = 0$$

Let $y = \log_x 2 \Rightarrow 4y - \frac{1}{y} - 3 = 0$

$$\Rightarrow 4y^2 - 1 - 3y = 0$$

$$\Rightarrow 4y^2 - 3y - 1 = 0$$

$$\Rightarrow (4y + 1)(y - 1) = 0$$

$$\Rightarrow y = -\frac{1}{4}, y = 1$$

Hence, $\log_x 2 = -\frac{1}{4}, \log_x 2 = 1$

$$\Rightarrow x^{-\frac{1}{4}} = 2, x^1 = 2$$

$$\Rightarrow x = 2^{-4} = \frac{1}{16}, x = 2$$

(ii) Solve $2 \log_4 x + 1 = \log_x 4$

$$\Rightarrow 2 \log_4 x + 1 = \frac{\log_4 4}{\log_4 x}$$

$$\Rightarrow 2 \log_4 x + 1 = \frac{1}{\log_4 x}$$

Let $y = \log_4 x \Rightarrow 2y + 1 = \frac{1}{y}$

$$\Rightarrow 2y^2 + y = 1$$

$$\Rightarrow 2y^2 + y - 1 = 0$$

$$\Rightarrow (2y - 1)(y + 1) = 0$$

$$\Rightarrow y = \frac{1}{2}, y = -1$$

Hence, $\log_4 x = \frac{1}{2}, \log_4 x = -1$

$$\Rightarrow 4^{\frac{1}{2}} = x, 4^{-1} = x$$


$$\Rightarrow x = \sqrt{4} = 2, x = \frac{1}{4}$$

Exercise 12.10

1. Property 1 $\Rightarrow \log_a 1 = 0 \Rightarrow a^0 = 1$, true
 Property 2 $\Rightarrow \log_2 2 = 1 \Rightarrow 2^1 = 2$, true
 $\Rightarrow \log_e e = 1 \Rightarrow e^1 = e$, true
 $\Rightarrow \log_{10} 10 = 1 \Rightarrow 10^1 = 10$, true
 Property 5 $\Rightarrow \log_a 0 = y \Rightarrow a^y = 0 \Rightarrow$ No solution here

2.

$x =$	0	1	2
$y = 10^x =$	1	10	100
$y = \log_{10} x =$	undef.	0	0.3




3. (i)

$x =$	$\frac{1}{9}$	$\frac{1}{3}$	1	3	9
$y = \log_3 x =$	-2	-1	0	1	2

(ii) Graph $y = \log_3 x$

(iii) $\log_3 2.5 = 0.8$, using graph

$$\begin{aligned} \text{(iv) } \log_3 2.5 &= \frac{\log_{10} 2.5}{\log_{10} 3} \\ &= \frac{0.39794}{0.47712} = 0.834 \end{aligned}$$




4. (i)

$x =$	0	1	2
$y = 5^x =$	1	5	25

(ii)

$x =$	0	1	5	10	25
$y = \log_5 x =$	undef.	0	1	1.4	2

(iii) One graph is the inverse of the other.




5.

$x =$	0	1	2	4
$y = \log_2 x =$	undef.	0	1	2

$x =$	0	1	2	4
$2x =$	0	2	4	8
$y = \log_2 2x =$	undef.	1	2	3

$x =$	2	3	4	6
$x - 2 =$	0	1	2	4
$y = \log_2 (x - 2) =$	undef.	0	1	2




6.

$x =$	0	1	10	20
$y = \log_{10} x =$	undef.	0	1	1.3

$x =$	0	2	10	20
$\frac{x}{2} =$	0	1	5	10
$y = \log_{10} \frac{x}{2} =$	undef.	0	0.7	1

$x =$	-2	-1	8	18
$x + 2 =$	0	1	10	20
$y = \log_{10} (x + 2) =$	u.	0	1	1.3



7. (i)

$$3^{x+2} = y + 5$$

$$\Rightarrow \log 3^{x+2} = \log (y + 5)$$

$$\Rightarrow (x + 2) \log 3 = \log (y + 5)$$

$$\Rightarrow x + 2 = \frac{\log (y + 5)}{\log 3}$$

$$\Rightarrow x = \frac{\log (y + 5)}{\log 3} - 2 \quad \text{or} \quad \log_3 (y + 5) - 2$$

$$(ii) \quad x = \frac{\log (y + 5)}{\log 3} - 2$$


$$\text{when } y = 30 \Rightarrow x = \frac{\log (30 + 5)}{\log 3} - 2 = \frac{\log 35}{\log 3} - 2$$

$$= \frac{1.544068}{0.47712} - 2$$


$$= 3.2362 - 2$$

$$x = 1.2362 = 1.236$$


8. (i) Graph of $y = \log_{10} x + 2$




- (ii) Graph of $y = \log_{10}(x + 2)$




- (iii) Graph of $y = \log_{10} x - 2$




- (iv) Graph of $y = 2 \log_{10} x$



- (v) $y = -\log_{10} x$




9. Graphs of (i) $y = \log_{10}(2x)$
(ii) $y = \log_{10}\left(\frac{x}{2}\right)$




10. Graphs for:


(i) $y = 2^x + 1$




(ii) $y = 1 - 2^x$



(iii) $y = 2^{x+1}$



(iv) $y = \left(\frac{1}{2}\right) \cdot 2^x$



Exercise 12.11

$$1. \quad (i) \quad (a) \quad A = 5000 + 5000\left(\frac{0.6}{100}\right)^1 = 5000(1 + 0.006) \\ = 5000(1.006) = \text{€}5030$$

$$(b) \quad A = 5000 + 5000\left(\frac{0.6}{100}\right)^2 = 5000(1.006)^2 = \text{€}5060.18$$

$$(c) \quad A = 5000 + 5000\left(\frac{0.6}{100}\right)^3 = 5000(1.006)^3 = \text{€}5090.54$$

$$(ii) \quad 5000(1.006)^t$$

$$(iii) \quad 5000(1.006)^t = 10\,000$$

$$\Rightarrow (1.006)^t = \frac{10\,000}{5000} = 2$$

$$\Rightarrow \ln(1.006)^t = \ln 2$$

$$\Rightarrow t \ln(1.006) = \ln 2$$

$$\Rightarrow t = \frac{\ln 2}{\ln 1.006} = 115.87 = 116 \text{ months}$$

$$2. \quad y = Ae^{bt}$$

$$t = 0 \Rightarrow 100 = Ae^{b(0)} = Ae^0 = A \cdot 1$$

$$\Rightarrow A = 100$$

$$\text{Hence, } y = 100e^{bt}$$

$$t = 6 \Rightarrow 100e^{b(6)} = 450$$

$$\Rightarrow e^{6b} = \frac{450}{100} = 4.5$$

$$\Rightarrow \ln e^{6b} = \ln 4.5$$

$$\Rightarrow 6b \ln e = 6b(1) = \ln 4.5$$

$$\Rightarrow b = \frac{\ln 4.5}{6} = 0.2507 = 0.25$$

$$3. \quad T = 15 + 30 \times 10^{-0.02t}$$

$$(i) \quad t = 0 \Rightarrow T = 15 + 30 \times 10^{-0.02(0)} \\ = 15 + 30 \times 10^0 = 15 + 30(1) = 45^\circ\text{C}$$

$$(ii) \quad T = 35^\circ\text{C} \Rightarrow 15 + 30 \times 10^{-0.02t} = 35$$

$$\Rightarrow 30 \times 10^{-0.02t} = 35 - 15 = 20$$

$$\Rightarrow 10^{-0.02t} = \frac{20}{30} = \frac{2}{3}$$

$$\Rightarrow \log_{10} 10^{-0.02t} = \log_{10} \frac{2}{3}$$

$$\Rightarrow -0.02t(\log_{10} 10) = -0.02t(1) = \log \frac{2}{3}$$

$$\Rightarrow t = \frac{\log \frac{2}{3}}{-0.02} \\ = \frac{-0.17609}{-0.02} \\ = 8.8 \text{ minutes}$$

$$(iii) \quad \text{As } t \text{ gets larger, } 30 \times 10^{-0.02t} \text{ gets smaller.}$$

$$\text{Hence, smallest value for } 30 \times 10^{-0.02t} = 0.$$

$$\Rightarrow \text{Room temperature} = 15^\circ\text{C}$$

$$4. \quad L = 10 \log_{10} \left(\frac{I}{I_0} \right) = 10 \log_{10} \left(\frac{I}{1 \times 10^{-12}} \right)$$

$$(i) \quad L = 100 \Rightarrow 100 = 10[\log_{10} I - \log_{10}(1 \times 10^{-12})]$$

$$\Rightarrow 10 = \log_{10} I + 12$$

$$\Rightarrow \log_{10} I = -2$$

$$I = 10^{-2} = 0.01 \text{ Wm}^{-2}$$

$$L = 110 \Rightarrow 110 = 10[\log_{10} I - \log_{10}(1 \times 10^{-12})]$$

$$11 = \log_{10} I + 12$$

$$\Rightarrow \log_{10} I = -1 \Rightarrow I = 10^{-1} = 0.1 \text{ Wm}^{-2}$$

$$\Rightarrow \text{Range is between } 0.01 \text{ Wm}^{-2} \text{ and } 0.1 \text{ Wm}^{-2}$$

$$(ii) \quad I = 10 \text{ Wm}^{-2} \Rightarrow L = 10 \log_{10} \left(\frac{10}{1 \times 10^{-12}} \right)$$

$$= 10 \log_{10} 10^{13}$$

$$= 10 \cdot 13 \log_{10} 10$$

$$= 10 \cdot 13(1) = 130 \text{ dB}$$

$$5. \quad A = 10^M \quad \text{and} \quad E \cong 10^{1.5M+4.8}$$

$$= 10^{1.5M} \cdot 10^{4.8}$$

$$= (10^M)^{1.5} \cdot 10^{4.8}$$

$$= A^{1.5} \cdot 10^{4.8} = 10^a A^b$$

$$\text{Hence,} \quad a = 4.8 \text{ and } b = 1.5$$

$$6. \quad \text{Value} = \text{€}100 \quad \text{in} \quad 2000$$

$$(i) \quad t \text{ years later} \Rightarrow \text{Value} = 100 \left(1 + \frac{4.5}{100} \right)^t$$

$$= 100(1.045)^t$$

$$(ii) \quad 10 \text{ years later} \Rightarrow t = 10 \Rightarrow \text{Value} = 100(1.045)^{10}$$

$$= 155.2969 = \text{€}155.30$$

$$(iii) \quad 5 \text{ years earlier} \Rightarrow t = -5 \Rightarrow \text{Value} = 100(1.045)^{-5}$$

$$= 80.2451 = \text{€}80.25$$

$$7. \quad W = 0.6 \times 1.15^t$$

$$(i) \quad t = 0 \Rightarrow W = 0.6 \times 1.15^0$$

$$= 0.6 \times 1 = 0.6 \text{ kg}$$

$$(ii) \quad 1.15 = 1 + 0.15 = 1 + \frac{15}{100}$$

$$\Rightarrow \text{growth constant} = 15\%$$

$$(iii) \quad W = 2(0.6) = 1.2 \Rightarrow 0.6 \times 1.15^t = 1.2$$

$$\Rightarrow 1.15^t = \frac{1.2}{0.6} = 2$$

$$\Rightarrow \ln 1.15^t = \ln 2$$

$$\Rightarrow t \ln 1.15 = \ln 2$$

$$\Rightarrow t = \frac{\ln 2}{\ln 1.15} = 4.959 = 5 \text{ months}$$

$$8. \quad M = M_0 e^{-kt}$$

$$(i) \quad M = 10 \text{ when } t = 0 \Rightarrow M_0 e^{-k(0)} = 10$$

$$\Rightarrow M_0 e^0 = 10$$

$$\Rightarrow M_0(1) = M_0 = 10$$

Hence, $M = 10e^{-kt}$

$$\begin{aligned}
 M = 5 \text{ when } t = 140 &\Rightarrow 10 \cdot e^{-k(140)} = 5 \\
 &\Rightarrow e^{-140k} = \frac{5}{10} = 0.5 \\
 &\Rightarrow \ln e^{-140k} = \ln 0.5 \\
 &\Rightarrow -140k(\ln e) = \ln 0.5 \\
 &\Rightarrow -140k(1) = -0.693147 \\
 &\Rightarrow k = \frac{-0.693147}{-140} \\
 &= 0.00495
 \end{aligned}$$

(ii) $M = 10e^{-0.00495t}$

$$\begin{aligned}
 t = 70 &\Rightarrow M = 10e^{-0.00495(70)} \\
 &= 10e^{-0.3465} \\
 &= 10(0.7071) \\
 &= 7.071 = 7g
 \end{aligned}$$

(iii) $M = 2g \Rightarrow 10e^{-0.00495t} = 2$

$$\begin{aligned}
 &\Rightarrow e^{-0.00495t} = \frac{2}{10} = 0.2 \\
 &\Rightarrow \ln e^{-0.00495t} = \ln 0.2 \\
 &\Rightarrow -0.00495t(\ln e) = \ln 0.2 \\
 &\quad -0.00495t(1) = -1.609438 \\
 &\Rightarrow t = \frac{-1.609438}{-0.00495} = 325.1 = 325 \text{ days}
 \end{aligned}$$

Exercise 12.12(A)

1. Proof:

- (i) $n = 1? \Rightarrow 2(1) = 1(1 + 1) \Rightarrow 2 = 2$, true $n = 1$
- (ii) Assume true for $n = k$.
 $\Rightarrow 2 + 4 + 6 + 8 + \dots + 2k = k(k + 1)$
- (iii) Also true for $n = k + 1$?
 $\Rightarrow 2 + 4 + 6 + 8 + \dots + 2k + 2(k + 1) = k(k + 1) + 2(k + 1)$
 $\Rightarrow 2 + 4 + 6 + 8 + \dots + 2k + 2(k + 1) = k(k + 1)(k + 2)$
 $\Rightarrow 2 + 4 + 6 + 8 + \dots + 2k + 2(k + 1) = k(k + 1)[(k + 1) + 1]$
 \therefore It is true for $n = k + 1$.
- (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
- (v) Therefore, it is true for all positive integer values of n .

2. Proof:

- (i) $n = 1? \Rightarrow 1 = \frac{1}{2}[3(1) - 1] \Rightarrow 1 = \frac{1}{2}(2) = 1$, true $n = 1$,
- (ii) Assume true for $n = k$.
 $\Rightarrow 1 + 4 + 7 + 10 + \dots + (3k - 2) = \frac{k}{2}(3k - 1)$
- (iii) Also true for $n = k + 1$?
 $\Rightarrow 1 + 4 + 7 + 10 + \dots + (3k - 2) + (3k + 1) = \frac{k}{2}(3k - 1) + (3k + 1)$
 $\Rightarrow 1 + 4 + 7 + 10 + \dots + (3k - 2) + (3k + 1) = \frac{k(3k - 1) + 2(3k + 1)}{2}$

$$\Rightarrow 1 + 4 + 7 + 10 + \dots (3k - 2) + (3k + 1) = \frac{3k^2 - k + 6k + 2}{2}$$

$$\Rightarrow 1 + 4 + 7 + 10 + \dots (3k - 2) + (3k + 1) = \frac{3k^2 + 5k + 2}{2}$$

$$\Rightarrow 1 + 4 + 7 + 10 + \dots (3k - 2) + (3k + 1) = \frac{(k + 1)(3k + 2)}{2}$$

$$\Rightarrow 1 + 4 + 7 + 10 + \dots (3k - 2) + (3k + 1) = \frac{(k + 1)[3(k + 1) - 1]}{2}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

3. Proof:

(i) $n = 1?$ $\Rightarrow 1 \cdot 2 = \frac{1}{3}(1 + 1)(1 + 2) \Rightarrow 2 = \frac{1}{3}(2)(3) \Rightarrow 2 = 2$, true $n = 1$.

(ii) Assume true for $n = k$.

$$\Rightarrow 1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots k(k + 2) = \frac{k}{3}(k + 1)(k + 2)$$

(iii) Also true for $n = k + 1$?

$$\Rightarrow 1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots k(k + 1) + (k + 1)(k + 2) = \frac{k}{3}(k + 1)(k + 2) + (k + 1)(k + 2)$$

$$\Rightarrow 1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots k(k + 1) + (k + 1)(k + 2) = \frac{k(k + 1)(k + 2) + 3(k + 1)(k + 2)}{3}$$

$$\Rightarrow 1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots k(k + 1) + (k + 1)(k + 2) = \frac{(k + 1)(k + 2)(k + 3)}{3}$$

$$\Rightarrow 1 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 4 \cdot 5 + \dots k(k + 1) + (k + 1)(k + 2) = \frac{(k + 1)[(k + 1) + 1][(k + 1) + 2]}{3}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

4. Proof:

(i) $n = 1?$ $\Rightarrow \frac{1}{2 \cdot 3} = \frac{1}{2(1 + 2)} \Rightarrow \frac{1}{2 \cdot 3} = \frac{1}{2 \cdot 3} \Rightarrow$ true $n = 1$,

(ii) Assume true for $n = k$.

$$\Rightarrow \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} + \dots \frac{1}{(k + 1)(k + 2)} + \frac{k}{2(k + 2)}$$

(iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} + \dots \frac{1}{(k + 1)(k + 2)} + \frac{1}{(k + 2)(k + 3)} &= \frac{k}{2(k + 2)} + \frac{1}{(k + 2)(k + 3)} \\ &= \frac{k(k + 3) + 2(1)}{2(k + 2)(k + 3)} \\ &= \frac{k^2 + 3k + 2}{2(k + 2)(k + 3)} \\ &= \frac{(k + 1)(k + 2)}{2(k + 2)(k + 3)} \\ &= \frac{k + 1}{2(k + 2)} = \frac{k + 1}{2[(k + 1) + 2]} \end{aligned}$$

\therefore It is true for $n = k + 1$

- (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

5. Proof:

- (i) $n = 1? \Rightarrow \frac{1}{4 \cdot 5} = \frac{1}{4(1+4)} \Rightarrow \frac{1}{4 \cdot 5} = \frac{1}{4 \cdot 5}$, true $n = 1$
 (ii) Assume true for $n = k$.

$$\Rightarrow \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} + \frac{1}{6 \cdot 7} + \dots + \frac{1}{(k+3)(k+4)} = \frac{k}{4(k+4)}$$

 (iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow \frac{1}{4 \cdot 5} + \frac{1}{5 \cdot 6} + \frac{1}{6 \cdot 7} + \dots + \frac{1}{(k+3)(k+4)} + \frac{1}{(k+4)(k+5)} &= \frac{k}{4(k+4)} + \frac{1}{(k+4)(k+5)} \\ &= \frac{k(k+5) + 4(1)}{4(k+4)(k+5)} \\ &= \frac{k^2 + 5k + 4}{4(k+4)(k+5)} \\ &= \frac{(k+1)(k+4)}{4(k+4)(k+5)} \\ &= \frac{k+1}{4(k+5)} = \frac{k+1}{4[(k+1)+4]} \end{aligned}$$

\therefore It is true for $n = k + 1$.

- (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

6. Proof:

- (i) $n = 1? \Rightarrow 1^3 = \frac{(1)^2}{4}(1+1)^2 \Rightarrow 1 = \frac{1}{4} \Rightarrow 1 = 1$, true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 1^3 + 2^3 + 3^3 + \dots + k^3 = \frac{k^2}{4}(k+1)^2$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 1^3 + 2^3 + 3^3 + \dots + k^3 + (k+1)^3 &= \frac{k^2}{4}(k+1)^2 + (k+1)^3 \\ &= \frac{k^2(k+1)^2 + 4(k+1)^3}{4} \\ &= \frac{(k+1)^2[k^2 + 4(k+1)]}{4} \\ &= \frac{(k+1)^2(k+2)(k+2)}{4} \\ &= \frac{(k+1)^2(k+2)^2}{4} = \frac{(k+1)^2[(k+1)+1]^2}{4} \end{aligned}$$

\therefore It is true for $n = k + 1$.

- (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

7.
$$\sum_{n=1}^n n(n+2) = 1 \cdot 3 + 2 \cdot 4 + 3 \cdot 5 + \dots + n(n+2) = \frac{n(n+1)(2n+7)}{6}$$

Proof:

- (i) $n = 1? \Rightarrow 1 \cdot 3 = \frac{1(1+1)(2+7)}{6} \Rightarrow 3 = \frac{2 \cdot 9}{6} \Rightarrow 3 = 3$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow 1 \cdot 3 + 2 \cdot 4 + 3 \cdot 5 + \dots k(k+2) = \frac{k(k+1)(2k+7)}{6}$

(iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 1 \cdot 3 + 2 \cdot 4 + 3 \cdot 5 + \dots k(k+2) + (k+1)(k+3) &= \frac{k(k+1)(2k+7)}{6} + (k+1)(k+3) \\ &= \frac{k(k+1)(2k+7) + 6(k+1)(k+3)}{6} \\ &= \frac{(k+1)[2k^2 + 7k + 6k + 18]}{6} \\ &= \frac{(k+1)[2k^2 + 13k + 18]}{6} \\ &= \frac{(k+1)(k+2)(2k+9)}{6} = \frac{(k+1)[(k+1)+1][2(k+1)+7]}{6} \end{aligned}$$

\therefore It is true for $n = k + 1$

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

8. Proof:

(i) $n = 1? \Rightarrow x = \frac{x(x^1 - 1)}{x - 1} \Rightarrow x = x$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow x + x^2 + x^3 + x^4 + \dots + x^k = \frac{x(x^k - 1)}{x - 1}$

(iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow x + x^2 + x^3 + \dots + x^k + x^{k+1} &= \frac{x(x^k - 1)}{x - 1} + x^{k+1} \\ &= \frac{x(x^k - 1) + x^{k+1}(x - 1)}{x - 1} \\ &= \frac{x \cdot x^k - x + x \cdot x^{k+1} - x^{k+1}}{x - 1} \\ &= \frac{x^{k+1} + x \cdot x^{k+1} - x - x^{k+1}}{x - 1} \\ &= \frac{x(x^{k+1} - 1)}{x - 1} \end{aligned}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

Exercise 12.12(B)

1. Proof:

(i) True for $n = 1? \Rightarrow 5$ is a factor of $6^1 - 1 = 5$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow 5$ is a factor of $6^k - 1$, $k \in \mathbb{N}$.

(iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 6^{k+1} - 1 &= 6^k \cdot 6^1 - 1 \\ &= 6^k(5 + 1) - 1 \\ &= 5 \cdot 6^k + (6^k - 1) \end{aligned}$$

Since $5 \cdot 6^k$ is divisible by 5 and $(6^k - 1)$ is assumed divisible by 5,

$\therefore 5 \cdot 6^k + (6^k - 1)$ is divisible by 5.

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

2. Proof:

- (i) True for $n = 1$? $\Rightarrow 4$ is a factor of $5^1 - 1 = 4$, true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 4$ is a factor of $5^k - 1$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 5^{k+1} - 1 &= 5^k \cdot 5^1 - 1 \\ &= 5^k(4 + 1) - 1 \\ &= 4 \cdot 5^k + (5^k - 1)\end{aligned}$$
 Since $4 \cdot 5^k$ is divisible by 4 and $(5^k - 1)$ is assumed divisible by 4,
 $\therefore 4 \cdot 5^k + (5^k - 1)$ is divisible by 4.
 \therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

3. Proof:

- (i) True for $n = 1$? $\Rightarrow 4$ is a factor of $9^1 - 5^1 = 4$, true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 4$ is a factor of $9^k - 5^k$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 9^{k+1} - 5^{k+1} &= 9^k \cdot 9^1 - 5^k \cdot 5^1 \\ &= 9^k(8 + 1) - 5^k(4 + 1) \\ &= 8 \cdot 9^k + 9^k - 4 \cdot 5^k - 5^k \\ &= 8 \cdot 9^k - 4 \cdot 5^k + (9^k - 5^k)\end{aligned}$$
 Since $8 \cdot 9^k - 4 \cdot 5^k$ is divisible by 4 and $9^k - 5^k$ is assumed divisible by 4,
 $\therefore 8 \cdot 9^k - 4 \cdot 5^k + (9^k - 5^k)$ is divisible by 4.
 \therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

4. Proof:

- (i) True for $n = 1$? $\Rightarrow 8$ is a factor of $3^{2(1)} - 1 = 9 - 1 = 8$, true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 8$ is a factor of $3^{2k} - 1$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 3^{2(k+1)} - 1 &= 3^{2k+2} - 1 \\ &= 3^{2k} \cdot 3^2 - 1 \\ &= 3^{2k} \cdot 9 - 1 \\ &= 3^{2k}(8 + 1) - 1 \\ &= 8 \cdot 3^{2k} + (3^{2k} - 1)\end{aligned}$$
 Since $8 \cdot 3^{2k}$ is divisible by 8 and $(3^{2k} - 1)$ is assumed divisible by 8,
 $\therefore 8 \cdot 3^{2k} + (3^{2k} - 1)$ is divisible by 8.
 (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .

5. Proof:

- (i) True for $n = 1$? $\Rightarrow 5$ is a factor of $7^1 - 2^1 = 5$, true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 5$ is a factor of $7^k - 2^k$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 7^{k+1} - 2^{k+1} &= 7^k \cdot 7^1 - 2^k \cdot 2^1 \\ &= 7^k(5 + 2) - 2 \cdot 2^k \\ &= 5 \cdot 7^k + 2 \cdot 7^k - 2 \cdot 2^k \\ &= 5 \cdot 7^k + 2(7^k - 2^k)\end{aligned}$$

Since $5 \cdot 7^k$ is divisible by 5 and $(7^k - 2^k)$ is assumed divisible by 5,
 $\therefore 5 \cdot 7^k + 2(7^k - 2^k)$ is divisible by 5.

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

6. Proof:

(i) True for $n = 1$? $\Rightarrow 8$ is a factor of $7^{2(1)+1} + 1 = 7^3 + 1 = 344 = 8 \cdot 43 \Rightarrow$ true $n = 1$.

(ii) Assume true for $n = k \Rightarrow 7^{2k+1} + 1$ is divisible by 8.

(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 7^{2(k+1)+1} + 1 &= 7^{2k+2+1} + 1 \\ &= 7^{2k+1} \cdot 7^2 + 1 \\ &= 7^{2k+1} \cdot 49 + 1 \\ &= 7^{2k+1}(48 + 1) + 1 \\ &= 48 \cdot 7^{2k+1} + (7^{2k+1} + 1)\end{aligned}$$

Since $48 \cdot 7^{2k+1}$ is divisible by 8 and $(7^{2k+1} + 1)$ is assumed divisible by 8,

$\therefore 48 \cdot 7^{2k+1} + (7^{2k+1} + 1)$ is divisible by 8.

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

7. Proof:

(i) True for $n = 1$? $\Rightarrow 7$ is a factor of $2^{3(1)-1} + 3 = 2^2 + 3 = 7$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow 7$ is a factor of $2^{3k-1} + 3$, $k \in N$.

(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 2^{3(k+1)-1} + 3 &= 2^{3k+3-1} + 3 \\ &= 2^{3k-1} \cdot 2^3 + 3 \\ &= 2^{3k-1} \cdot 8 + 3 \\ &= 2^{3k-1}(7 + 1) + 3 \\ &= 7 \cdot 2^{3k-1} + (2^{3k-1} + 3)\end{aligned}$$

Since $7 \cdot 2^{3k-1}$ is divisible by 7 and $2^{3k-1} + 3$ is assumed divisible by 7,

$\therefore 7 \cdot 2^{3k-1} + (2^{3k-1} + 3)$ is divisible by 7.

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it must now be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

8. Proof:

(i) True for $n = 1$? $\Rightarrow 4$ is a factor of $5^1 - 4(1) + 3 = 4$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow 4$ is a factor of $5^k - 4k + 3$, $k \in N$.

(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 5^{k+1} - 4(k + 1) + 3 &= 5^k \cdot 5^1 - 4k - 4 + 3 \\ &= 5^k(4 + 1) - 4k - 4 + 3 \\ &= 4 \cdot 5^k + 5^k - 4k - 4 + 3 \\ &= 4 \cdot 5^k - 4 + (5^k - 4k + 3)\end{aligned}$$

Since $4 \cdot 5^k - 4$ is divisible by 4 and $(5^k - 4k + 3)$ is assumed divisible by 4,

$\therefore 4 \cdot 5^k - 4 + (5^k - 4k + 3)$ is divisible by 4.

\therefore It is true for $n = k + 1$.

- (iv) But since it is true for $n = 1$, it must now be true for $n = 1 + 1 = 2$.
And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
(v) Therefore, it is true for all positive integer values of n .

9. Proof:

- (i) True for $n = 1$? $\Rightarrow 6$ is a factor of $7^1 + 4^1 + 1 = 12 \Rightarrow$ true $n = 1$.
(ii) Assume true for $n = k \Rightarrow 6$ is a factor of $7^k + 4^k + 1$, $k \in N$.
(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow 7^{k+1} + 4^{k+1} + 1 &= 7^k \cdot 7^1 + 4^k \cdot 4^1 + 1 \\ &= 7^k(6 + 1) + 4^k(3 + 1) + 1 \\ &= 6 \cdot 7^k + 7^k + 3 \cdot 4^k + 4^k + 1 \\ &= 6 \cdot 7^k + 3 \cdot 4^k + (7^k + 4^k + 1)\end{aligned}$$
 4^k is an even number $\Rightarrow 3 \cdot 4^k$ is divisible by 6 and
 $7^k + 4^k + 1$ is assumed divisible by 6;
 $\therefore 6 \cdot 7^k + 3 \cdot 4^k + (7^k + 4^k + 1)$ is divisible by 6.
 \therefore It is true for $n = k + 1$.
(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
(v) Therefore, it is true for all positive integer values of n .

10. Proof:

- (i) True for $n = 1$? $\Rightarrow 1(1+1)[2(1) + 1] = (1)(2)(3) = 6 \Rightarrow$ true $n = 1$.
(ii) Assume true for $n = k \Rightarrow 3$ is a factor of $k(k+1)(2k+1) = 2k^3 + 3k^2 + k$, $k \in N$.
(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow (k+1)[(k+1) + 1][2(k+1) + 1] &= (k+1)(k+2)(2k+3) \\ &= (k+1)(2k^2 + 7k + 6) \\ &= 2k^3 + 9k^2 + 13k + 6 \\ &= 2k^3 + 6k^2 + 3k^2 + 12k + k + 6 \\ &= 6k^2 + 12k + 6 + (2k^3 + 3k^2 + k)\end{aligned}$$
Since $6k^2 + 12k + 6$ is divisible by 3, and $(2k^3 + 3k^2 + k)$ is assumed divisible by 3,
 $\therefore 6k^2 + 12k + 6 + (2k^3 + 3k^2 + k)$ is divisible by 3.
(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
(v) Therefore, it is true for all positive integer values of n .

11. Prove $n^3 - n$ is divisible by 3 for $n \in N$.

Proof:

- (i) True for $n = 1$? $\Rightarrow 3$ is a factor of $1^3 - 1 = 1 - 1 = 0 \Rightarrow$ true $n = 1$.
(ii) Assume true for $n = k \Rightarrow k^3 - k$ is divisible by 3.
(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow (k+1)^3 - (k+1) &= k^3 + 3k^2 + 3k + 1 - k - 1 \\ &= 3k^2 + 3k + (k^3 - k)\end{aligned}$$
Since $3k^2 + 3k$ is divisible by 3, and $(k^3 - k)$ is assumed divisible by 3,
 $\therefore 3k^2 + 3k + (k^3 - k)$ is divisible by 3.
 \therefore It is true for $n = k + 1$.
(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
(v) Therefore, it is true for all positive integer values of n .

12. Prove $13^n - 6^{n-2}$ is divisible by 7 for $n \in N$.

Proof:

- (i) True for $n = 2$? $\Rightarrow 7$ is a factor of $13^2 - 6^{2-2} = 169 - 1 = 168 = 7 \cdot 24 \Rightarrow$ true $n = 2$.
 (ii) Assume true for $n = k \Rightarrow 13^k - 6^{k-2}$ is divisible by 7.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 13^{k+1} - 6^{(k+1)-2} &= 13^k \cdot 13^1 - 6^{k-2} \cdot 6^1 \\ &= 13^k(7 + 6) - 6^{k-2} \cdot 6 \\ &= 7 \cdot 13^k + 6 \cdot 13^k - 6 \cdot 6^{k-2} \\ &= 7 \cdot 13^k + 6(13^k - 6^{k-2}) \end{aligned}$$

Since $7 \cdot 13^k$ is divisible by 7, and $(13^k - 6^{k-2})$ is assumed divisible by 7,
 $\therefore 7 \cdot 13^k + 6(13^k - 6^{k-2})$ is divisible by 7.
 \therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n excluding 1.

Exercise 12.12(C)**1.** Proof:

- (i) True for $n = 3$? $\Rightarrow 2^3 > 2(3) + 1 \Rightarrow 8 > 7 \Rightarrow$ true $n = 3$.
 (ii) Assume true for $n = k \Rightarrow 2^k > 2k + 1, k \in N, k \geq 3$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 2^{k+1} &= 2 \cdot 2^k \\ &> 2(2k + 1) && \text{since } 2^k > 2k + 1 \text{ (assumed)} \\ &= 4k + 2 \\ &= 2k + 2k + 2 \\ &> 2k + 3 && \text{since } 2k + 2 > 3, \text{ for } k \geq 3 \\ &= 2(k + 1) + 1 \end{aligned}$$

\therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 3$, it now must be true for $n = 3 + 1 = 4$.
 And if it is true for $n = 4$, it is true for $n = 4 + 1 = 5 \dots$ etc.
 (v) Therefore, it is true for all values of $n, n \geq 3, n \in N$.

2. Proof:

- (i) True for $n = 2$? $\Rightarrow 3^2 > 2^2 \Rightarrow 9 > 4 \Rightarrow$ true $n = 2$.
 (ii) Assume true for $n = k \Rightarrow 3^k > k^2, k \in N, \text{ and } k \geq 2$.
 (iii) Also true for $n = k + 1$?

$$\begin{aligned} \Rightarrow 3^{k+1} &= 3^1 \cdot 3^k \\ &> 3 \cdot k^2 && \text{since } 3^k > k^2 \text{ (assumed)} \\ &= k^2 + k^2 + k^2 \\ &> k^2 + 2k + 1 && \text{since } 2k^2 > 2k + 1, \text{ for } k \geq 2 \\ &= (k + 1)(k + 1) \\ &= (k + 1)^2 \end{aligned}$$

\therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 2$, it now must be true for $n = 2 + 1 = 3$.
 And if it is true for $n = 3$, it is true for $n = 3 + 1 = 4 \dots$ etc.
 (v) Therefore, it is true for all values of $n, n \geq 2, n \in N$.

3. Proof:(i) True for $n = 2$? $\Rightarrow 3^2 > 2(2) + 2 \Rightarrow 9 > 6 \Rightarrow \text{true } n = 2.$ (ii) Assume true for $n = k \Rightarrow 3^k > 2k + 2, k \in N, k \geq 2.$ (iii) Also true for $n = k + 1$?

$$\begin{aligned}
 \Rightarrow 3^{k+1} &= 3 \cdot 3^k \\
 &> 3(2k + 2) && \text{since } 3^k > 2k + 2 \text{ (assumed)} \\
 &= 6k + 6 \\
 &= 2k + 4k + 2 + 4 \\
 &= 2k + 4 + 4k + 2 \\
 &> 2k + 4 && \text{since } 4k + 2 > 0, \text{ for } k \geq 2 \\
 &= 2(k + 1) + 2
 \end{aligned}$$

 \therefore It is true for $n = k + 1.$ (iv) But since it is true for $n = 2$, it now must be true for $n = 2 + 1 = 3.$ And if it is true for $n = 3$, it is true for $n = 3 + 1 = 4 \dots$ etc.(v) Therefore, it is true for all values of $n, n \geq 2, n \in N.$ **4. Proof:**(i) True for $n = 3$? $\Rightarrow 3! > 2^{3-1} \Rightarrow 6 > 4 \Rightarrow \text{true } n = 3.$ (ii) Assume true for $n = k \Rightarrow k! > 2^{k-1}, k \in N, k \geq 3.$ (iii) Also true for $n = k + 1$?

$$\begin{aligned}
 \Rightarrow (k + 1)! &= (k + 1)k! \\
 &> (k + 1) \cdot 2^{k-1} && \text{since } k! > 2^{k-1} \text{ (assumed)} \\
 &> (1 + 1) \cdot 2^{k-1} && \text{since } k + 1 > 2, \text{ for } k \geq 3 \\
 &= 2 \cdot 2^{k-1} \\
 &= 2^k \\
 &= 2^{(k+1)-1}
 \end{aligned}$$

 \therefore It is true for $n = k + 1.$ (iv) But since it is true for $n = 3$, it now must be true for $n = 3 + 1 = 4.$ And if it is true for $n = 4$, it is true for $n = 4 + 1 = 5 \dots$ etc.(v) Therefore, it is true for all values of $n, n \geq 3, n \in N.$ **5. Proof:**(i) True for $n = 2$? $\Rightarrow (2 + 1)! > 2^2 \Rightarrow 6 > 4 \Rightarrow \text{true } n = 2.$ (ii) Assume true for $n = k \Rightarrow (k + 1)! > 2^k.$ (iii) Also true for $n = k + 1$?

$$\begin{aligned}
 \Rightarrow (k + 1 + 1)! &= (k + 2)! \\
 &= (k + 2)(k + 1)! \\
 &> (k + 2) \cdot 2^k && \text{since } (k + 1)! > 2^k \text{ (assumed)} \\
 &> 2 \cdot 2^k && \text{since } (k + 2) > 2, \text{ for } k \geq 2 \\
 &= 2 \cdot 2^{k+1}
 \end{aligned}$$

 \therefore It is true for $n = k + 1.$ (iv) But since it is true for $n = 2$, it now must be true for $n = 2 + 1 = 3.$ And if it is true for $n = 3$, it is true for $n = 3 + 1 = 4 \dots$ etc.(v) Therefore, it is true for all values of $n, n \geq 2, n \in N.$ **6. Proof:**(i) True for $n = 1$? $\Rightarrow (1 + 2x)^1 = 1 + 2(1)x \Rightarrow 1 + 2x = 1 + 2x \Rightarrow \text{true } n = 1.$ (ii) Assume true for $n = k \Rightarrow (1 + 2x)^k \geq 1 + 2kx$ for $x > 0, k \in N.$

(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow (1 + 2x)^{k+1} &= (1 + 2x)(1 + 2x)^k \\ &\geq (1 + 2x)(1 + 2kx) && \text{since } (1 + 2x)^k \geq 1 + 2kx \text{ (assumed)} \\ &= 1 + 2kx + 2x + 4kx^2 \\ &\geq 1 + 2kx + 2x && \text{since } 4kx^2 \geq 0, \text{ for } x > 0, k \geq 1 \\ &= 1 + 2(k+1)x\end{aligned}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all values of n , $n \geq 1$, $n \in N$.

7. Prove $(1 + ax)^n \geq 1 + anx$ for $a > 0$, $x > 0$, $n \in N$.

Proof:

(i) True for $n = 1$? $\Rightarrow (1 + ax)^1 \geq 1 + a(1)x \Rightarrow 1 + ax \geq 1 + ax$, true $n = 1$.

(ii) Assume true for $n = k \Rightarrow (1 + ax)^k \geq 1 + akx$.

(iii) Also true for $n = k + 1$?

$$\begin{aligned}\Rightarrow (1 + ax)^{k+1} &= (1 + ax)(1 + ax)^k \\ &\geq (1 + ax)(1 + akx) && \text{since } (1 + ax)^k \geq 1 + akx \text{ (assumed)} \\ &= 1 + akx + ax + a^2kx^2 \\ &\geq 1 + akx + ax && \text{since } a^2kx^2 \geq 0, \text{ for } a > 0, x > 0 \\ &= 1 + a(k+1)x\end{aligned}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all values of n , $n \geq 1$, $n \in N$.

Revision Exercise 12 (Core)

1. $-1 \leq \frac{2x+4}{3} \leq 2, x \in R$

$$\Rightarrow -3 \leq 2x + 4 \leq 6$$

$$\Rightarrow -7 \leq 2x \leq 2$$

$$\Rightarrow -3.5 \leq x \leq 1$$

2. (a) (i) $10^{3.5} = 3162.278 = 3162$

(ii) $\log_{10} 4.5 = 0.6532 = 0.65$

(iii) $t = 0.04 \Rightarrow 10^{3t} = 10^{3(0.04)} = 10^{0.12} = 1.318 = 1.32$

(iv) $n = 100 \Rightarrow \log 5n = \log 5(100) = \log 500 = 2.69897 = 2.7$

(b) (i) $e^{3.4} = 29.964 = 30$

(ii) $\ln 589 = 6.378 = 6.38$

(iii) $t = 40 \Rightarrow e^{-0.02t-4} = e^{-0.02(40)-4} = e^{-0.8-4} = e^{-4.8}$
 $= 0.0082297 = 0.00823$

(iv) $k = 3.7 \Rightarrow \ln\left(\frac{10}{k}\right) = \ln\left(\frac{10}{3.7}\right) = 0.994 = 0.99$

3. (i) $f(x) = 3 \times 4^x$

Point $(a, 6) \Rightarrow f(a) = 3 \times 4^a = 6$

$$\Rightarrow 4^a = 2$$

$$\Rightarrow (2^2)^a = 2$$

$$\Rightarrow 2^{2a} = 2^1$$

$$\Rightarrow 2a = 1 \Rightarrow a = \frac{1}{2}$$

$$\begin{aligned}
 \text{(ii) Point } \left(-\frac{1}{2}, b\right) &\Rightarrow f\left(-\frac{1}{2}\right) = 3 \times 4^{-\frac{1}{2}} = b \\
 &\Rightarrow 3 \times \frac{1}{2} = b \\
 &\Rightarrow b = \frac{3}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{4. } x - 8 = -3 \quad \text{or} \quad x - 8 = 3 \\
 \Rightarrow x = 5 \quad \text{or} \quad x = 11
 \end{aligned}$$

$$\begin{aligned}
 \text{5. (i) } 5^{2n} \times 25^{2n-1} &= 625 \\
 \Rightarrow 5^{2n} \times (5^2)^{2n-1} &= 5^4 \\
 \Rightarrow 5^{2n} \times 5^{4n-2} &= 5^4 \\
 \Rightarrow 5^{2n+4n-2} &= 5^{6n-2} = 5^4 \\
 &\Rightarrow 6n - 2 = 4 \\
 &\Rightarrow 6n = 6 \Rightarrow n = 1
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } 27^{n-2} &= 9^{3n+2} \\
 \Rightarrow (3^3)^{n-2} &= (3^2)^{3n+2} \\
 \Rightarrow 3^{3n-6} &= 3^{6n+4} \\
 \Rightarrow 3n - 6 &= 6n + 4 \\
 \Rightarrow -3n &= 10 \Rightarrow n = -\frac{10}{3}
 \end{aligned}$$

$$\text{6. } y = a2^x + b$$

$$\begin{aligned}
 \text{(i) Point } (0, 2.5) &\Rightarrow 2.5 = a \cdot 2^0 + b = a \cdot 1 + b \Rightarrow a + b = 2.5 \\
 \text{Point } (2, 4) &\Rightarrow 4 = a \cdot 2^2 + b = 4a + b \Rightarrow 4a + b = 4
 \end{aligned}$$

$$\begin{aligned}
 \text{(ii) } 4a + b &= 4 \\
 a + b &= 2.5 \\
 \Rightarrow \frac{a + b}{3a} &= \frac{2.5}{1.5} \Rightarrow a = \frac{1.5}{3} = 0.5 \\
 \text{and } 0.5 + b &= 2.5 \Rightarrow b = 2.5 - 0.5 = 2
 \end{aligned}$$

$$\begin{aligned}
 \text{7. (i) Curve } C = \ln(x) &\Rightarrow \text{Point } (1, 0) \Rightarrow \ln 1 = 0, \text{ true} \\
 \text{(ii) Curve } B = \ln(x + 1) &\Rightarrow \text{Point } (0, 0) \Rightarrow \ln(0 + 1) = \ln 1 = 0, \text{ true} \\
 \text{(iii) Curve } A = \ln(x) + 1 &\Rightarrow \text{Point } (1, 1) \Rightarrow \ln 1 + 1 = 0 + 1 = 1, \text{ true}
 \end{aligned}$$

$$\begin{aligned}
 \text{8. Solve } \ln(x - 1) + \ln(x + 2) &= \ln(6x - 8) \\
 \Rightarrow \ln(x - 1)(x + 2) &= \ln(6x - 8) \\
 \Rightarrow (x - 1)(x + 2) &= 6x - 8 \\
 \Rightarrow x^2 + x - 2 &= 6x - 8 \\
 \Rightarrow x^2 - 5x + 6 &= 0 \\
 \Rightarrow (x - 2)(x - 3) &= 0 \\
 \Rightarrow x = 2, x = 3
 \end{aligned}$$

$$\text{9. } y = Ae^{bt}$$

$$\begin{aligned}
 y = 6 \text{ when } t = 1 &\Rightarrow 6 = Ae^{b(1)} \Rightarrow Ae^b = 6 \Rightarrow e^b = \frac{6}{A} \\
 y = 8 \text{ when } t = 2 &\Rightarrow 8 = Ae^{b(2)} \Rightarrow Ae^{2b} = 8 \\
 &\Rightarrow A(e^b)^2 = 8 \\
 &\Rightarrow A\left(\frac{6}{A}\right)^2 = 8 \\
 &\Rightarrow A \cdot \frac{36}{A^2} = \frac{36}{A} = 8 \\
 &\Rightarrow 8A = 36 \Rightarrow A = \frac{9}{2}
 \end{aligned}$$

$$e^b = \frac{6}{\frac{9}{2}} = \frac{4}{3}$$

$$\Rightarrow \ln e^b = \ln \frac{4}{3}$$

$$\Rightarrow b(\ln e) = b = \ln \frac{4}{3}$$

10. $y = a \log_2(x - b)$

$$(5, 2) \Rightarrow 2 = a \log_2(5 - b)$$

$$\Rightarrow 2 = \log_2(5 - b)^a \Rightarrow (5 - b)^a = 2^2 = 4$$

$$\Rightarrow 5 - b = 4^{\frac{1}{a}}$$

$$\Rightarrow b = 5 - 4^{\frac{1}{a}}$$

$$(7, 4) \Rightarrow 4 = a \log_2(7 - b)$$

$$\Rightarrow 4 = \log_2(7 - b)^a \Rightarrow (7 - b)^a = 2^4 = 16$$

$$\Rightarrow 7 - b = 16^{\frac{1}{a}} = (4^2)^{\frac{1}{a}} = 4^{\frac{2}{a}}$$

$$\Rightarrow b = 7 - 4^{\frac{2}{a}}$$

Hence, $5 - 4^{\frac{1}{a}} = 7 - 4^{\frac{2}{a}}$

$$\Rightarrow 4^{\frac{2}{a}} - 4^{\frac{1}{a}} = 2$$

$$\Rightarrow (4^{\frac{1}{a}})^2 - 4^{\frac{1}{a}} - 2 = 0$$

Let $y = 4^{\frac{1}{a}} \Rightarrow y^2 - y - 2 = 0$

$$\Rightarrow (y - 2)(y + 1) = 0$$

$$\Rightarrow y = 2, y = -1$$

$$\Rightarrow 4^{\frac{1}{a}} = 2 \quad \text{or} \quad 4^{\frac{1}{a}} = -1 \text{ (Not valid)}$$

$$\Rightarrow 4 = 2^a$$

$$\Rightarrow 2^2 = 2^a \Rightarrow a = 2$$

$$\Rightarrow b = 5 - 4^{\frac{1}{2}} = 5 - 2 = 3$$

11. Solve $32^{x-1} = 28$

$$\Rightarrow \ln 32^{x-1} = \ln 28$$

$$\Rightarrow (x - 1) \ln 32 = \ln 28$$

$$\Rightarrow x - 1 = \frac{\ln 28}{\ln 32} = 0.96147$$

$$\Rightarrow x = 1.96147$$

$$\Rightarrow x = 1.96$$

12. Proof:

(i) $n = 1? \Rightarrow 3 = \frac{3(1)}{2}(1 + 1) \Rightarrow 3 = 3, \text{ true } n = 1.$

(ii) Assume true for $n = k \Rightarrow 3 + 6 + 9 + \dots + 3k = \frac{3k}{2}(k + 1).$

(iii) Also true for $n = k + 1?$

$$\Rightarrow 3 + 6 + 9 + \dots + 3k + (3k + 3) = \frac{3k}{2}(k + 1) + 3(k + 1)$$

$$\Rightarrow 3 + 6 + 9 + \dots + 3k + 3(k + 1) = \frac{3k(k + 1) + 2 \cdot 3(k + 1)}{2}$$

$$\Rightarrow 3 + 6 + 9 + \dots + 3k + 3(k + 1) = \frac{3}{2}(k + 1)(k + 2)$$

$$\Rightarrow 3 + 6 + 9 + \dots + 3k + 3(k + 1) = \frac{3}{2}(k + 1)[(k + 1) + 1]$$

$$\therefore \text{It is true for } n = k + 1.$$

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2.$

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of $n.$

13. Proof:

- (i) True for $n = 1$? $\Rightarrow 7$ is a factor of $8^1 + 6 = 14 \Rightarrow$ true $n = 1$.
 (ii) Assume true for $n = k \Rightarrow 8^k + 6$ is divisible by 7.
 (iii) Also true for $n = k + 1$?
 $\Rightarrow 8^{k+1} + 6 = 8^k \cdot 8 + 6$
 $= 8^k(7 + 1) + 6$
 $= 7 \cdot 8^k + (8^k + 6)$
 Since $7 \cdot 8^k$ is divisible by 7, and $(8^k + 6)$ is assumed divisible by 7,
 $\therefore 7 \cdot 8^k + (8^k + 6)$ is divisible by 7.
 (iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.
 And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.
 (v) Therefore, it is true for all positive integer values of n .


14. Prove by induction that $n^2 > 4n + 3$ for $n \geq 5, n \in N$.

Proof:

- (i) True for $n = 5$? $\Rightarrow 5^2 > 4(5) + 3 \Rightarrow 25 > 23 \Rightarrow$ true $n = 5$.
 (ii) Assume true for $n = k \Rightarrow k^2 > 4k + 3$ for $k \geq 5, k \in N$.
 (iii) Also true for $n = k + 1$?
 $\Rightarrow (k + 1)^2 = k^2 + 2k + 1$
 $> 4k + 3 + 2k + 1$ since $k^2 > 4k + 3$ (assumed)
 $= 4k + 4 + 2k$
 $> 4(k + 1) + 3$ since $2k > 3$, for $k \geq 5$
 \therefore It is true for $n = k + 1$.
 (iv) But since it is true for $n = 5$, it now must be true for $n = 5 + 1 = 6$.
 And if it is true for $n = 6$, it is true for $n = 6 + 1 = 7 \dots$ etc.
 (v) Therefore, it is true for all values of $n, n \geq 5, n \in N$.

Revision Exercise 12 (Advanced)

- 1.** (i) $3x + 4 < x^2 - 6$
 $x^2 - 3x - 10 > 0$
 Let $x^2 - 3x - 10 = 0$
 $(x - 5)(x + 2) = 0$
 $x = 5$ or $x = -2$
 Solution: $x < -2$ or $x < 5$
 (ii) $x^2 - 6 < 9 - 2x$
 $x^2 + 2x - 15 < 0$
 Let $x^2 + 2x - 15 = 0$
 $(x + 5)(x - 3) = 0$
 $x = -5$ or $x = 3$
 Solution: $-5 < x < 3$
 (iii) The solution of
 $3x + 4 < x^2 - 6 < 9 - 2x$
 is then the intersection of the above solution sets, i.e.
 $-5 < x < -2$.



2. $M = 30 \times 2^{-0.001t}$

(i) $t = 0 \Rightarrow M = 30 \times 2^{-0.001(0)} = 30 \times 2^0 = 30 \times 1 = 30 \text{ g}$

(ii) $M = 10 \Rightarrow 30 \times 2^{-0.001t} = 10$

$$\Rightarrow 2^{-0.001t} = \frac{10}{30} = \frac{1}{3}$$

$$\Rightarrow \log 2^{-0.001t} = \log \frac{1}{3}$$

$$\Rightarrow -0.001t \cdot \log 2 = \log \frac{1}{3}$$

$$\Rightarrow -0.001t = \frac{\log \frac{1}{3}}{\log 2} = -1.5849625$$

$$\Rightarrow t = \frac{-1.5849625}{-0.001} = 1584.9625$$

$$\Rightarrow t = 1585 \text{ years}$$

(iii) 1% of 30 = 0.3

$$\Rightarrow 30 \times 2^{-0.001t} = 0.3$$

$$\Rightarrow 2^{-0.001t} = \frac{0.3}{30} = 0.01$$

$$\Rightarrow \log 2^{-0.001t} = \log 0.01$$

$$\Rightarrow -0.001t \cdot \log 2 = \log 0.01$$

$$\Rightarrow -0.001t = \frac{\log 0.01}{\log 2} = -6.643856$$

$$\Rightarrow t = \frac{-6.643856}{-0.001} = 6643.856 = 6644 \text{ years}$$

3. $I = I_0 \times 10^{0.1S}$

(i) $S = 30 \Rightarrow I = I_0 \times 10^{0.1(30)}$

$$= I_0 \times 10^3 = 1000 I_0 \Rightarrow \text{Answer} = 1000$$

(ii) $S = 28 \Rightarrow I = I_0 \times 10^{0.1(28)} = I_0 \times 10^{2.8}$

$$S = 15 \Rightarrow I = I_0 \times 10^{0.1(15)} = I_0 \times 10^{1.5}$$

$$\Rightarrow \text{No. of times} = \frac{I_0 \times 10^{2.8}}{I_0 \times 10^{1.5}} = 10^{1.3} = 19.95 = 20 \text{ Times}$$

4. Solve $\log_5 x - 1 = 6 \log_x 5$.

$$\Rightarrow \log_5 x - 1 = 6 \frac{\log_5 5}{\log_5 x} = 6 \cdot \frac{1}{\log_5 x} = \frac{6}{\log_5 x}$$

Let $y = \log_5 x \Rightarrow y - 1 = \frac{6}{y}$

$$\Rightarrow y^2 - y = 6$$

$$\Rightarrow y^2 - y - 6 = 0$$

$$\Rightarrow (y - 3)(y + 2) = 0$$

$$\Rightarrow y = 3, y = -2$$

Hence, $\log_5 x = 3$ or $\log_5 x = -2$

$$\Rightarrow x = 5^3 = 125 \text{ or } x = 5^{-2} = \frac{1}{25}$$

5. Solve $(0.7)^x \geq 0.3$

$$\Rightarrow \log(0.7)^x \geq \log(0.3)$$

$$\Rightarrow x \cdot \log(0.7) \geq \log(0.3)$$

$$\Rightarrow x(-0.1549) \geq -0.5228787$$

$$\Rightarrow x(0.1549) \leq 0.5228787$$

$$\Rightarrow x \leq \frac{0.5228787}{0.1549} = 3.3755$$


$$\Rightarrow x \leq 3.38$$

6. (i)

$x =$	-3	-2	-1	0	1	2	3
(ii) $f(x) = x =$	3	2	1	0	1	2	3
$g(x) = x + 2$	5	4	3	2	3	4	5
(iii) $x + 2 =$	-1	0	1	2	3	4	5
$h(x) = x + 2 $	1	0	1	2	3	4	5

(iv) $f(x) \cap h(x) \Rightarrow x = -1$

(v) $g(x) > h(x) \Rightarrow -3 \leq x < 0$




7.

$x =$	4	5	6	7	8	9
$x - 3 =$	1	2	3	4	5	6
$\ln(x - 3) =$	0	0.7	1.1	1.4	1.6	1.8

$\ln(x - 3) = \log_e(x - 3) = y$

$\Rightarrow e^y = x - 3$

$\Rightarrow x = e^y + 3$




8.

$x = 1$	-30	-24	-18	-12	-6	0	6
$\frac{1}{4}x =$	-7.5	-6	-4.5	-3	-1.5	0	1.5
$\frac{1}{4}x + 3 =$	-4.5	-3	-1.5	0	1.5	3	4.5
$f(x) = \left \frac{1}{4}x + 3\right =$	4.5	3	1.5	0	1.5	3	4.5

Solve $\left|\frac{1}{4}x + 3\right| \geq 3$

$\Rightarrow -24 \geq x \geq 0$



9. $\frac{x^{\frac{3}{2}} - x^{-\frac{1}{2}}}{x^{\frac{1}{2}} - x^{-\frac{1}{2}}}$

$$= \frac{x^{-\frac{1}{2}}(x^2 - 1)}{x^{-\frac{1}{2}}(x - 1)} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1$$

10. Proof:

(i) True for $n = 1$? $\Rightarrow \frac{1}{(1 + r)^1} \leq \frac{1}{1 + (1)r} \Rightarrow \frac{1}{1 + r} \leq \frac{1}{1 + r} \Rightarrow \text{true } n = 1.$

(ii) Assume true for $n = k \Rightarrow \frac{1}{(1 + r)^k} \leq \frac{1}{1 + kr}$ for $k \geq 1, r > 0, k \in \mathbb{N}.$

(iii) Also true for $n = k + 1$?

$$\Rightarrow \frac{1}{(1 + r)^{k+1}} = \frac{1}{(1 + r)^k(1 + r)} \leq \frac{1}{(1 + kr)(1 + r)} \quad \text{since } \frac{1}{(1 + r)^k} \leq \frac{1}{1 + kr} \text{ (assumed)}$$

$$= \frac{1}{1 + kr + r + kr^2}$$

$$\leq \frac{1}{1 + kr + r} \quad \text{since } kr^2 > 0 \text{ for } k \geq 1, r > 0$$

$$= \frac{1}{1 + (k+1)r}$$

\therefore It is true for $n = k + 1$.

(iv) But since it is true for $n = 1$, it now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \dots$ etc.

(v) Therefore, it is true for all positive integer values of n .

11. Prove $\frac{4x}{(x+1)^2} \leq 1$ for all $x \in R, x \neq -1$.

Proof: $\frac{4x}{(x+1)^2} \cdot (x+1)^2 \leq 1(x+1)^2$

$$\Rightarrow 4x \leq x^2 + 2x + 1$$

$$\Rightarrow -x^2 + 2x - 1 \leq 0$$

$$\Rightarrow x^2 - 2x + 1 \leq 0$$

$$\Rightarrow (x-1)^2 \leq 0 \quad \text{true for all } x \in R, x \neq -1$$

Hence, $\frac{4x}{(x+1)^2} \leq 1$.

12. $(1+2k)x^2 - 10x + (k-2) = 0$

(i) Real Roots $\Rightarrow b^2 - 4ac \geq 0$

$$\Rightarrow (-10)^2 - 4(1+2k)(k-2) \geq 0$$

$$\Rightarrow 100 - 4(2k^2 - 3k - 2) \geq 0$$

$$\Rightarrow 100 - 8k^2 + 12k + 8 \geq 0$$

$$\Rightarrow -8k^2 + 12k + 108 \geq 0$$

$$\Rightarrow 8k^2 - 12k - 108 \leq 0$$

$$\Rightarrow 2k^2 - 3k - 27 \leq 0$$

Factors: $(k+3)(2k-9) = 0$

Roots: $k = -3, k = 4\frac{1}{2}$

Hence, $-3 \leq k \leq 4\frac{1}{2}$.

(ii) Sum of roots $> 5 \Rightarrow \frac{10}{1+2k} > 5$

$$\Rightarrow \frac{10}{1+2k}(1+2k)^2 > 5(1+2k)^2$$

$$\Rightarrow 10(1+2k) > 5(1+2k)^2$$

$$\Rightarrow 2(1+2k) > (1+2k)^2$$

$$\Rightarrow 2 + 4k > 1 + 4k + 4k^2$$

$$\Rightarrow -4k^2 + 1 > 0$$

$$\Rightarrow 4k^2 - 1 < 0$$

Factors: $(2k+1)(2k-1) = 0$

Roots: $k = -\frac{1}{2}, k = \frac{1}{2}$

Hence, $-\frac{1}{2} < k < \frac{1}{2}$.

13. Proof:

(i) True for $n = 1$? $\Rightarrow 1 = (1-1)2^1 + 1 \Rightarrow 1 = 0 + 1 \Rightarrow 1 = 1 \Rightarrow \text{true } n = 1$.

(ii) Assume true for $n = k \Rightarrow 1 + 2 \cdot 2 + 3 \cdot 2^2 + \dots + k \cdot 2^{k-1} = (k-1) \cdot 2^k + 1$.

(iii) Also true for $n = k + 1$?

$$\begin{aligned}
 &\Rightarrow 1 + 2 \cdot 2 + 3 \cdot 2^2 + \cdots + k \cdot 2^{k-1} + (k+1) \cdot 2^{(k+1)-1} \\
 &= (k-1)2^k + 1 + (k+1)2^{(k+1)-1} \\
 &= (k-1)2^k + 1 + (k+1)2^k \\
 &= k \cdot 2^k - 2^k + 1 + k \cdot 2^k + 2^k \\
 &= 2k \cdot 2^k + 1 \\
 &= k \cdot 2^{k+1} + 1 = [(k+1) - 1] \cdot 2^{k+1} + 1 \\
 &\therefore \text{It is true for } n = k + 1.
 \end{aligned}$$

(iv) But since it is true for $n = 1$, it is now must be true for $n = 1 + 1 = 2$.

And if it is true for $n = 2$, it is true for $n = 2 + 1 = 3 \cdots$ etc.

(v) Therefore, it is true for all positive integer values of n .

14. $u_n = (n - 20) \cdot 2^n$

$$\Rightarrow u_{n+1} = (n + 1 - 20) \cdot 2^{n+1} = (n - 19) \cdot 2 \cdot 2^n$$

$$\Rightarrow u_{n+2} = (n + 2 - 20) \cdot 2^{n+2} = (n - 18) \cdot 2^2 \cdot 2^n = (n - 18) \cdot 4 \cdot 2^n$$

$$\begin{aligned}
 \text{Hence, } u_{n+2} - 4u_{n+1} + 4u_n &= (n - 18) \cdot 4 \cdot 2^n - 4(n - 19) \cdot 2 \cdot 2^n + 4 \cdot (n - 20)2^n \\
 &= 2^n[4n - 72 - 8n + 152 + 4n - 80] \\
 &= 2^n[8n - 8n + 152 - 152] = 0
 \end{aligned}$$

15. $2 \log y = \log 2 + \log x$ and $2^y = 4^x$ for $y > 0$

$$\Rightarrow \log y^2 = \log 2x \quad \Rightarrow \quad 2^y = (2^2)^x$$

$$\Rightarrow y^2 = 2x \quad \Rightarrow \quad 2^y = 2^{2x}$$

$$\Rightarrow y^2 = y \quad \Rightarrow \quad y = 2x$$

$$\Rightarrow y^2 - y = 0$$

$$\Rightarrow y(y - 1) = 0$$

$$\Rightarrow y = 0 \text{ (Not Valid)} \text{ or } y = 1 \Rightarrow 2x = 1 \Rightarrow x = \frac{1}{2}$$

16. (i) An exponential function : $P = 40\,000(1.03)^n$

$$\begin{aligned}
 \text{(ii) } n = 12 &\Rightarrow P = 40\,000(1.03)^{12} \\
 &= 40\,000(1.42576) \\
 &= 57\,030.43 = 57\,030
 \end{aligned}$$

$$\begin{aligned}
 \text{(iii) } n = 0 &\Rightarrow P = 40\,000(1.03)^0 \\
 &= 40\,000(1) = 40\,000
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv) } P = 80\,000 &\Rightarrow 40\,000(1.03)^n = 80\,000 \\
 &\Rightarrow (1.03)^n = \frac{80\,000}{40\,000} = 2 \\
 &\Rightarrow \log(1.03)^n = \log 2 \\
 &\Rightarrow n \log(1.03) = \log 2 \\
 &\Rightarrow n = \frac{\log 2}{\log 1.03} = 23.449 = 23.5 \text{ years}
 \end{aligned}$$

17. (i) $P = Ae^{kt}$, where $A = 8000$, $P = 15\,000$, $t = 8$; find k .

$$\Rightarrow 15\,000 = 8000e^{k(8)}$$

$$\Rightarrow e^{8k} = \frac{15\,000}{8000} = 1.875$$

$$\Rightarrow \ln e^{8k} = \ln 1.875$$

$$\Rightarrow 8k(\ln e) = 8k(1) = 8k = 0.62860865$$

$$\Rightarrow k = \frac{0.62860865}{8} = 0.078576$$

Hence, $P = Ae^{kt}$, where $k = 0.078576$, $t =$ number of years
and $A = 8000$.

$$\begin{aligned}
 \text{(ii) } t = 10 &\Rightarrow P = 8000 e^{0.078576(10)} \\
 &= 8000 e^{0.78576} = 8000(2.1940738) \\
 &= 17\,552.59 = 17\,553 \\
 \text{(iii) } P = 30\,000 &\Rightarrow 15\,000 e^{0.078576t} = 30\,000 \\
 &\Rightarrow e^{0.078576t} = \frac{30\,000}{15\,000} = 2 \\
 &\Rightarrow \ln e^{0.078576t} = \ln 2 \\
 &\Rightarrow 0.078576t \ln e = 0.69314718 \\
 &\Rightarrow 0.078576t(1) = 0.69314718 \\
 &\Rightarrow t = \frac{0.69314718}{0.078576} = 8.82 = 9 \text{ years} \\
 \text{Ans} &= 2016
 \end{aligned}$$

Revision Exercise 12 (Extended-Response Questions)

1. $N = 5000e^{-0.15t}$

$$\begin{aligned}
 \text{(i) } t = 0 &\Rightarrow N = 5000e^{-0.15(0)} = 5000e^0 = 5000(1) = 5000 \\
 t = 5 &\Rightarrow N = 5000e^{-0.15(5)} = 5000e^{-0.75} \\
 &= 5000(0.47236655) \\
 &= 2361.83 = 2362
 \end{aligned}$$

\Rightarrow Claim is justified.


$$\begin{aligned}
 \text{(ii) } t = 10 &\Rightarrow N = 5000e^{-0.15(10)} \\
 &= 5000e^{-1.5} \\
 &= 5000(0.22313) \\
 &= 1115.65
 \end{aligned}$$

$$\text{(iii) } t = 0 \Rightarrow N = 5000e^{-0.15(0)} = 5000 \cdot e^0 = 5000 \cdot 1 = 5000$$

$$\begin{aligned}
 \text{(iv) } N = 100 &\Rightarrow 5000e^{-0.15t} = 100 \\
 &\Rightarrow e^{-0.15t} = \frac{100}{5000} = 0.02 \\
 &\Rightarrow \ln e^{-0.15t} = \ln(0.02) \\
 &\Rightarrow -0.15t(\ln e) = \ln(0.02) \\
 &\Rightarrow -0.15t(1) = -0.15t = -3.912 \\
 &\Rightarrow t = \frac{-3.912}{-0.15} = 26.08 = 26.1 \text{ days}
 \end{aligned}$$

(v)

$t =$	0	2	4	6	8	10	12	14	16	18	20
$N = 5000e^{-0.15t}$	5000	3704	2744	2033	1506	1116	826	612	454	336	249



2. (i) $A = 0.02(0.92)^{\frac{x}{10}}$

$$\begin{aligned}
 \text{(ii) Length} &= \frac{5}{3} = 1.66667 \\
 \Rightarrow A &= 0.02(0.92)^{\frac{1.66667}{10}} = 0.02(0.92)^{0.166667} \\
 &= 0.01972 = 0.0197
 \end{aligned}$$

$$(iii) S = (0.92)^{10-3x}$$

$$\begin{aligned} W &= S \times A \\ &= 0.02(0.92)^{\frac{x}{10}} \cdot (0.92)^{10-3x} \\ &= 0.02(0.92)^{0.1x+10-3x} \\ &= 0.02(0.92)^{10-2.9x} \end{aligned}$$

$$\begin{aligned} (iv) W &< (0.02)(0.92)^{2.5} \\ \Rightarrow 0.02(0.92)^{10-2.9x} &< (0.02)(0.92)^{2.5} \\ \Rightarrow 10 - 2.9x &< 2.5 \\ \Rightarrow -2.9x &< 2.5 - 10 = -7.5 \\ \Rightarrow 2.9x &> 7.5 \\ \Rightarrow x &> \frac{7.5}{2.9} = 2.586 \\ \Rightarrow x &> 2.59 \end{aligned}$$

$$3. (i) A = (0.83)^n I; \quad B = (0.66)(0.89)^n I$$

$$\begin{aligned} (ii) (0.83)^n I &= (0.66)(0.89)^n I \\ \Rightarrow \log(0.83)^n &= \log(0.66)(0.89)^n \\ \Rightarrow n \log(0.83) &= \log(0.66) + \log(0.89)^n \\ \Rightarrow n \log(0.83) &= \log(0.66) + n \log(0.89) \\ \Rightarrow n[\log(0.83) - \log(0.89)] &= \log(0.66) \\ \Rightarrow n[-0.0809219 + 0.05061] &= -0.180456 \\ \Rightarrow n(-0.0303119) &= -0.180456 \\ \Rightarrow n &= \frac{-0.180456}{-0.0303119} = 5.9533 \\ \Rightarrow n &= 6 \text{ stations} \end{aligned}$$

$$4. (i) P_g = A \left(1 + \frac{11}{100} \right)^t = A(1.11)^t$$

$$(ii) P_r = 10A \left(1 - \frac{5}{100} \right)^t = 10A(0.95)^t$$


$$\begin{aligned} (iii) A(1.11)^t &= 10A(0.95)^t \\ \Rightarrow \frac{(1.11)^t}{(0.95)^t} &= 10 \\ \Rightarrow \left(\frac{1.11}{0.95} \right)^t &= 10 \\ \Rightarrow (1.168421)^t &= 10 \\ \Rightarrow \log(1.168421)^t &= \log 10 \\ \Rightarrow t \log(1.168421) &= 1 \\ \Rightarrow t &= \frac{1}{\log(1.168421)} = 14.793 = 14.8 \text{ years} \end{aligned}$$

$$(iv) A(1.11)^t = 100A(0.95)^t \quad \dots [\text{Note: } P_g = 10P_r \text{ when proportions are reversed.}]$$

$$\begin{aligned} \Rightarrow \left(\frac{1.11}{0.95} \right)^t &= 100 \\ \Rightarrow (1.168421)^t &= 100 \\ \Rightarrow \log(1.168421)^t &= \log 100 \\ \Rightarrow t \log(1.168421) &= 2 \\ \Rightarrow t &= \frac{2}{\log(1.168421)} = 29.586 \\ \Rightarrow t &= 29.6 \text{ years} \end{aligned}$$

(v)

$t =$	0	5	10	15	20	25	30
$P = 10(0.95)^t$	10	7.7	6.0	4.6	3.6	2.8	2.1
$P = (1.11)^t$	1	1.7	2.8	4.8	8.1	13.6	23.9



5. $n = A(1 - e^{-bt})$

(i) Growth; as t increases, n also increases.

(ii) $t = 2$ when $n = 10\,000 \Rightarrow 10\,000 = A(1 - e^{-2b})$

$$\Rightarrow \frac{10\,000}{A} = 1 - e^{-2b}$$

$$\Rightarrow e^{-2b} = 1 - \frac{10\,000}{A}$$

$t = 4$ when $n = 15\,000 \Rightarrow 15\,000 = A(1 - e^{-4b})$

$$\Rightarrow \frac{15\,000}{A} = 1 - e^{-4b}$$

$$\Rightarrow e^{-4b} = 1 - \frac{15\,000}{A}$$

Hence, $2e^{-4b} - 3e^{-2b} + 1$

$$= 2\left(1 - \frac{15\,000}{A}\right) - 3\left(1 - \frac{10\,000}{A}\right) + 1$$

$$= 2 - \frac{30\,000}{A} - 3 + \frac{30\,000}{A} + 1 = 0$$

(iii) $a = e^{-2b} \Rightarrow e^{-4b} = (e^{-2b})^2 = a^2$

Hence, $2e^{-4b} - 3e^{-2b} + 1 = 0$

becomes $2a^2 - 3a + 1 = 0$

(iv) Factors: $(a - 1)(2a - 1) = 0$

Roots: $a = 1, a = \frac{1}{2}$

(v) $e^{-2b} = 1$

or

$$e^{-2b} = \frac{1}{2}$$

$$\Rightarrow e^{-2b} = e^0$$

\Rightarrow

$$\ln e^{-2b} = \ln \frac{1}{2}$$

$$\Rightarrow -2b = 0$$

\Rightarrow

$$-2b(\ln e) = \ln 1 - \ln 2$$

$$\Rightarrow b = 0 \text{ (Not valid)}$$

\Rightarrow

$$-2b = -\ln 2$$


\Rightarrow

$$b = \frac{1}{2} \ln 2$$

$$\begin{aligned}
 \text{(vi)} \quad n &= A(1 - e^{-t \cdot \frac{1}{2} \ln 2}) \\
 t = 2 \text{ when } n &= 10\,000 \Rightarrow 10\,000 = A(1 - e^{-2 \cdot \frac{1}{2} \ln 2}) \\
 &\Rightarrow 10\,000 = A(1 - e^{-\ln 2}) \\
 &\Rightarrow 10\,000 = A(1 - 0.5) \\
 &\Rightarrow 10\,000 = A(0.5) \\
 &\Rightarrow 20\,000 = A(1) \\
 &\Rightarrow A = 20\,000
 \end{aligned}$$

$$\text{(vii)} \quad n = 20\,000(1 - e^{-t \cdot \frac{1}{2} \ln 2})$$

$t =$	0	1	2	3	4	5
$n =$	0	5858	10 000	12 929	15 000	16 464



$$\begin{aligned}
 \text{(viii)} \quad n &= 18\,000 \Rightarrow 20\,000(1 - e^{-t \cdot \frac{1}{2} \ln 2}) = 18\,000 \\
 &\Rightarrow (1 - e^{-t \cdot \frac{1}{2} \ln 2}) = \frac{18\,000}{20\,000} = 0.9 \\
 &\Rightarrow -e^{-t \cdot \frac{1}{2} \ln 2} = 0.9 - 1 = -0.1 \\
 &\Rightarrow e^{-t \cdot \frac{1}{2} \ln 2} = 0.1 \\
 &\Rightarrow \ln e^{-t \cdot \frac{1}{2} \ln 2} = \ln 0.1 \\
 &\Rightarrow -t \cdot \frac{1}{2} \ln 2 (\ln e) = -2.3 \\
 &\Rightarrow t = \frac{2.3}{\frac{1}{2} \ln 2} \\
 &\Rightarrow t = \frac{2.3}{0.34657} = 6.636 = 6.64 \text{ hours}
 \end{aligned}$$