

Algebra 2R Problem List 2

Weronika Jakimowicz

EXERCISE 3.

Assume that $f : K \rightarrow K$ is a non-zero endomorphism (e.g. the Frobenius function). Prove that $\text{Fix}(f) = \{x \in K : f(x) = x\}$ is a subfield of the field K


EXERCISE 4.

Assume that K is a finite field, characteristic p .

(a) Prove that every irreducible polynomial $f \in K[x]$ divides the polynomial $w_n(x) = x^n - 1$ for some n not divisible by p . (hint: prove that the splitting field of f is finite.)

Let f be an irreducible polynomial $f \in K[x]$ of degree $n = \deg(f) > 0$. Without loss of generality assume that f is monic. Let $a \in L \supseteq K$ be one of its roots, where L is the splitting field of f over K . Because K is finite, I can say that $|K| = p^k$.

"Proof graph"



Lemaczysko: An irreducible monic polynomial $f \in K[X]$ is the minimal polynomial for some root a , $f(a) = 0$

As K is a field, the ring $K[X]$ is an euclidean domain. Let us suppose that $h \in K[X]$ is the minimal polynomial of a in K such that $\deg(h) < \deg(f)$. We have that there exists $p, r \in K[X]$ such that

$$f = hp + r$$

but notice that $f(a) = 0$ and $h(a) = 0$, so $r = 0$ and we would have $f = hp$ but f was irreducible.

Lemat: The splitting field of f is finite.

The ideal

$$I(a/K) = \{w \in K[X] : w(a) = 0\} = (f)$$

because f is irreducible. We showed that f is minimal in [Lemaczysko](#) and so from Remark 4.5. (below) we have that $[L : K] = \deg(f) = n$.

Lemacik: *This is not really a lemma but the third step in the diagram: $w_m(a) = 0$ for $m = p^{kn} - 1$.*

Now let us look at L^* , which is the multiplicative group of L . Because L was a field, we know that

$$|L| = p^{kn} = p^l$$

($[L : K] = n$ and there were p^k elements in K) and that

$$|L^*| = |L \setminus \{0\}| = p^l - 1.$$

Furthermore, we know that every finite group is isomorphic to the field \mathbb{Z}_p so we must have that L^* is a cyclic group with $a \in L^*$ as one of its generators. We know that $a^{p^l} = a$ will "loop back" inside of L^* and so $a^{p^l-1} = 1$ inside of L^* . This gives us the following equality:

$$w_{p^l-1}(a)a^{p^l-1} - 1 = 1 - 1 = 0$$

with $p \nmid p^l - 1$.

Lemaciuś: *Once again not a lemma but showing that f divides w_m , m as above.*

What remains now is to show that $f|w_m$. Suppose that this is untrue and that their "gcd" is equal to 1. Then by Bezout's identity we have that there exist $c, d \in K[X]$ such that

$$f(x)c(x) + w_m(x)d(x) = 1$$

but for $x = a$ we would have $0 = 1$ which is a contradiction. Hence, one has to divide the other. f is irreducible so it cannot be divided by anything but itself and so $f|w_m$.

Remark 4.5. *Suppose that $I(a/K) = (f)$ and f is monic. Then:*

1. f is the minimal monic polynomial such that $f(a) = 0$
2. $\deg(f) = [K(a) : K]$, thus the degree of the minimal polynomial is equal to the dimension of the linear space $K(a)$ over K .

EXERCISE 5.

(a) *Prove that if $K \subseteq L$ are finite fields, $|K| = p^m$, $|L| = p^n$, then $m|n$.*

Let $[L : K] = d$. Then we have that the basis of L over K has d elements. Every element of L can be expressed as a linear combination of elements from the basis with coefficients from K . There are

$$|K|^d = p^{md}$$

such combinations. Hence $|L| = p^{md} = p^n \implies n = md \implies m|n$.

(b) *Prove that every field with p^n elements contains a unique subfield with p^m elements, where $m|n$.*

"Proof graph" of existence

3. if $\text{char}(K) = p > 0$ then $p \nmid n$.

EXERCISE 6.

Let $F(p^n)$ be a field with p^n elements. From Problem 5 it follows from that

$$F(p) \subseteq F(p^2) \subseteq F(p^{3!}) \subseteq \dots \subseteq F(p^{n!}) \subseteq \dots$$

(after suitable identifications of isomorphic fields). Let

$$F = \bigcup_{n>0} F(p^{n!})$$

Prove that the field F is algebraically closed. (hint: use Problem 4.)

A field is algebraically closed if every non-constant polynomial $f \in F[X]$ has a root in F .

"Proof graph"

$$\begin{array}{c} \text{Ex. 4: } (\forall f \in F[X]) \text{ } f\text{-irreducible} \implies f|w_m \\ \downarrow \\ (\forall n \in \mathbb{N})(\exists a_1, \dots, a_n \in F) \ w_n(a_i) = 0. \ i = 1, \dots, n \\ \downarrow \\ w_n(a_i) = 0 \implies f(a_i) = 0 \text{ for some } i \in \{1, \dots, n\} \end{array}$$

Because all polynomials in $F[X]$ are either irreducible or a product of irreducible polynomials, it is sufficient to show that every irreducible polynomial in $F[X]$ has a root in F . Let $f \in F[X]$ be irreducible and $n = \deg(f)$. From Ex. 4 we know that $f|w_{p^{nk}-1}$ for some $k \in \mathbb{N}$ and so they must have a common root. Thus, it will be sufficient to show that all roots of $w_{p^{nk}-1}$ are within F .

Take $n \in \mathbb{N}$ and consider $w_n(x) \in F[X]$. The field $F(p^k)$ such that $n < p^k$ will have all roots of $w_n(x)$. But $F(p^k) \subseteq F$ so we have that F also contains all roots of $w_n(x)$.

The above reasoning was conducted for arbitrary chosen n , so it will be true for $p^{nk} - 1$ and so F contains all roots of $p^{nk} - 1$, meaning that f contains at least one root in F and so F is algebraically closed.