

Algebra 2R

Problem List 2

Weronika Jakimowicz

EXERCISE 4.

Assume that K is a finite field, characteristic p .

(a) Prove that every irreducible polynomial $f \in K[x]$ divides the polynomial $w_n(x) = x^n - 1$ for some n not divisible by p . (hint: prove that the splitting field of f is finite.)

Let f be an irreducible polynomial $f \in K[x]$ and $n = \deg(f) > 0$ and let $a_1, \dots, a_r \in L \supseteq K$ be its roots, where L is the splitting field of f over K . Because K is finite, I can say that $|K| = q$.

For my convenience, I will consider $g = b_n^{-1}f$, where b_n is the leading coefficient in f . So now g is a monic polynomial and considering the splitting field of f is the same as considering the splitting field of g - I just multiplied a polynomial by a nonzero constant.

Lemaczysko: An irreducible polynomial $g \in K[X]$ is the minimal polynomial for some root a , $f(a) = 0$

As K is a field, the ring $K[X]$ is an euclidean domain. Let us suppose that $h \in K[X]$ is the minimal polynomial of a in K such that $\deg(h) < \deg(g)$. We have that there exists $p, r \in K[X]$ such that

$$f = hp + r$$

but notice that $f(a) = 0$ and $h(a) = 0$, so $r = 0$ and we would have $f = hp$ but f was irreducible.

Lemat: The splitting field of g (equivalently, of f) is finite.

We will construct the splitting field of K as such:

$$L_1 = K(a_1)$$

$$L_2 = L_1(a_2)$$

$$L_i = L_{i-1}(a_i)$$

and then $L = L_r$.

1. $[L_1 : K] = n$. The ideal

$$I(a_1/K) = \{w \in K[X] : w(a_1) = 0\} = (g)$$

because g is irreducible. We showed that g is minimal in **Lemaczysko** and so from Remark 4.5. (below) we have that $[L_1 : K] = \deg(g) = n$.

2. $[L_{i+1} : L_i] = n$. Once again, g is irreducible over L_i (because not all roots of g are in L_i)

$$I(a_{i+1}/K) = \{w \in K[X] \subseteq L_i[X] : w(a_{i+1}) = 0\} = (g)$$

and it follows from Remark 4.5. (once again) that $[L_{i+1}, L_i] = \deg(g) = n$.

Now, using Fact 4.6. (even below) We have that

$$[L : K] = [L_r : L_{r-1}][L_{r-1} : L_{r-2}] = \dots = \prod_{i=1}^r [L_i : L_{i-1}] = n^r < \infty.$$

Let $m = p^k + 1$ such that $m > [L : K]$. It is clear that $p \nmid m$. Now, consider the polynomial

$$w_m(x) = x^m - 1 = (x - 1) \underbrace{(x^{m-1} + x^{m-2} + \dots + x + 1)}_{v_m(x)}$$

and we will show that $(x - a_i) \mid v_m$ for every $i = 1, \dots, r$.

Remark 4.5. Suppose that $I(a/K) = (f)$ and f is monic. Then:

1. f is the minimal monic polynomial such that $f(a) = 0$
2. $\deg(f) = [K(a) : K]$, thus the degree of the minimal polynomial is equal to the dimension of the linear space $K(a)$ over K .

Fact 4.6 Let $K \subseteq L \subseteq M$ be extensions of fields. Then

$$[M : K] = [M : L][L : K]$$