




Supplementary Table S1. The literature search identified 237 records from PubMed, Embase, Scopus, Web of Science, and the Cochrane Central Register of Controlled Trials (CENTRAL) (see [Supplementary Table S2a](#)). After removal of 45 duplicate records (see [Supplementary Table S2b](#)), 192 titles and abstracts were screened according to the predefined eligibility criteria. Potentially relevant reports underwent full-text assessment by two independent reviewers. Reports were excluded if they investigated an inappropriate intervention, used an ineligible study design, addressed diseases outside the predefined scope of this review, or if the full text could not be retrieved.

The database searches yielded eight eligible published clinical studies (see [Supplementary Table S3](#)). An additional six eligible clinical studies were identified through hand searching, resulting in a total of 14 studies included in the qualitative synthesis ([Tables 1, 2](#)).

The CENTRAL search required particular consideration because it indexed both published articles and trial registry

records. Twenty-five clinical records were identified, of which 18 were considered potentially relevant. These comprised five published full-text articles included in the review and 13 trial registry records referring to completed and published, ongoing, completed but unpublished, or otherwise unavailable studies. Seven records did not meet the predefined eligibility criteria. Registry records without an eligible published full-text article available by the final search date were not included in the qualitative synthesis.

3.2 In gingivitis/mucositis

The characteristics of the included clinical studies are summarized in [Table 1](#). Detailed domain-specific RoB 2 assessments and the rationale for the overall judgements are presented in [Supplementary Tables S4, S5](#).