

Quantum

↳ confinement: when physical dim $\propto$ de Broglie

↳ confinement $\propto \frac{1}{L^n}$

- i.e. 10nm: weak conf.

- $d \approx 4$ nm: strong conf.

2(a) =

(b) $p = \frac{h}{\lambda}$ $E = hc/\lambda$
 $\lambda = \frac{hc}{E}$
 $p = \frac{h}{\lambda} = \frac{hE}{hc}$

F-N Tunneling

$$T \propto \rho \omega^2 \int_{-\infty}^{\infty} u(x) dx$$

$\uparrow$, $T \uparrow$

Calculus

Interference

reflecting the, MO

~~NIL~~ → requires more
in the object → DATA

low set of thick, - Aug 19

Queso SEM: BFE, SE

~~NA (4140) 4-1-1~~
~~Winter feed program~~

Energy distribution

simple algorithm

2. EELS (capture secondary & inelastic)

Step: Focussed ion beam

is a connection $\rightarrow$ (Emission)

$$q = \Delta f = \frac{f_0}{\sqrt{2}} \Delta f$$

force $\propto 1/r^2$ (Coulomb) +
in electric field

rating model:

9) Feeds. cont. Next

Stau nach:

2

TEAM - (card)

2011/10  HALL & FRESH
FRANKE & BROSCH

Handwritten scribbles and lines.

(Retention Hold & Data Damage)

Gate-Gate-Gate-Gate

Fe_2O_3 , NiO

1 = 1000

11 $\rightarrow$ can resist

$$N_{\text{MAN}} \approx N_{\text{TOT}} - N_{\text{H}_2\text{O}}$$

Applied volt. $\uparrow$ $\rightarrow$ $N_1(E_F)$
current $\uparrow$ $\rightarrow$ $N_1(E_F)$

Restaurant → less than 1000

Elektronenstrom
Ic = Strom im TND (Hohlrohr)

→ perlopa

3. TAT. (ATC)

Wahlkreis (Wahlbezirk) (FN):