

of E
must be continuous

(16/1/25)


(Mag & LC)

Orbital & spin L momentum.

Orbital motion of e around the nucleus known as orbital L momentum.

$l \rightarrow$ orbital L momentum quantum no.

$m_l \rightarrow$ orbital magnetic moment quantum no.



$$\hbar L = \vec{r} \times \vec{p}$$


$$= -i\hbar(\vec{r} \times \nabla)$$

$$\hbar L_z = i\hbar \left[\hat{x} \frac{\partial}{\partial y} - \hat{y} \frac{\partial}{\partial x} \right]$$

$$\hbar L_z |\psi\rangle = i\hbar \frac{\partial}{\partial \phi} \hbar e^{im_l \phi} = m_l \hbar e^{im_l \phi}$$

$$\begin{aligned} \hbar^2 l(l+1) |\psi\rangle &= \hbar^2 l(l+1) |\psi\rangle \\ &= \left\{ \hbar \sqrt{l(l+1)} \right\}^2 |\psi\rangle \end{aligned}$$

$$\hbar L_z |\psi\rangle = m_l \hbar |\psi\rangle$$



(P1)

$$L_{\pm} = L_x \pm iL_y$$

$$L_{\pm} |\psi\rangle$$

$$= \sqrt{l(l+1) - m_l(m_l \pm 1)} \hbar |\psi\rangle$$

$$\vec{\mu} = \gamma \vec{L}$$

$\vec{\mu} \parallel \vec{L}$ should be \parallel .

$$(\mu_z) = \gamma L_z \quad (\text{projection along } z \text{ (direc}^n \text{ of } \vec{B}))$$

$$= \gamma m_l \hbar$$

$$= \left(\frac{e \hbar}{2m} \right) m_l$$

$$= m_l \mu_B \quad (\text{Bohr magneton})$$

$$\mu_B = \frac{e \hbar}{2m}$$

Total μ_z (mag dipole moment):

$$\vec{\mu} = \gamma \sqrt{L(L+1)} \hbar$$

$$= \left(\frac{e \hbar}{2m} \right) \sqrt{L(L+1)} = \sqrt{L(L+1)} \mu_B$$

Spin L momentum.

Intrinsic L momentum is called SPIN;

$$\hat{S}^2 |s, m_s\rangle = \hbar^2 s(s+1) |s, m_s\rangle$$

$$S_z |s, m_s\rangle = m_s \hbar |s, m_s\rangle$$

component of \vec{S} along field axis.
magnitude of total $\mu_s = \hbar \sqrt{s(s+1)}$.

$$(\mu_s)_z = \gamma g m_s \hbar$$


$$= \left(\frac{e \hbar}{2m} g \right) m_s = g \mu_B m_s$$

\rightarrow Lande g factor.

$$g = 2 \left(1 + \frac{\alpha}{2} \right)$$

$$\alpha \text{ (fine structure const)} = \frac{e^2}{4\pi\epsilon_0 \hbar c}$$

$$g = 2.0023$$



Zeeman Splitting

$$(\mu_z)_2 = -g \mu_B m_s$$

(B' along z) Energy = $-(\mu_z B)$.

$$= -(\mu_z)_2 \cdot B'$$

$$= g \mu_B m_s \cdot B'$$

$$E_1 = \frac{g \mu_B B}{2}$$

$$E_2 = -\frac{g \mu_B B}{2}$$



This is Zeeman splitting.

$$\vec{S} = \begin{bmatrix} S_x \\ S_y \\ S_z \end{bmatrix}$$

$$[S_x, S_y] = i \hbar S_z$$

$$[S^2, S_z] = 0$$

$$[S^2, S_x] = 0$$

$$[S_x, S_z] = 2 \hbar S_y$$

$$[S_z, S_x] = \pm \hbar S_y$$

$$\hat{S}_x = \frac{\hbar}{2} \hat{\sigma}_x$$

$$\hat{S}_y = \frac{\hbar}{2} \hat{\sigma}_y$$

$$\sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$$

$$\sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

An Atom in a magnetic field.

Magnetization & susceptibility.

$$\vec{H}_0 = \sum_{i=1}^Z \left(\frac{\vec{p}_i^2}{2m_e} + U_i \right)$$

$$\vec{B} = \nabla \times \vec{A}$$

~~Electromagnetic~~

Let's assume that the interaction of spin or L/S moment & their em are completely ignored.
electron momentum \longleftrightarrow Canonical momentum

$$\vec{p} = \vec{p}_i + e\vec{A}(\vec{r}) \quad | \quad T_0 = \frac{p^2}{2me}$$

$$\lambda \text{ to unperturbed KE} = \frac{p^2}{2me} = \left(\vec{p} + e\vec{A} \right)^2 / 2me$$

In the presence of magnetic field.

$$\hat{H} = \sum_i \left\{ \frac{(\vec{p}_i + e\vec{A}(\vec{r}_i))^2}{2m} + V_i \right\} \rightarrow \text{perturbed.}$$

$$\hat{f} = i\hbar \nabla \rightarrow i\hbar (\partial_x \hat{f} + \partial_y \hat{f} + \partial_z \hat{f})$$

$$\hbar \nabla \vec{A} \rightarrow i\hbar (\nabla \cdot \vec{A}) \hat{f} - i\hbar \vec{A} \cdot \nabla \hat{f} = \boxed{i\hbar \vec{A} \cdot \nabla \hat{f}}$$

$$\begin{aligned} (\vec{p}_i + e\vec{A})^2 &= p_i^2 + e^2 A^2 + 2(\vec{p}_i \cdot \vec{A})e + 2(\vec{A} \cdot \vec{p}_i)e \\ &= p_i^2 + e^2 A^2 + \boxed{2(\vec{A} \cdot \vec{p}_i)e} \xrightarrow{\text{TO}} \end{aligned}$$

Choose a gauge such that $\vec{A}(\vec{r}) = \frac{\vec{B} \times \vec{r}}{2}$

$$\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k}$$

let \vec{B} be along \hat{z} .

$$\vec{B} = \nabla \times \vec{A} \Rightarrow \begin{bmatrix} \partial_x A_z - \partial_z A_x \\ \partial_y A_z - \partial_z A_y \\ \partial_x A_y - \partial_y A_x \end{bmatrix}$$

$$A_z = \frac{1}{2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \cdot \vec{r} = \frac{1}{2} (x^2 + y^2)$$

$$A_x = -\frac{By}{2}, \quad A_y = \frac{Bx}{2}$$

$$B_x = \partial_x A_z - \partial_z A_x = 0$$

$$B_y = \partial_y A_z - \partial_z A_y = 0$$

$$B_z = \partial_x A_y - \partial_y A_x = B$$

$$\vec{B} \Rightarrow \text{for } \vec{A} = \frac{\vec{B} \times \vec{r}}{2}$$

$\vec{A} \leftrightarrow \frac{\vec{B} \times \vec{r}}{2}$
gauge invariant
gauge choice

15. A.p for our gauge

$$\begin{aligned}
 &= i\hbar \left(A_x \frac{\partial}{\partial x} + A_y \frac{\partial}{\partial y} + A_z \frac{\partial}{\partial z} \right) \\
 &= i\hbar \left(-\frac{B}{2} \left\{ y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right\} \right) \\
 &= \left(-\frac{\hbar}{2} \right) \left(\frac{B}{2} \right) \left[y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y} \right] \rightarrow L_z
 \end{aligned}$$

$$\hat{H} = \hat{H}_0 + \sum_i \frac{e^2 A^2}{2m_e} + e \hbar \frac{1}{2} \vec{L} \cdot \vec{B}$$

$$\begin{aligned}
 \hat{H} &= \hat{H}_0 + \left(\frac{e\hbar}{2m_e} \right) \vec{L} \cdot \vec{B} + \frac{e^2}{8m_e} \sum_i (B \cdot \vec{r}_i)^2 \\
 &= \hat{H}_0 + \frac{e\hbar}{2m_e} (\vec{L} \cdot \vec{B}) + \frac{e^2 B^2}{8m_e} \sum_i (x_i^2 + y_i^2)
 \end{aligned}$$

If spin is considered

$$\hat{H} = \hat{H}_0 + \frac{e\hbar}{2m_e} (\vec{L} + g\vec{S}) \cdot \vec{B} + \frac{e^2 B^2}{8m_e} \sum_i (x_i^2 + y_i^2)$$

This is the effect of atomic spin magnetic moment.

paramagnetic term.

diamagnetic term.

The energy shift produced by the above eqn is quite small

$$E_n \approx E_n + \Delta E_n.$$

$$\Delta E_n = \langle n | \Delta H_B | n \rangle + \sum_{n' \neq n} \frac{|\langle n' | \Delta H_B | n \rangle|^2}{E_n - E_{n'}}$$

$$= \mu_B \langle n | L + gS | n \rangle \quad \xrightarrow{\text{PM}} \quad \text{---} \quad \xrightarrow{\text{PM}}$$

$$+ \frac{e^2 B^2}{8m_e} \langle n | \sum_i (x_i^2 + y_i^2) | n \rangle \quad \text{---} \quad \xrightarrow{\text{DM}}$$

$$+ \frac{\mu_B^2 B^2}{E_n - E_{n'}} |\langle n | (L + gS) | n' \rangle|^2$$

$$= \mu_B B \langle n | L + g_s | n \rangle + \frac{e^2 B^2}{8 m_e} \langle n | \sum_{i=1}^2 (x_i^2 + y_i^2) | n \rangle$$

(1st) 2nd
(PM) (DM)

and (higher order term)

$$+ \frac{|\langle n | \mu_B B (L + g_s) | n' \rangle|^2}{E_n - E_{n'}}$$

Van-Vleck Term

Date: 17/01/2025

$$\Delta E_n = \mu_B B \cdot \langle n | L + g_s | n \rangle + \frac{e^2 B^2}{8 m_e} \langle n | \sum_{i=1}^2 (x_i^2 + y_i^2) | n \rangle$$

PM

$$+ \frac{|\langle n | \mu_B B (L + g_s) | n' \rangle|^2}{E_n - E_{n'}}$$

Dominating Term,

linear in B, is going to be much larger than the other 2 terms

$$\mu_B B \langle n | L + g_s | n \rangle \approx \bigcirc (\mu_B B)$$

$$\mu_B = \frac{e \hbar}{2 m_e}$$

$$\sim \frac{\hbar e B}{2 m_e} \sim \hbar \omega_c$$

$$\bigcirc 10^{-4} \text{ eV}$$

when $B = 1 \text{ tesla}$

This energy shift is very small

Diamagnetic Term \rightarrow

$$-\frac{e^2 B^2}{8mc} \langle n | \sum_i (x_i^2 + y_i^2) | n \rangle$$

$$\sim \frac{e^2 B^2}{8mc} \left(\frac{eB}{me} \right)^2 \cdot me^2 \cdot a_0^2$$

$$\because a_0 = \text{atomic radius} \\ = 5 \text{ \AA}$$

This term is smaller than the linear in B term, by a factor of 10^{-5} (smaller than the linear term) even for $B = 1$ tesla.

☆

Inert gas

$$\left\{ \begin{array}{l} \text{He} \rightarrow \sum L_i = L = 0 \\ \text{Ne} \\ \text{Ar} \\ \vdots \end{array} \right. \quad \begin{array}{l} \sum S_i = S = 0 \end{array}$$

Diamagnetic contribution (only contribution)

☆ Van-Vleck Term

↓
Also very small (since

$$\frac{\hbar \omega_c}{\Delta}$$

$$n \neq n'$$

$\text{Na}^+ \rightarrow$ no unfilled shell \rightarrow Fully filled shell

• Pure Diamagnetic Term —

☆ $PM=0$, Only Diamagnetic term \rightarrow

First order shift in the ground state energy due to diamagnetic term \rightarrow



$$\Delta E_0 = \frac{e^2 B^2}{8mc} \sum_{i=1}^2 \langle 0 | x_i^2 + y_i^2 | 0 \rangle \quad \text{--- (1)}$$

where, $|0\rangle \rightarrow$ ground state wave function

☆ electronic orbit

revolution of e^-

L



μ precession (Additional motion in opposite direction)

origin of diamagnetism

Lenz's law

$$\mu_{\text{precession}} = \text{diamagnetic moment} = -\frac{e\omega_L}{2} \langle R^2 \rangle$$

$$\mu = -ve$$

If we assume spherically symmetric atom —

$$\langle x_i^2 \rangle = \langle y_i^2 \rangle = \langle z_i^2 \rangle = \frac{1}{3} \langle r_i^2 \rangle$$

$$x_i^2 + y_i^2 + z_i^2 = r_i^2$$


$$\therefore \langle 0 | x_i^2 | 0 \rangle = \langle 0 | y_i^2 | 0 \rangle = \langle 0 | z_i^2 | 0 \rangle$$

\therefore from eqn 1 \rightarrow


$$\Delta E_0 = \frac{e^2 B^2}{8 m e^2} \sum_{i=1}^N \frac{2}{3} \langle 0 | r_i^2 | 0 \rangle$$

From the Thermodynamics —

$$dF = -SdT - PdV - MdB$$


 $M = - \left(\frac{\partial F}{\partial B} \right)_{T, V}$
 Diamagnetic magnetization

F = Helmholtz free energy


 $S = - \left(\frac{\partial F}{\partial T} \right)_{V, B}$


 $\chi = M/H$

$$\frac{\partial M}{\partial H} = -4\pi \frac{\partial^2 F}{\partial B^2}$$

- Consider a solid composed of N ions each with Z electron of mass m_e in volume V with all shell filled —

$$M = -\frac{\partial E}{\partial B} = -\frac{N}{V} \cdot \frac{\partial}{\partial B} (4E_0)$$

$$= -\frac{N}{V} \cdot \frac{e^2 B}{6m_e} \sum_{i=1}^Z \langle r_i^2 \rangle$$

at ground state

$$\chi_{\text{dia}} = M/H = \mu_0 M/B \quad \left[\begin{array}{l} \text{as } B = \mu_0 H \\ H = B/\mu_0 \end{array} \right]$$

$$= -\frac{\mu_0 N}{V} \cdot \frac{e^2}{6m_e} \cdot \sum_{i=1}^Z \langle r_i^2 \rangle$$

In S.I. \rightarrow unitless

In C.G.S. $\chi_{\text{dia}} = -\frac{4\pi N e^2}{c^2 6m_e V} \sum_{i=1}^Z \langle r_i^2 \rangle$

- $\chi_{\text{dia}} = -ve$ in sign

Temp independent
Property of all matter

$$\chi_{\text{dia}} \propto \langle r_i^2 \rangle$$


$\langle r_i^2 \rangle \rightarrow$ mean square of electron distance from nucleus

II Z_{eff} = no. of electron in the outer shell of an atom ion and r = measured ionic radius,

All the electron in the outer shell of the ion have roughly, same value of $\langle r_i^2 \rangle$

$$\sum_{i=1}^{Z_{eff}} \langle r_i^2 \rangle \sim Z_{eff} r^2$$

$$\therefore \chi_{dia} \propto Z_{eff} \times r^2 \quad \text{[Langevin]}$$



Larmour Diamagnetic Calculation.

- Not all electrons in an atom ion have same $\langle r^2 \rangle$ but the agreement is quite impressive.

Measured diamagnetic susceptibility —

$$\chi_m \propto \chi_{\text{eff}} \cdot n^2 \quad [\text{plot shows linearity}]$$

For Ne,

$$Z = 10, \quad k = 2, \quad l = 8$$

$$\langle r_i^2 \rangle \approx (10^{-8})^2 \text{ cm}^2 = 10^{-16} \text{ cm}^2$$

$$N = 5 \times 10^{22} \text{ cm}^{-3}$$

$$\chi = 10^{-6}$$

$$J|0\rangle = L|0\rangle = S|0\rangle = 0$$

- Larmor Diamagnets

↑ Solid composed of...
atoms with all electron shell filled.

Negative
A Ions

Elements

Susceptibility (molar susceptibility)

F⁻

$$-9.4 \times 10^{-6} \text{ cm}^3/\text{mol}$$

Cl⁻

$$-24.2 \times 10^{-6}$$

Br⁻

$$-34.5 \times 10^{-6}$$

I⁻

$$-50.6 \times 10^{-6}$$

Positive
Ions

Li ⁺	-0.7×10^{-6}	cm ³ /mol
Na ⁺	-6.1×10^{-6}	
K ⁺	-14.6×10^{-6}	
Rb ⁺	-22.0×10^{-6}	
Ca ²⁺	-35.1×10^{-6}	

Inert
gas

He	-1.9×10^{-6}
Ne	-7.2×10^{-6}
Ar	-19.4×10^{-6}
Kr	-28.0×10^{-6}
Xe	-43.0×10^{-6}

↓
radius

• density: 2220 kg/m³

$$\sqrt{(n^2)_{m}}$$