Supplementary Material

Adsorption of Methylene Blue Dye in the Thiol-Norbornene Crosslinked Polyvinyl Alcohol Hydrogel Surface: A Molecular Dynamics Study

Rontu Das, Debashis Kundu *

Department of Chemical Engineering, Institute of Chemical Technology Marathwada Campus, Jalna, Maharashtra, 431203, India

* Corresponding author's email: d.kundu@marj.ictmumbai.edu.in

Table/Figure No.	Description	Page No.
Figure S1.	Average number of H-bonds (a) polymer to methylene blue (b) solvent to methylene blue	S2
Figure S2.	. RDF between the different moiety of PVA-NTC framework to sulfur moiety of methylene blue (a) unreacted ene group presents in norbornene to the thiol moiety of methylene blue (b) oxygen moiety of norbornene to the thiol moiety of methylene blue (i) Schematic representation of norbornene functionalised PVA chain of PVA-NTC polymer and methylene blue. (II) Schematic representation of norbornene functionalised PVA chain of PVA-NTC polymer and methylene blue.	S3




Figure S1. Average number of H-bonds (a) polymer to methylene blue (b) solvent to methylene blue




Figure S2. RDF between the different moiety of PVA-NTC framework to sulfur moiety of methylene blue (a) unreacted ene group presents in norbornene to the thiol moiety of methylene blue (b) oxygen moiety of norbornene to the thiol moiety of methylene blue (i) Schematic representation of norbornene functionalised PVA chain of PVA-NTC polymer and methylene blue. (II) Schematic representation of norbornene functionalised PVA chain of PVA-NTC polymer and methylene blue.