Supplementary Material

Lithium Ion Transportation in Lignin Infused Polyvinyl Alcohol Based Eutectogel: A Molecular Dynamics Framework

Aradhana V. Thombre, Rontu Das, Debashis Kundu *

Department of Chemical Engineering, Institute of Chemical Technology Marathwada Campus, Jalna, Maharashtra, 431203, India

* Corresponding author's email: d.kundu@marj.ictmumbai.edu.in

Table/Figure No.	Description	Page No.
Figure S1.	Average number of H-bonds (a) PVA-PVA (b) EG-EG (c) PVA-EG (d) Lignin-lignin (e) Lignin-PVA (f) Lignin-EG	S2
Figure S3.	RDF of (a) H^+ (EG) $-O^-$ (lignin) (e) H^+ (EG) $-OCH_3^-$ (lignin) (d) H^+ (EG) $-OH^-$ (lignin)	S6
Figure S4.	RDF of (a) H^+ (PVA) $-O^-$ (lignin) (e) H^+ (PVA) $-OCH_3^-$ (lignin) (d) H^+ (PVA) $-OH^-$ (lignin)	S7
Figure S5.	MSD curve for EG moieties present in systems (a) 0-L (b) 1-L (c) 2-L (d) 3-L	S8




Figure S1. Average number of H-bonds (a) PVA-PVA (b) EG-EG (c) PVA-EG (d) Lignin-lignin (e) Lignin-PVA (f) Lignin-EG




Figure S3: RDF of (a) H^+(EG) -O^- (lignin) (e) H^+(EG) -OCH_3^-(lignin) (d) H^+(EG) -OH^- (lignin)




Figure S4: RDF of (a) H^+(PVA) -O^- (lignin) (e) H^+(PVA) -OCH_3^-(lignin) (d) H^+(PVA) -OH^- (lignin)




Figure S5: MSD curve for EG moieties present in systems (a) 0-L (b) 1-L (c) 2-L (d) 3-L