

Breast cancerReceived in formalin is a (right / left) ☐ modified radical mastectomy specimen☐ simple mastectomy specimen☐Measuring18x.....9.....cm. ☐ with axillary content , x x cm.The skin ellipse ,15..... x cm⁷ and ☒ appears normal.....☐ shows an old surgical scar cm in length at (areola / upper / lower / inner / outer) (quadrant).☐ shows an ulceration x cm . at (areola / upper / lower / inner / outer) (quadrant).The nipple ☒ is everted . ☐ shows inverted . ☐ shows ulceration

There (is a / are / two / are multiple)

☒ infiltrative firm yellow white mass3.6..... x3..... x2.8..... cm.☐ well-defined firm white mass with slit like appearance x x cm.☐ previous surgical cavity with adjacent fibrous tissue x x cm.☐ previous surgical cavity with adjacent fibrous tissue x x cm. and firm yellow white residual mass , x x cm.located ☐ beneath the nipple . ☐ beneath the scar . ☐ in the central portion (subareolar) .☒ in (upper / lower / inner / outer) quadrant . ☐**Tumor is located**

.....0.7..... cm. from deep margin ,3.5..... cm. from superior margin ,
.....1..... cm. from inferior margin , cm. from medial margin ,
.....5..... cm. from lateral margin ,	and0.4..... cm. from skin .

☐ The uninvolved breast parenchyma has a fat to fibrous tissue ratio of approximately:.....The remaining of breast tissue ☒ is unremarkable . ☐☐ There are multiple lymph nodes ranging from cm . to cm . in diameter.**Representative sections are submitted as**

.....A15..... = nipple

.....A2:1.to.A4:1..... = mass

..... = old biopsy cavity with fibrosis

.....A15..... = deep resected margin

.....A15.(Inferior)..... = nearest resected margin ,

..... = sampling upper inner quadrant

..... = sampling upper outer quadrant

..... = sampling lower inner quadrant

..... = sampling lower outer quadrant

..... = sampling central region

..... = axillary lymph nodes

.....Prosecutor

Date