

MATH240 Assignment 2 - Real Analysis

Emma Hogan

53837798

October 8, 2019

Question 1

Let S be a non-empty subset of the real numbers which is bounded above. Let c be a real number and define $T = \{cx \mid x \in S\}$. Show that, if $c > 0$, then T is non empty, bounded above and that $\sup T = c\sup S$. Give an example of a set S as above, such that, with $c = -1$, the set T is still bounded above but $\sup T \neq c\sup S$.

To show T is non-empty, observe $S \neq \emptyset$, so $T = \{cx \mid x \in S\} \neq \emptyset$.

Since S is bounded above, by definition there exists some upper bound U such that $x \leq U \forall x \in S$. So, by transitivity, $cx \leq cU \forall x \in S$. Therefore, cU is an upper bound for T .

Since S is non-empty and bounded above, we know $\sup S$ exists. By definition of $\sup S$, $\forall s \in S, s \leq \sup S$. So, since c is positive, $cs \leq c\sup S$. This means that $t \leq c\sup S \forall t \in T$ and therefore $c\sup S$ is an upper bound for T .

Since T is also non-empty and bounded above, we know $\sup T$ also exists. Assume $\sup T < c\sup S$. We will find a contradiction.

Since $c > 0$:

$$\begin{aligned} \sup T &< c\sup S \\ \Rightarrow \frac{\sup T}{c} &< \sup S \end{aligned}$$

By definition of $\sup S$, for any value $X < \sup S$, $\exists s \in S$ such that $s > X$. So, $\exists s' \in S$ such that

$$\begin{aligned} s' &> \frac{\sup T}{c} \\ \Rightarrow cs' &> \sup T \end{aligned}$$

$cs' \in T$ by definition of T , so $cs' > \sup T$ is a contradiction.

$$\therefore \sup T \geq c\sup S \tag{1}$$

But $c\sup S$ is an upper bound for T .

$$\therefore \sup T \leq c\sup S \tag{2}$$

And so from (1) and (2), we know:

$$\sup T = c\sup S$$

As an example of $\sup T \neq \sup S$ when $c = -1$, take $S = \{-1, -2\}$, so $\sup S = -1$. With $c = -1$, $T = \{1, 2\}$ and $\sup T = 2$. But $c\sup S = 1 \neq 2$, so $c\sup S \neq \sup T$.

This is true for any bounded S with $|S| > 1$.

Question 2

(i) If (x_n) is a sequence such that $\lim |x_n| = 0$, prove that $\lim x_n = 0$ also.

By definition of the limit, $\lim |x_n| = 0 \Rightarrow \forall \mathcal{E} \geq 0, \exists N \in \mathbb{N}$ such that $\forall n \geq N$,

$$\begin{aligned} ||x_n| - 0| &\leq \mathcal{E} \\ \Rightarrow ||x_n|| &\leq \mathcal{E} \\ \Rightarrow |x_n| &\leq \mathcal{E} \\ \Rightarrow |x_n - 0| &\leq \mathcal{E} \\ \Rightarrow \lim x_n &= 0 \end{aligned}$$

(ii) Give an example of a sequence (x_n) that does not converge but that $(|x_n|)$ converges.

Define (x_n) by $x_n = (-1)^n$ which does not converge.

$|x_n| = |(-1)^n| = |-1|^n = 1$ so $(|x_n|)$ converges to 1.

Question 3

Let $\sum x_n$ be a series that converges absolutely. Show that $\sum x_n^2$ converges.

Since $\sum |x_n|$ converges, we know that $\lim_{n \rightarrow \infty} |x_n| = 0$. So, by definition of the limit, we know $\exists N$ such that $\forall n \geq N, |x_n| \leq 1$.

We can remove a finite number of terms for the start of a series without change its behaviour. So, we consider $\sum_{n=N}^{\infty} |x_n|$, so $|x_n| \leq 1$ for every term in the sequence.

Note that since $x_n^2 \geq 0$, $|x_n|^2 = x_n^2 = |x_n^2|$.

$|x_n| \leq 1$ so $|x_n|^2 = |x_n^2| \leq 1$.

$|x_n^2| \leq |x_n| \forall n \geq N$, so, since $\sum_{n=N}^{\infty} |x_n|$ is convergent, $\sum_{n=N}^{\infty} |x_n^2|$ is also convergent. Once again, N is just a finite number so it does not change the behaviour of the series.

Therefore x_n^2 is absolutely convergent, which implies it is convergent.

Question 4

Write question 4 here

Solve question 4 here.

Question 5

Write question 5 here

Solve question 5 here.