

Species Summary *Splendid alfonsino*

NPFC Bottom Fisheries Small Working Group

2021-10-11

Splendid alfonsino (*Beryx splendens*)

Common names: Splendid alfonsino (English); 红眼金鲷 (Chinese); キンメダイ (Japanese); 빛금눈돔 (Korean); Низкотельный берикс (Russian)

Biological Information

Global distribution ranges from tropical to temperate oceans. Historical catch records in the Emperor Seamount suggest the distribution from Nintoku (45 °N) to Hancock (30 °N). Settlement occurs following a certain period of the pelagic life stage. Adults show a vertical distribution from 200 to 800 m with diel vertical migration, feeding on crustaceans, cephalopods, and fish during the night. Limited information is available for recruitment and reproduction processes in the Emperor Seamounts, whereas the population in the Japanese coast shows 4–5 years to sexually mature and spawning occurs during summer (Shotton 2016).




Figure 1: Photographs of *Beryx splendens* on different developmental stages A) postlarva, B) juvenile, C) young, D) adult (from Watari et al. 2017)




Figure 2: Known distribution of *Beryx splendens* around NPFC waters. Points indicate observation data from original sources (AquaMaps 2019, October)

Fishery

Since the discovery of large populations of North Pacific armorhead in the Emperor Seamount in the late 1960s, Splendid alfonso has been exploited as an alternative resource to the armorhead due to the large temporal fluctuation of the armorhead population. The main fishing methods are bottom trawls and gillnets.

Historical catch record (Figure 3) shows the highest catch proportion by Japan, followed by Korea and Russia. Russia terminated their fishery nearly a decade ago. Fishing pressure somewhat reflects the recruitment condition of North Pacific armorhead. In 2010 and 2012, when high recruitment of the armorhead occurred, the annual catch decreased below 1,000 tons, whereas it increased up to 4,000 tons ever since then.

Size composition analysis from the catch data by Japanese trawlers suggests the substantial decrease in size of fish in catches over the past decade, raising the concern about recruitment overfishing (Sawada et al. 2018).




Figure 3: Historical trends of *Splendid alfonsino* catches in NPFC waters. The annual amounts of catch by each country are shown by the bar plot.




Figure 4. Historical fishing efforts for *Splendid alfonsino*. The annual fishing efforts by each country are shown by barplot. The efforts are calculated by the total fishing days operated during the year

Assessment

There are no biomass estimates available for Splendid alfonsino in NPFC waters.

An age- or length-structured stock assessment may be feasible given the life history of this species. Surplus production models developed by Japan in 2008 showed that the average fishing mortality is 20–28 % higher than the MSY level (Nishimura and Yatsu 2008). This analysis, however, remains unreliable as the estimated CPUE is biased due to target shifts between North Pacific armorhead and Splendid alfonsino and the estimated intrinsic population growth rate parameter was too high for long-lived deep-sea fish.

Data limited approaches, such as YPR or SPR analysis that do not require detailed resource parameters or fishing data, should be explored in the future.

Management

Active Management Measures

The following NPFC conservation and management measures pertain to this species:

- CMM 2021-05 For Bottom Fisheries and Protection of VMEs in the NW Pacific Ocean
- CMM 2019-06 For Bottom Fisheries and Protection of VMEs in the NE Pacific Ocean

Available from <https://www.npfc.int/active-conservation-and-management-measures>

	Status	Comment
Biological reference	Not accomplished	Not established
Stock status	Unknown	Status determination criteria not
Catch limit	Intermediate	Recommended catch, effort limits
Harvest control rule	Not accomplished	Not established
Other	Intermediate	No expansion of fishing beyond

Currently, there is no accepted harvest control rule for this species.

In 2016, the management measures were implemented, which includes limiting the fishing effort to the 2007's level, prohibiting fisheries from November to December (which corresponds to the spawning season for North Pacific armorhead) and not allowing fisheries in C-H Seamount and the southeastern part of Koko Seamount (for the protection of VMEs).

In 2019, an additional measure was adopted, which includes the regulation of the mesh size (trawl: > 10 cm) to protect juvenile fish. Still, this measure may be insufficient as the substantial catch of young fish has been reported by trawlers even after being implemented.

Data Summary

Data type	Source	Years available	Comment
Catch	Japan, trawl	1969-present	May not be publicly
	Japan,	1990-present	
	Korea, trawl	2004-2019	
	Russia,	1969-1988, 2002, 2005,	
CPUE	Japan, trawl	Varies	Log book and observer
	Japan,	Varies	Log book and observer
	Korea, trawl	2013-2019	Log book from one
Survey	Japan	2019-present	Monitoring survey:
Age data	Japan	2013-present	
	Korea	2013-2017,2019	
Length data	Japan	2018-present	Survey data, see

Maturity/fecun	Japan	2013-present	Gonad mass/GSI,
	Korea	2013-2017, 2019	

References

- Watari, S., Yonezawa, J., Takeuchi, H., Kato, M., Yamakawa, M., Hagiwara, Y., & Ochi, Y. (2017). Fisheries biology and resource management of Splendid alfonsino *Beryx splendens*. Bulletin of Japan Fisheries Research and Education Agency, 44, 1-46.
- Kaschner, K., Kesner-Reyes, K., Garilao, C., Segschneider, J., Rius-Barile, J. Rees, T., & Froese, R. (2019, October). AquaMaps: Predicted range maps for aquatic species. Retrieved from <https://www.aquamaps.org>.
- Shotton, R. (2016). Global review of alfonsino (*Beryx* spp.), their fisheries, biology and management. FAO Fisheries and Aquaculture Circular, (C1084), I.
- Sawada, K., Nishida, K., Yonezaki, S. and Kiyota, M. (2018). Review of biology and fisheries of Splendid alfonsino *Beryx splendens*, especially in the Emperor seamounts area. NPFC-2018-SSC- BF01-WP03. 26 pp.
- Nishimura, A., & Yatsu, A. (2008, October). Application of surplus-production models to splendid alfonsin stock in the Southern Emperor and Northern Hawaiian Ridge (SE-NHR). In Fifth Intergovernmental Meeting on Establishment of New Mechanism for Management of High Seas Bottom Trawl Fisheries in the North Western Pacific Ocean (NWPBT/SWG-05), Tokyo, 17-18 October 2008 (pp. 1-11).