

# Comments on Uranium Content Anomalies in Groundwaters of Fazilka District of Punjab (India) for the Assessment of Excess Cancer Risk

Munish Kumar<sup>1,2,\*</sup>

<sup>1</sup>Radiological Physics and Advisory Division, Bhabha Atomic Research Centre, Mumbai, Maharashtra, India

<sup>2</sup>Department of Physical Sciences, Homi Bhabha National Institute, Mumbai, Maharashtra, India  
E-mail: drmunishkumar@gmail.com

In a recent article in “*Research & Reviews: Journal of Oncology and Hematology*”, Prof. Virk has evaluated radiological risk due to uranium content present in the groundwater of Fazilka district for the Malwa belt of Punjab, India [1]. As per the report, the uranium content of water samples of the studied villages varied from 121.95–366.00 ppb ( $\mu\text{g l}^{-1}$ ) with an average value of 198.198 ppb ( $\mu\text{g l}^{-1}$ ) [1]. The excess cancer risk varied from  $(3.45-10.37) \times 10^{-4}$  with a mean value of  $5.01 \times 10^{-4}$  and works out to be 5 per 10,000 people or 500 per million [1].

In this regard, the author would like to add that the typical lifetime dose associated with average uranium concentration of ~200 ppb is negligible. Even for high background radiation areas (doses upto ~70 mGy/year), the studies did not reveal any excess cancer risk [2]. The author would further like to stress that it is inappropriate to estimate the number of cancer cases at such dose values using ICRP's adopted risk factor of 5 %/Sv or any other methodology especially for low dose values [3, 4]. This is in line with the ICRP as well as UNSCEAR report where it is categorically mentioned that the aggregation of very low individual doses over extended time periods is inappropriate for use in risk calculations and projections [5, 6]. In view of above, the researchers should avoid performing any risk calculation or predicting (hypothetical) number of cancer cases, as the number predicted is not of any use and against the philosophy as well as guidelines of the ICRP. The projection of such hypothetical cancer numbers unnecessarily scare public.

## REFERENCES

1. Virk HS. Uranium Content Anomalies in Groundwaters of Fazilka District of

Punjab (India) for the Assessment of Excess Cancer Risk. *Research & Reviews: Journal of Oncology and Hematology*. 2017; 6: 21–6p.

2. Nair RR, Rajan K, Akiba S, *et al.* Background radiation and cancer incidence in Kerala, India—Karanagappally cohort study. *Health Physics*. 2009; 96: 55–66p.
3. Kumar M, Datta D. Comments on Radiological risk assessment of cosmic radiation at aviation altitudes (a trip from Houston Intercontinental Airport to Lagos International Airport). *J Med Phys*. 2017; 42(4): 279–80p.
4. Kumar M. Comments on Assessment of radiation dose and excessive life-time cancer risk from the Bunair granite, northern Pakistan. *Radiat Prot Dosim*. 2017; [Epub ahead of print]. Available from: <https://doi.org/10.1093/rpd/ncx227>.
5. UNSCEAR 2016. *Report on sources, effects and risks of ionising radiation, United Nations Scientific Committee on the effects of Atomic Radiation (report to General Assembly), 1–512, January. USA: United Nations; 2017.*
6. ICRP. *The 2007 recommendations of the International Commission on Radiological Protection*. ICRP Publication 103. Ann ICRP. 2007; 37(2–4).

## Cite this Article

Kumar M. Comments on “Uranium Content Anomalies in Groundwaters of Fazilka District of Punjab (India) for the Assessment of Excess Cancer Risk”. *Research & Reviews: Journal of Oncology & Hematology*. 2017; 6(3): 1–1p.