

# F/ $\alpha$ TRACK-ETCH METHOD FOR U, Th AND ISOTROPIC DISEQUILIBRIUM STUDY OF GEOLOGICAL SAMPLES

N.P. SINGH, SURINDER SINGH and H.S. VIRK

SSNTD-Laboratory, Department of Physics, Guru Nanak Dev University  
 Amritsar-143 005, India

**Abstract:** The relation connecting concentration of uranium (U) and thorium (Th) atoms in geological samples with alpha track production rate in plastic track detector, LR-115 type 2, is used for analysis of some rock and fossil bone samples. The alpha track measurements from the samples under secular equilibrium define the total U and Th contents, while fission track measurements yield U concentration alone, thus by combining the results of both these measurements, U, Th and Isotopic disequilibrium is determined. Uranium concentration determined by fission track analysis in radioactive fossil bones is found to vary from 93.8 to 418.0 ppm. The results of U concentration as determined by alpha autoradiography are pronouncedly lower than those determined by fission track analysis indicating the presence of radioactive disequilibrium. However, the quartzite and phosphorite samples analysed using this technique show the presence of radioactive equilibrium.

## 1. INTRODUCTION

Uranium and thorium are natural alpha emitters and an emulsion detector when placed against the surface of a geological sample should record the alpha disintegrations. Coppen<sup>1</sup> had given the uranium (U) and thorium (Th) content of a sample as a function of alpha track production rate (N):

$$10^6 N = 9.20kU \quad (1)$$

$$\text{or } 10^6 N = 2.80kTh \quad (2)$$

$$\text{and } 10^6 N = (9.20U + 2.80Th)k \quad (3)$$

Eq (3) has been modified by Singh et al.<sup>2</sup> for use with solid state nuclear track detectors which can record alpha particle tracks more precisely. We used cellulose nitrate plastic (LR-115 type 2) and modified eq (3) for this plastic track recorder as follows:

$$10^6 N = (6.77U + 1.60Th)k \quad (4)$$

$$\text{where } k = \frac{0.85}{\sum \frac{CS}{A}} \quad (5)$$

In eq (5), C is the concentration of an element of atomic weight A and S is the stopping power of an alpha particle for that element. Knowing the value of k and estimating U by homogenized fission track technique, Th concentration can be easily determined from eq(4).

A comparison between the U concentrations determined by autoradiography and those by other methods (viz. fission track method, fluorimetry, etc.) allows the indication of existence of a radioactive equilibrium/disequilibrium.

In the present investigations the modified relation is used for U, Th and isotopic disequilibrium study of some radioactive Siwalik fossil bones and phosphorites and quartzites collected from Kullu area of Himachal Pradesh (India).

## 2. EXPERIMENTAL

The homogenized fission track technique<sup>3</sup> is used for uranium estimation in geological samples. In this method, the sample is powdered and pressed into a pellet (13 mm dia and 1 mm thickness). The pellet sand-wiched between lexan plastic discs is irradiated with a known dose of thermal neutrons from a reactor alongwith a suitable U-rich standard glass dosimeter pellet. The lexan plastic is then etched and scanned. A comparison of induced fission track densities recorded in the lexan covering the unknown and standard pellets gives the U content in the samples as follows<sup>4</sup>:

$$U_x = U_s T_x / T_s$$

(6)

where x and s stand for unknown and standard respectively, and T, the induced fission track density.

A sheet of LR-115 plastic is then placed against the polished surface of the sample and kept undisturbed for a period depending upon the alpha activity, to allow the alpha tracks to accumulate. After the accumulation period, the plastic detector is etched for 3 h in 2.5N NaOH at 50°C. The tracks are counted using Olympus binocular microscope at a magnification of 600X. Since in this method the concentration of U is estimated using fission track technique and concentration of Th is calculated by alpha autoradiography, this technique is called F/α techniques. The mechanism of the technique is shown in Fig.




Fig. 1. Mechanism of F/α Technique

## 3. RESULTS AND DISCUSSION

Since Th is absent<sup>6</sup> in Siwalik fossil bones, the simplified relation

$$10^6 N = 6.77 KU$$

is used for U estimation in these samples. The results of U measurements by alpha autoradiography and fission track analysis are reported in Table 1. U content values obtained using alpha-autoradiography are

**Table 1**  
U concentration by Alpha-autoradiography and fission track analysis  
in radioactive fossil bones.

Sample Number	Specimen	$10^6 N$	$U_{\alpha}$ (ppm)	$U_f$ (ppm)
<b>SAHARANPUR*</b>				
FS1	Elephant tusk	6100	58.3	118.6
FS2	Elephant tusk	2951	28.2	93.8
FS3	Molar teeth	4976	47.6	102.9
FS4	Unidentified	6620	63.3	159.7
<b>NARAINGARH</b>				
FS3	Tortise bone	7998	76.5	165.1
FS4	Unidentified	5799	55.4	119.7
<b>NALAGARH</b>				
FV1	Horn piece	15961	152.6	418.0
FV2	Elephant tusk	11180	106.9	292.8
FV3	Unidentified	6377	61.0	131.7

$U_{\alpha}$  - U content determined by alpha autoradiography

$U_f$  - U content determined by fission track analysis

Value of k for fossil bone = 15.44

\* Sample location

pronouncedly lower than those determined by fission track analysis, which reveals the presence of radioactive disequilibrium of U in radioactive fossil bones. These results thus confirms the hypothesis of secondary enrichment of U in Siwalik fossil bones reported earlier<sup>7</sup>. Our recent X-ray fluorescence spectrometric analysis<sup>8</sup> of these samples have detected Ti, V, Cu, Y and As (Table 2) only in some of the bone samples with anomalous U content (threshold value for U content in these samples is already reported<sup>9</sup> to be 56.2 ppm). These are pathfinder elements of U and their presence suggests that besides uranium, Ti, V, Cu, Y and As also got enriched in bone samples with anomalous U content. The uranium anomalies found in the samples from Saharanpur (U.P.), Naraingarh (Haryana) and Nalagarh (HP) are typical of those found in Siwalik Himalayas. Moreover, the U concentration in fossil bones is controlled by availability of uranium in contemporary environment in which the vertebrates lived and the nature of the mineralizing solution acting as the carrier of uranium.

Alpha-autoradiographic analysis predicted a linear relationship between U content and alpha track rate satisfied by the relation<sup>9</sup>.

Comparison of results shows that U determined by both these methods is in good agreement. This indicates that the U present in these samples is in radioactive equilibrium. This fact is also confirmed from the results of gamma ray spectrometry.

#### ACKNOWLEDGEMENTS

The authors acknowledge the financial assistance of Third World Academy of Sciences, Trieste (Italy) under Project No. TWAS RG 86-27.

#### REFERENCES

1. R.Coppens, P. Richard and S. Bashir. Nucl Inst Methods 147, 87 (1977)
2. N.P. Singh, M. Singh, S. Singh and H.S. Virk. Indian J Pure Appl Phys 24, 143 (1986)
3. D.E. Fisher, Anal Chem 41, 414 (1970)
4. R.L. Fleischer, P.B. Price and R.M. Walker. Nuclear Tracks in Solids-Principles and Applications, University of California Press, California, U.S.A. (1975)
5. D.E. Fisher. J Radioanal Chem 38, 477 (1977)
6. G.R. Udas and T. M. Mahadevan. Proc IAEA Symp, Vienna, 426 (1976)
7. K.K. Sharma, O.P. Sharma, V. M. Choubey and K.K. Nagpaul. Trans Nucl Soc Am 33, 148 (1983)
8. N.P. Singh, S.Singh, H.S. Virk, G.S. Lodha and K.J.S.Sawhney. Indian J Pure Appl Phys 25, 411 (1987)
9. N.P. Singh, S.Singh, M. Singh and H.S. Virk. Nucl Tracks Rad Meas 12, 793 (1986)