# Steady state era result:

# worsened performance, even with $v_x(t = 0)$ as input

11

model identifies clumps of particle displacements but exhibits a cluster bias towards these clumps and a lack of sensitivity to minority modes


Parity Plot for Best R²
steady state (vₓ-aware)
Time Step: Dt1000, R²: 0.5826, Epoch: 122
Predicted X Displacement
Actual X Displacement (standardized)
3
2
1
0
−1
−2
−3


Parity Plot for Best R²
steady state (vₓ-naive)
Time Step: Dt1000, R²: 0.6145, Epoch: 79
Predicted X Displacement
Actual X Displacement (standardized)
3
2
1
0
−1
−2
−3


Parity Plot for Best R²
steady state (vₓ-aware)
Time Step: Dt1000, R²: 0.5826, Epoch: 122
Predicted X Displacement
Actual X Displacement (standardized)
3
2
1
0
−1
−2
−3


Parity Plot for Best R²
steady state (vₓ-naive)
Time Step: Dt1000, R²: 0.6145, Epoch: 79
Predicted X Displacement
Actual X Displacement (standardized)
3
2
1
0
−1
−2
−3

# Steady state era result:

## worsened performance, even with $v_x(t = 0)$ as input







model identifies clumps of particle displacements but exhibits a cluster bias towards these clumps and a lack of sensitivity to minority modes

# Steady state era result:

## worsened performance, even with $v_x(t=0)$ as feature




$1$ MD step $=$ strain $\gamma$ of $10^{-8}$