Conclusions:

13

- GNN ability to emulate MD by using a pair of snapshots and predicting displacement in direction of shear:

- promising for transient era with $t_0 = 0$

- limited for steady state with $t_0 > 0$, even when $v_x(t_0)$ introduced as feature

- More to explore in predicting displacement:

- RMSE loss function not physical?

- is wall distance highest gradient region in feature-space? (implies success in transient era but challenges at later times)

- Given extensive literature on this question, further review is warranted

Conclusions:

- GNN ability to emulate MD by using a pair of snapshots and predicting displacement in direction of shear:
  - promising for transient era with $t_0 = 0$
  - limited for steady state with $t_0 > 0$, even when $v_x(t_0)$ introduced as feature
- More to explore in predicting displacement:
  - RMSE loss function not physical?
  - is wall distance highest gradient region in feature-space? (implies success in transient era but challenges at later times)
- Given extensive literature on this question, further review is warranted

contact: rorlija1.github.io

Thanks!

We gratefully acknowledge:

- *scientific contributions of Junhyuk Kim, Jean-Luc Ishimwe, Cacey Bester, Xiang Li*

- *support from NSF (DMR-1905474, DMR-1905737), and time on TACC via ACCESS (PHY230003, TRA100004) and Swarthmore College's Firebird Cluster.*

A.A. and D.M. gratefully acknowledge partial support of this research by NSF through the University of Wisconsin Materials Research Science and Engineering Center (DMR-2309000).