

# Derivations and Equations

## 1 Deriving the bound

The standard variational formulation is

$$\begin{aligned}\log p(y) &= \int q(z) \log \frac{p(y|z)p(z)}{q(z)} \frac{q(z)}{p(z|y)} dz \\ &= \underbrace{\mathbb{E}_{q(z)} [\log p(y|z)p(z) - \log q(z)]}_{\mathcal{L}_1} + \text{KL}[q(z) \| p(z|y)]\end{aligned}\quad (1)$$

The GP-LVM-CMF variational joint in the augmented latent space is:

$$q(z, f, u, X) = \prod_{i=1}^N \{q(z_i | f(X_i))q(f(X_i) | u)\} q(u) q(X) \quad (2)$$

which induces an intractable variational marginal

$$q(z) = \int \prod_{i=1}^N \{q(z_i | f(X_i))q(f(X_i) | u)\} q(u) q(X) df du dX. \quad (3)$$

However we can apply the variational formulation again, noting that  $q(z) = \frac{q(z|f,u,X)q(f,u,X)}{q(f,u,X|z)}$ ,

and integrating 1 wrt the auxiliary variables, to obtain a further lower bound:

$$\log p(y) = \int q(z|f, u, X) q(f, u, X) \log \frac{p(y|z)p(z)r(f, u, X|z)}{q(z|f, u, X)q(f, u, X)} \frac{q(f, u, X|z)}{r(f, u, X|z)} \frac{q(z)}{p(z|y)} df du dX dz$$

$$= \mathbb{E}_{q(z,f,u,X)} [\log p(y|z)p(z)r(f, u, X|z) - \log q(z|f, X)q(f, u, X)] \quad (4)$$

$$+ \mathbb{E}_q(z) [\text{KL}[q(f, u, X|z) \| r(f, u, X|z)]] + \text{KL}[q(z) \| p(z|y)] \quad (5)$$

in which the new auxiliary lower bound,  $\mathcal{L}_{aux}$  is given by the expression 4, and where we have introduced the auxiliary distribution  $r(f, u, X|z)$  which serves to approximate the variational posterior,  $q(f, u, X|z)$ , of the auxiliary variables conditioned on the latent variables.

We may re-express  $\mathcal{L}_{aux}$  in a way which makes use of the analytical expression for the K-L divergence between two Gaussians,  $q(f, u, x)$  and  $r(f, u, X|z)$  and, in the case that the prior of the generative model,  $p(z)$ , is also Gaussian distributed - as is the case for the continuous latent variable MLP model we'll consider first - then the bound contains a second Gaussian KL term:

$$\mathcal{L}_{aux} = \underbrace{\mathbb{E}_{q(z)} [\log p(y|z)]}_A - \underbrace{\mathbb{E}_{q(f,u,X)} [\text{KL}[q(z|f, X) \| p(z)]]}_B - \underbrace{\mathbb{E}_{q(z)} [\text{KL}[q(f, u, X) \| r(f, u, X|z)]]}_C. \quad (6)$$

## 2 Expressions

### 2.1 Generative model

### 2.2 Variational model

$$q(X) = \prod_{i=1}^N \mathcal{N}(x_{i,:} \mid 0, I_R) = \prod_{i=1}^N \prod_{r=1}^R \mathcal{N}(x_{i,r} \mid 0, 1) \quad (7)$$

$$q(u) = \prod_{m=1}^M \mathcal{N}(u_{:,m} \mid 0, K_{X_u X_u}) \quad (8)$$

$$q(f(X)|u) = \mathcal{N}(f(X) \mid K_{X X_u} K_{X_u X_u}^{-1} u, K_{X X} - K_{X X_u} K_{X_u X_u}^{-1} K_{X_u X}) \quad (9)$$

$$q(z|f(X)) = \prod_{i=1}^N \mathcal{N}(z_i \mid f(X_i), \sigma^2) \quad (10)$$

## 2.3 Auxiliary model

$$r(f, u, X|z) = q(f(X)|u, X)r(u, X|z) \tag{11}$$

$$r(u, X|z) = \tag{12}$$