SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Dogwood

1 existing; 1 proposed. Products: leaf brewing flavour - specialist use.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December-June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Buying advice: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Source: SANBI PlantZAfrica: Rhamnus prinoides

Bushman's grape

1 existing; 1 proposed. Products: edible berries; roots excluded.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Food is the berry; the same SANBI account reports poisonous tuberous roots. September-June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Buying advice: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, rhoicissus-tridentata

Rosemary

1 existing; 1 proposed. Products: fresh or dried leaves for seasoning.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Buying advice: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: DAFF: Rosemary production brochure (2012), p. 2

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Karoo boer-bean

1 existing; 1 proposed. Products: edible seeds.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Buying advice: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Little Karoo and drier Cape botanical account - edible seed production reference: May-Jun.

Source: SANBI PlantZAfrica, schotia-afra-var-afra

Kuni bush

1 existing; 1 proposed. Products: ripe dried fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Buying advice: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Source: SANBI Red List: Searsia undulata

Dune spinach

1 existing; 1 proposed. Products: young leafy tops and stems.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August-November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Buying advice: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Tetragonia decumbens

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Kinkelbossie

1 existing; 1 proposed. Products: young shoots, cooked.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September-January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Buying advice: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

1 existing; 1 proposed. Products: flower buds.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July-September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Buying advice: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Trachyandra ciliata

Pigeonwood

1 existing; 1 proposed. Products: young leaves for cooked greens.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Buying advice: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Trema orientalis

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Wild garlic

1 existing; 1 proposed. Products: leaves and edible flowers.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Buying advice: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Eastern South African botanical account - edible flowers only: Jan-Apr.

Source: SANBI PlantZAfrica, tulbaghia-violacea

Ginger

1 existing; 1 proposed. Products: fresh rhizomes; dried ginger recorded separately.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Buying advice: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Source: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Source: University of Pretoria author manuscript: Gatabazi et al. (2022)

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and ground.

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.