SYNTHETIC GRAPH SOURCE {COUNT} $&

Pele o reka kapa o qala

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Sehlopha sa tlelaemete sa dipatlisiso: Sebaka sa temo ha se eso hlake. Sehlopha sena ha se keletso e netefaditsweng bakeng sa polasi ena.

Seo o se tlalehileng ka polasi ena

Metsi nakong ya komello

Ha ho a rekotwa. Qetella phuputso ya sebaka mme o netefatse metsi pele o kgetha moralo o hlokang disebediswa tse ngata.

Serame polasing ena

Se bonweng ka serame sebakeng ha se a rekotwa. Dikarolelano tsa kgwedi tsa sathelaete di ke ke tsa tiisa matsatsi a pele le a ho qetela a serame.

Letsatsi le phallo ya metsi mobung

Letsatsi: ha ho a rekotwa.

Phallo ya metsi: ha ho a rekotwa. Diphetho tsa teko ya mobu le ditlhoko tsa sejalo ka seng di ntse di hloka ho hlahlojwa.

Direkoto tsa selemo sa tlaleho - eseng ponelopele

Dikakaretso le dikgwedi tsa selemo sa tlaleho se fetileng di hlalosa se tlalehilweng. Ha di tshepise dijo moralong o tlang.

Original eggs {year} $&

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 240 eggs.

Tshebediso ya lelapa: 0 eggs. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: Php, Phu. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Fish

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: ha ho a rekotwa.

Tshebediso ya lelapa: ha ho a rekotwa. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: ha ho a rekotwa. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Meroho e meng

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 12 kg.

Tshebediso ya lelapa: 7 kg. Ho rekisitsweng: 3 kg.

Dikgwedi tsa kotulo tse tlalehilweng: Phe, Tsh. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae

Dogwood

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: tatso ya makgasi ha ho ritela — tshebediso ya setsebi.

Tlhahlobo ya tshebediso e le dijo: netefatsa hore ke semela sefe, karolo e jewang le tokiso e nepahetseng le moeletsi wa sebaka ya nang le tsebo pele o se kenya dijong tsa lelapa. Se tlohetswe dikakaretsong tsa dijo tse tlwaelehileng le mesebetsing ya kotulo.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Keletso ya ho reka: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Mohlodi: SANBI PlantZAfrica: Rhamnus prinoides

Bushman's grape

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: monokotshwai o jewang; metso ha e kenyeletswe.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Keletso ya ho reka: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica, rhoicissus-tridentata

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Rosemary

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi a foreshe kapa a omisitsweng bakeng sa ho noka dijo.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Keletso ya ho reka: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: DAFF: Rosemary production brochure (2012), p. 2

Karoo boer-bean

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dipeo tse jewang.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Keletso ya ho reka: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Nako ya kotulo e lebeletsweng: ha ho nako e le nngwe e tiisitsweng ya mofuta wa sebaka. Dinako tsa ditshupiso tse ka tlase ha se ponelopele.

Tshupiso feela — tlaleho ya botaniki ya Little Karoo le Cape e ommeng ho feta — tshupiso ya tlhahiso ya peo e jewang: Mot–Php.

Mohlodi: SANBI PlantZAfrica, schotia-afra-var-afra

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Kuni bush

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: ditholwana tse butswitseng tse omisitsweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Keletso ya ho reka: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Mohlodi: SANBI Red List: Searsia undulata

Dune spinach

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dintlha tse ntjha tse nang le makgasi le dikutu.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Keletso ya ho reka: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Tetragonia decumbens

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Kinkelbossie

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: mahlomela a matjha, a phehilweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Keletso ya ho reka: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dithunthung tsa dipalesa.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Keletso ya ho reka: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Trachyandra ciliata

Pigeonwood

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi a matjha bakeng sa moroho o phehilweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Keletso ya ho reka: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Trema orientalis

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Wild garlic

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi le dipalesa tse jewang.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Keletso ya ho reka: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Nako ya kotulo e lebeletsweng: ha ho nako e le nngwe e tiisitsweng ya mofuta wa sebaka. Dinako tsa ditshupiso tse ka tlase ha se ponelopele.

Tshupiso feela — tlaleho ya botaniki ya botjhabela ba South Africa — dipalesa tse jewang feela: Phe–Mme.

Mohlodi: SANBI PlantZAfrica, tulbaghia-violacea

Ginger

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dikutu tse foreshe tse melang ka tlasa mobu; ginger e omisitsweng e rekotwa ka thoko.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Keletso ya ho reka: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Mohlodi: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Mohlodi: University of Pretoria author manuscript: Gatabazi et al. (2022)

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Mefuta ya dijalo e lokelang ho hlahlojwa sebakeng seo

Boloka lebitso la mofuta oo o o kgethang ka nnete. Manane a makgutshwane a dipatlisiso a hloka netefatso bakeng sa nako ya sebaka, metsi le mobu.

Tomatoes

Mofuta o rekotilweng: ha ho eso kgethwe.

Maemo a lehodimo a sebaka sena ha a eso hlake, kahoo ha ho bolelwe lenane le lekgutshwane la sebaka. Bula karete ya sejalo ho bona mehlodi le ho rekota kgetho ya hao.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.