SYNTHETIC GRAPH SOURCE {COUNT} $&

Musi ni sa athu u renga kana u thoma

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Tshigwada tsha kilima tsha ṱhoḓisiso: Fhethu ha u lima a hu athu u ṱalutshedzea. Tshigwada itshi a si nyeletshedzo yo khwaṱhisedzwaho ya bulasi iḽi.

Zwe na vhiga nga bulasi iḽi

Maḓi nga khalaṅwaha yo oma

A zwo ngo rekhodwa. Fhedzisani ṱhoḓisiso ya fhethu ni khwaṱhisedze maḓi ni sa athu u nanga pulane ine ya ṱoḓa zwishumiswa zwinzhi.

Murotholo kha bulasi iḽi

Zwe zwa vhonwa nga murotholo wa henefho a zwo ngo rekhodwa. Zwikati zwa ṅwedzi zwa sathelaithi a zwi koni u khwaṱhisedza maḓuvha a murotholo wa u thoma na wa u fhedza.

Ḓuvha na u bva ha maḓi mavuni

Ḓuvha: a zwo ngo rekhodwa.

U bva ha maḓi: a zwo ngo rekhodwa. Mvelelo dza u ṱola mavu na ṱhoḓea dza tshiliṅwa tshiṅwe na tshiṅwe zwi kha ḓi ṱoḓa u ṱolwa.

Rekhodo dza ṅwaha wa u vhiga - hu si mbonelaphanḓa

Zwivhalo na miṅwedzi zwa ṅwaha wo fhiraho wa u vhiga zwi ṱalutshedza zwo vhigwaho. A zwi fulufhedzisi zwiḽiwa kha pulane i ḓaho.

Original eggs {year} $&

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: 240 eggs.

Zwo shumiswaho muṱani: 0 eggs. Zwo rengiswaho: a zwo ngo rekhodwa.

Miṅwedzi ya khaṇo yo vhigwaho: Fulw, Fwan. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

Fish

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: a zwo ngo rekhodwa.

Zwo shumiswaho muṱani: a zwo ngo rekhodwa. Zwo rengiswaho: a zwo ngo rekhodwa.

Miṅwedzi ya khaṇo yo vhigwaho: a zwo ngo rekhodwa. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

Miṅwe miroho

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: 12 kg.

Zwo shumiswaho muṱani: 7 kg. Zwo rengiswaho: 3 kg.

Miṅwedzi ya khaṇo yo vhigwaho: Phan, Nyen. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṇu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani

Dogwood

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: muthetshelo wa maṱari musi hu tshi khou ukwa — u shumiswa nga muḓivhi.

U sedzulusa tshumiso ya zwiḽiwa: khwaṱhisedzani vhuvha, tshipiḓa tshi ḽiwaho na u lugisela ho teaho na mueletshedzi wa henefho a re na ndalukanyo ni sa athu u katela tshimela itshi kha zwiḽiwa zwa muṱa. A tshi katelwi kha zwivhalo zwa zwiḽiwa zwo ḓoweleaho na mishumo ya u kaṋa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Nyeletshedzo ya u renga: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Tshiko: SANBI PlantZAfrica: Rhamnus prinoides

Bushman's grape

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo miṯuku i ḽiwaho; midzi a i katelwi.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Nyeletshedzo ya u renga: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica, rhoicissus-tridentata

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṯuṯuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Rosemary

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari maswa kana o omiswaho a u naka zwiḽiwa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Nyeletshedzo ya u renga: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: DAFF: Rosemary production brochure (2012), p. 2

Karoo boer-bean

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mbeu dzi ḽiwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Nyeletshedzo ya u renga: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu athu u khwaṱhisedzwa tshifhinga tshithihi tsha lushaka lwa henefho. Khalaṅwaha dza ndaedzi dzi re afho fhasi a si mbonelaphanḓa.

Ndaedzi fhedzi — tshiko tsha botaniki tsha Little Karoo na Cape ho omaho nga maanḓa — ndaedzi ya vhubveledzi ha mbeu i ḽiwaho: Shun–Fulw.

Tshiko: SANBI PlantZAfrica, schotia-afra-var-afra

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Kuni bush

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo yo vhibvaho yo omiswaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Nyeletshedzo ya u renga: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Tshiko: SANBI Red List: Searsia undulata

Dune spinach

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: ṱhodzi ntswa dzi re na maṱari na tsinde.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Nyeletshedzo ya u renga: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Tetragonia decumbens

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Kinkelbossie

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: matavhi maṱuku, o bikwa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Nyeletshedzo ya u renga: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Khalaṅwaha ya u kaṇa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṇa henefho.

Tshiko: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maluvha a sa athu u vulea.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Nyeletshedzo ya u renga: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Khalaṅwaha ya u kaṇa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṇa henefho.

Tshiko: SANBI PlantZAfrica: Trachyandra ciliata

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Pigeonwood

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari maṱuku a muroho wo bikwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Nyeletshedzo ya u renga: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Trema orientalis

Wild garlic

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari na maluvha a ḽiwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Nyeletshedzo ya u renga: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu athu u khwaṱhisedzwa tshifhinga tshithihi tsha lushaka lwa henefho. Khalaṅwaha dza ndaedzi dzi re afho fhasi a si mbonelaphanḓa.

Ndaedzi fhedzi — tshiko tsha botaniki tsha vhubvaḓuvha ha South Africa — maluvha a ḽiwaho fhedzi: Phan–Lamb.

Tshiko: SANBI PlantZAfrica, tulbaghia-violacea

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Ginger

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: tsinde ntswa dzi aluwaho fhasi ha mavu; ginger yo omiswaho i rekhodiwa nga thungo.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Nyeletshedzo ya u renga: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Tshiko: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Tshiko: University of Pretoria author manuscript: Gatabazi et al. (2022)

Tshaka dza zwiliṅwa dza u sedzulusa henefho

Vhulungani dzina ḽa lushaka lune na nanga zwa vhukuma. Mitevhe mipfufhi ya ṱhoḓisiso i ṱoḓa khwaṱhisedzo ya khalaṅwaha ya henefho, maḓi na mavu.

Tomatoes

Lushaka lwo rekhodwaho: a lu athu u nangwa.

Kilima ya fhethu hafha a i athu u ṱalutshedzea, ngauralo a hu buliwi mutevhe mupfufhi wa henefho. Vulani garaṱa ya tshiliṅwa u vhona zwiko na u rekhoda zwe na nanga.

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.