BED CALENDAR

Growing plan: Nov 2026 - Oct 2027

Read across to see what occupies each bed. Each line is one planting, not a measured strip of land.

Bed reserved / growing Pick fresh food Finish a cover crop

Tray crops reserve the bed from their first readiness month. Follow the monthly jobs. An arrow means the crop continues after Oct 2027.

CROP / SHARE OF BED	Nov 2026	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Bed 1												6 m²
 **Onions** 61% area - sow Mar 2026	Pick											
 **Tomatoes** 1/4 area - sow Sep 2027												→
 **Sweetcorn** 1/2 area - sow Jan 2027						Pick						
 **Green beans** 1/4 area - sow Feb 2027						Pick						
Bed 2												6 m²
 **Lettuce** 62% area - sow Aug 2027												→
 **Carrots** 86% area - sow Apr 2027										Pick		
 **Green beans** 1/3 area - sow Sep 2027												→

BED CALENDAR - CONTINUED

Growing plan: Nov 2026 - Oct 2027

Read across to see what occupies each bed. Each line is one planting, not a measured strip of land.

Bed reserved / growing Pick fresh food Finish a cover crop

Tray crops reserve the bed from their first readiness month. Follow the monthly jobs. An arrow means the crop continues after Oct 2027.

CROP / SHARE OF BED	Nov 2026	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Bed 3												6 m²
Butternut Whole area - sow Mar 2027									Pick			
Carrots 1/3 area - sow Sep 2027												
Beetroot 1/3 area - sow Sep 2027												
Brinjal (eggplant) 1/3 area - sow Aug 2027												
Bed 4												6 m²
Tomatoes 75% area - sow Sep 2027												
Carrots 20% area - sow Apr 2027										Pick		
Sweet potato 1/4 area - sow Sep 2027												

BED CALENDAR - CONTINUED

Growing plan: Nov 2026 - Oct 2027

Read across to see what occupies each bed. Each line is one planting, not a measured strip of land.

Bed reserved / growing Pick fresh food Finish a cover crop

Tray crops reserve the bed from their first readiness month. Follow the monthly jobs. An arrow means the crop continues after Oct 2027.

CROP / SHARE OF BED	Nov 2026	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Bed 5												6 m²
Green beans 69% area - sow Feb 2027						Pick						
Tomatoes 1/4 area - sow Aug 2027												
Parsley 1/3 area - sow Sep 2027												
Cucumber 1/4 area - sow Sep 2027												
Bed 6												8 m²
Garlic 1/2 area - sow May 2026		Pick										
Broccoli 47% area - sow Feb 2027									Pick			
Sweet potato 46% area - sow Oct 2027												

BED CALENDAR - CONTINUED

Growing plan: Nov 2026 - Oct 2027

Read across to see what occupies each bed. Each line is one planting, not a measured strip of land.

Bed reserved / growing Pick fresh food Finish a cover crop

Tray crops reserve the bed from their first readiness month. Follow the monthly jobs. An arrow means the crop continues after Oct 2027.

CROP / SHARE OF BED	Nov 2026	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Bed 7												6 m²
Maize (mielies) 83% area - sow Oct 2027												▶
Plot 1												21 m²
Butternut Whole area - sow Sep 2027											▶	▶
Plot 2												21 m²
Groundnuts (peanuts) Whole area - sow Oct 2027												▶
Plot 3												21 m²
Maize (mielies) Whole area - sow Nov 2026	▶	▶	▶	▶	▶	Pick						
Plot 4												21 m²
Sweet potato Whole area - sow Sep 2027											▶	▶

UBHEJANE CRÈCHE

Nov 2026 - Oct 2027

Fruit, nuts and berries: solid marks are local dates. Outlined marks: references when established, not food promised this year.

This site (satellite climate records) - Hot area, no frost - summer rain (sowing calendar set for the sample farm)

CROP / FOOD SOURCE	Nov NOW	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Fresh veg												
Broccoli									●	●		
Carrots										●		
Garlic		●										
Green beans						●						
Onions	●											
Sweetcorn						●						
Stored veg												
Garlic			●	●	●							
Onions		●	●									
Staple crops												
Butternut									●	●		
Maize (mielies)						●						
Stored staples												
Butternut											●	●
Maize (mielies)							●	●	●	●	●	●

UBHEJANE CRÈCHE

Nov 2026 - Oct 2027

Fruit, nuts and berries: solid marks are local dates. Outlined marks: references when established, not food promised this year.

This site (satellite climate records) - Hot area, no frost - summer rain (sowing calendar set for the sample farm)

CROP / FOOD SOURCE	Nov NOW	Dec 2026	Jan 2027	Feb 2027	Mar 2027	Apr 2027	May 2027	Jun 2027	Jul 2027	Aug 2027	Sep 2027	Oct 2027
Food forest - fruit, nuts & berries and other foods												When established
Avocado Plan: Hass								●	●	●	●	●
Mango Regional season	●	●	●	●	●							
Moringa Months to confirm	Dates to confirm - see plant ages and food sources											
Num-num Months to confirm	Dates to confirm - see plant ages and food sources											
Animal products												

No confirmed production months shown. Check animal products and housing on your map in the app.

Growing space used	22%	20%	19%	23%	31%	36%	13%	13%	13%	13%	47%	75%

Outlined bands are planning references when established; they do not enter food totals or picking jobs. Blank months have no picking window marked. Recorded planting ages keep young plants out of early reference months. Check plant ages and Food sources to check in the app. Gold rows require each crop's storage conditions; check the full plan or app. Red space figures mean the plan needs more growing space than is mapped.

UBHEJANE CRÈCHE

Food sources to check

These are on your design. Their picking or production months still need checking on site.



Avocado

Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.



Mango

Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.



Moringa

Plants on map: 0 existing; 2 proposed. Picking months and fruit-bearing plants need local confirmation.



Num-num

Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.

UBHEJANE CRÈCHE

Why these months are marked

References describe established plants. Confirm local dates; new plants need time to produce.

Mango - Regional reference: Mango

South Africa (general; Limpopo + Mpumalanga = 96% of SA mango area). Regional industry season, not a Tommy Atkins-only harvest window. Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: National Agricultural Marketing Council (NAMC), South African Fruit Trade Flow Report, Issue 56, December 2024 (citing FPEF 2024), p. 3

Avocado - Assumed variety: Hass

South Africa - warm subtropical regions (Hass cultivar). Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: Directorate Communication: Cultivation of Avocados (South African Dept. of Agriculture / ARC-Institute for Tropical and Subtropical Crops), p. 5

UBHEJANE CRÈCHE

Before using stored food

The storage row applies only when these sourced conditions are met. Check the food before using it.

Garlic - sourced storage window: 3 months

Well-cured bulbs (dried several weeks, dark/dry/ventilated), then kept unrefrigerated in a cool, dry, dark, well-ventilated place (e.g. mesh bags).

Source guide

Onions - sourced storage window: 2 months

Field-dried (cured, necks fully dry) bulbs held at ambient temperature in well-ventilated trays/racks.

Source guide

Butternut - sourced storage window: 2 months

Mature hard-rind fruit, cured, clean, dry, undamaged, kept ventilated at 10-15°C/50-70% RH.

Source guide

Maize (mielies) - sourced storage window: 6 months

Shelled grain dried to 14% moisture, kept in a clean sealed container or silo with active weevil/larger-grain-borer control.

Source guide

Sweet potato - sourced storage window: 3 months

Only mature, undamaged roots; cured 30-32°C/85-90% RH for 4-7 days, then stored with minimal handling as cool as achievable (source optimum 13°C) at ~85-90% RH. The sweet potato weevil (Cylas spp.) is the main storage threat.

Source guide

Groundnuts (peanuts) - sourced storage window: 6 months

Unshelled (in-shell) pods dried to in-shell moisture ~10% or less, stored in gunny bags, mud bins, baskets or a ventilated dry room, protected from pests and damp.

Source guide

Not checked by an agronomist - a planning guide, not an instruction. Show it to your extension officer, and get a soil test before spending on soil.

Published optimum-condition maturity endpoints are planning slots, not promises. Before any successor is sown or transplanted, confirm the previous crop is finished and the bed is actually clear; crops may mature later.

How this plan was put together

From the plan suggested in November 2026. Anything changed by hand since is not described here.

WHAT TO WATCH

Check Bed 1 before sowing there: something is already growing in that bed and this whole-year plan may overlap it in its first months.

Rain-fed plan: a crop is only sown where every month it grows gets rain of at least half the estimated evaporation (the rainfall test of the FAO growing-period method, using this site's monthly rainfall; evaporation estimated from temperature by Thornthwaite's method, which can under-estimate it in dry, windy places). Average rain is not every year's rain - water in a dry spell if you can.

Pumpkin, Watermelon want more room to sprawl than a veg bed can give - grow them in a dedicated plot, along your property edges, or in a food forest area instead. Turn on "Grow big vines in a veg bed anyway" if you'd rather use one of your beds for them.

Hot-area sowing months for Groundnuts (peanuts), Maize (mielies) are not yet from a published local table - check those dates with your extension officer before relying on them.

Dry beans (sugar beans), Swiss chard (spinach), Kale, Cabbage, Onions, Potato, Garlic, Peas, Broad beans (fava beans), Broccoli, True spinach (English spinach), Cauliflower were left out: none of their sowing windows has enough rain through the whole crop without irrigation.

CHOICES THIS PLAN MADE

Butternut gets Plot 1 to itself - a staple plot is exactly the dedicated sprawling room it wants.

Staple plots each take one crop at full area - Plot 2: Groundnuts (peanuts) (sow Oct) - Plot 3: Maize (mielies) (sow Nov) - Plot 4: Sweet potato (sow Sep). No earlier crop is recorded on these plots, so rotation could not steer this first choice - record what grew there before, if you know it.

GROUND WITH NO NEW SOWING

7 growing areas have a stretch with no new sowing: Bed 1 (May-Sep), Bed 2 (Feb-Mar), Bed 3 (Feb), Bed 6 (May-Jun, Sep), Bed 7 (Apr-Sep), Plot 2 (Mar, May-Sep), Plot 3 (Mar, May-Oct). Well-documented crops have a sowing window for those stretches, but the months around them are already fully planted, or crop rotation rules out the families that would fit. Clearing a nearby month, or letting the ground rest, are both fine choices.

3 growing areas have a stretch with no new sowing: Bed 5 (May), Plot 1 (Mar-Aug), Plot 4 (Apr-Aug). No crop the catalog can plan properly - one with a checked growing time, spacing and yield - has a sowing window that reaches those stretches. Ask locally what else does; this plan is not proof that nothing can grow then.

HOW THIS PLAN WAS MADE

Sowing calendar set for the sample farm: Hot area, no frost - summer rain.

This plan follows a repeating whole-year cycle. It was chosen by comparing plans starting in each of the 12 months and keeping the one with the fewest months without a fresh harvest - not just what fits best from Nov.

These sowing months follow the KZN DARD hot-area calendar. Outside KwaZulu-Natal, confirm them with your local extension officer.

Broad automatic choices use crops with verified yield, duration and field spacing so their productivity comparison has a common evidence basis.

How many people you feed is not used to guess how much to plant. The planner tries the exact crops you chose; your mapped space, the sowing windows and crop rotation decide what fits.

Some sowing months were skipped for this site's climate - too hot, or not enough rain through the crop: Maize (mielies) (not Dec); Green beans (not Mar, Apr, May, Jun, Jul, Aug); Butternut (not Mar, Apr, May, Jul, Aug); Pumpkin (not Mar, Apr, May, Jul, Aug); Carrots (not Feb, Mar, Apr, May, Jun, Jul, Aug); Beetroot (not Feb, Mar, Apr, May, Jun, Jul, Aug); Tomatoes (not Jan, Feb, Mar, Jul, Dec); Peppers (not Jan, Feb, Mar, Jul, Oct, Nov, Dec); Chilli (not Jan, Feb, Mar, Jul, Oct, Nov, Dec); Sweet potato (not Aug, Nov, Dec, Jan, Feb, Mar); Lettuce (not Feb, Mar, Apr, May, Jun, Jul); Groundnuts (peanuts) (not Nov); Cucumber (not Mar, Apr, May, Jul, Aug); Watermelon (not Aug); Turnips (not Feb, Mar, Apr, May, Jun, Jul, Aug); Parsley (not Feb, Mar, Apr, May, Jun, Jul, Aug); Brinjal (eggplant) (not Jan, Feb, Mar, Apr, May, Jun, Jul); Gem squash (not Mar, Apr, May, Jul, Aug); Baby marrow (courgette) (not Mar, Apr, May, Jul, Aug); Spanspek (sweet melon) (not Mar, Apr, May, Jul, Aug); Sweetcorn (not Feb, Mar, Jul, Aug).


Some months here are warmer than the crop's ideal range in FAO ECOCROP, which rates growth as reduced then (not failed) - mulch and afternoon shade help: Carrots (Jan, Dec); Parsley (Sep, Oct, Nov, Dec).

UBHEJANE CRÈCHE


Fresh food, kept food and space

Known food options by month. Counted once per crop or product, not kilograms, portions or meals.


1. Fresh food (food kinds)



2. Fresh food + kept food (food kinds)



3. Used and free crop-growing space (m2)



The two food charts share a scale. A crop present fresh and kept still counts once; kept colours add other food kinds. Dated output records update fresh counts. Research season references and sources without local dates are outside food counts.

Pantry storage requires each crop's source conditions. Preserved stock is shown only for recorded months; no amount, recipe or safe shelf life is inferred.

No preserved stock is recorded. This is a record status, not proof that the farm has no stock.

Space is for 128 m2 of mapped beds and plots only; trees, animal housing and ponds are outside this area. Numbers above space bars show used m2.

Some beds or plots have overlapping or invalid planting shares. Used space is capped at their mapped area; resolve the layout before relying on this comparison.


UBHEJANE CRÈCHE

Production weights and missing records

Monthly records, crop-cycle benchmarks and food kept in stock are separate measures. Missing quantities are not zero.

Monthly production kg are not recorded. The crop-cycle graph below is a planning benchmark; it does not fill in missing monthly harvest weights.

Crop-cycle kg by first picking month (full-cycle kg)



Each bar groups the entire crop-cycle benchmark by the month picking starts. It is not the weight harvested in that month. The source does not supply a monthly picking curve; quantities are not divided across a picking window.

No monthly quantities are recorded for this dated year in kg, eggs, litres, items or kept-stock kg. Enter expected or actual amounts in the app; the printed monthly table appears once a quantity is recorded, including zero.

? means quantity not recorded, not zero. * marks a known stock subtotal with missing batch weights. Egg counts and litres use actual records where entered, otherwise expected records. They are never converted to kg. Stock kg are food already held; they are never added to new production.

Monthly quantities still needed: Onions, Mango, Garlic, Green beans, Sweetcorn, Maize (mielies), Avocado, Broccoli, Butternut, Carrots. Research season references do not establish a harvest weight.

Food sources with no dated quantity: Avocado, Mango, Moringa, Num-num.

Kept food without recorded stock kg: Onions, Garlic, Maize (mielies), Butternut. Catalogue pantry windows and preservation months supply no stock weight.

Annual tree yield schedules are incomplete for Nov 2026 - Oct 2027: Avocado, Mango, Moringa, Num-num. No annual tree kg subtotal is known.


Tree kg by age are annual farm assumptions. They are shown in the future harvest pages and are not spread across monthly production bars.

UBHEJANE CRÈCHE

As your plants grow

Vegetables and staples alongside fruit, nuts and berries. Ten-year projection; not a harvest promise.

Green = vegetable and staple crop cycles. Gold = your tree yield schedule. Striped ends show harvest that changes as plants age during the year. * = known subtotal only; missing yields are not zero.



UBHEJANE CRÈCHE

What this forecast assumes

Projection, not a harvest promise. Vegetables and staples repeat only saved recurring sowings; dated and existing crops happen once.

Vegetable and staple crop-cycle kilograms are counted in the year of first picking, not divided into monthly quantities. Carry-in harvests need actual records.

Vegetable and staple estimates hold bed area, sowings and crop benchmarks constant. Revise the plan if growing trees change shade, water or available bed space.

Tree ages run from planting on this site. Separate older and younger plants into age groups. Research first-crop ages are references, not guaranteed dates.

Tree kg are your farm or nursery assumptions per plant per year. Values hold until the next age you enter; no growth multiplier is invented. A range covers changes within the year. Review the schedule as plants age.

Check that the variety, local climate, pollination, water and care match your yield assumptions. Ages alone do not establish site suitability or survival.

Known subtotals leave out missing yields. Eggs, meat, milk, fish and honey are not included in these crop kg totals. Picking months and food-gap checks still need local confirmation.

UBHEJANE CRÈCHE

Fruit, nuts and berries by age

Each column covers twelve months. Species total = all mapped plants. Group rows give kg for that age group. ? = missing information.

PLANT / AGE GROUPS	Nov 2026	Nov 2027	Nov 2028	Nov 2029	Nov 2030	Nov 2031	Nov 2032	Nov 2033	Nov 2034	Nov 2035
 **Avocado** 1 plant on map	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg
 **Mango** 1 plant on map	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg
 **Moringa** 2 plants on map	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg
 **Num-num** 1 plant on map	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg	Species total ? kg

Avocado: 1 plant needs age groups

Mango: 1 plant needs age groups

Moringa: 2 plants need age groups

Num-num: 1 plant needs age groups

Not checked by an agronomist - a planning guide, not an instruction. Show it to your extension officer, and get a soil test before spending on soil.

UBHEJANE CRÈCHE

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Subtropical coast or Lowveld / Bushveld. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

UBHEJANE CRÈCHE

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Avocado

Assumed variety: Hass - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
					■	■	■	■	■		

0 existing; 1 proposed. Products: fruit.

First crop reference: 2-3 years. Bruyns Hill, Natal Midlands bioclimatic group 2; irrigated Hass on clonal Duke 7 rootstock, 7 x 7 m (200 trees/ha), planted 1982-1984. The paper reports Hass starting in its second season and Fuerte in its third. The two-to-three-year reference describes those trial cultivars, not every avocado or nursery starting age. Actual timing depends on variety and site care.

Established production reference: 6-8 years. In this irrigated, intensively managed Natal Midlands trial, Hass reached about 20 t/ha at age 6 and orchards approached maximum sustainable average yield by age 8. This is a trial-specific bearing reference; there is no national plateau or automatically interpolated age-yield curve.

Production reference: 100-112.5 kg/plant. Same-trial arithmetic at the initial 200 trees/ha: 20,000 kg/ha / 200 = 100 kg/tree at age 6; sustainable average 22,500 kg/ha / 200 = 112.5 kg/tree. At age 3, 10,000 kg/ha / 200 = 50 kg/tree equivalent. These are average-equivalent reference values for the named Hass/Duke 7 orchard conditions, not measured individual-tree yields or default production for a different density, cultivar, climate or management. Intermediate ages are not inferred.

Hass age benchmarks - irrigated Natal Midlands trial: Hass on Duke 7, initially 200 trees/ha at Bruyns Hill (Natal Midlands bioclimatic group 2), with intensive irrigation and nutrition. Age 3: 10,000 kg/ha; age 6: 20,000 kg/ha; sustainable average about 22,500 kg/ha. Same-study density permits equivalents of 50 kg/tree at age 3 and 100 kg/tree at age 6; there is no reference for every intermediate age. These orchard benchmarks are conditional on good resources, comparable climate and timely orchard management, including Phytophthora control and thinning; do not auto-fill farm quantities.

Keep roots well drained; water matters during flowering.

Buying advice: KZN sequence: earlier Fuerte (May-Nov), middle Hass (Aug-Dec), later Ryan (Nov-Feb). These seasons overlap. Buy labelled grafted trees; confirm rootstock and pollination partners.

Pollination: partly self fertile. Check the chosen cultivar's requirements with the nursery.

Assumed variety: Hass. South Africa - warm subtropical regions (Hass cultivar). Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: KZN DARD, Fruit and Nut Production, PDF pp. 6-7 (regional reference; confirm current nursery stock)

Source: SA Avocado Industry Overview - South African Avocado Growers' Association (SAAGA)

Source: Directorate Communication: Cultivation of Avocados (South African Dept. of Agriculture / ARC-Institute for Tropical and Subtropical Crops), p. 5

Source: Wolstenholme, Kaiser & Palmer, SAAGA Yearbook 1991, 14:15-18

Source: Wolstenholme, Kaiser & Palmer, 'Yield Potential of Intensively Managed Avocados in the Natal Midlands - The Early Bearing Years', South African Avocado Growers' Association Yearbook 1991, 14:15-18, p. 6

Source: Wolstenholme, Kaiser & Palmer, SAAGA Yearbook 1991, 14:15-18, p. 7

Source: Wolstenholme, Kaiser & Palmer (1991), SAAGA Yearbook 14:15-18; reflowed PDF, p. 6

UBHEJANE CRÈCHE

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Mango

Regional reference: Mango - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

0 existing; 1 proposed. Products: fruit.

First crop reference: 3-4 years. Grafted mango stock in the KZN guide generally begins bearing after 3-4 years. Context is frost-free tropical/warm subtropical production with low humidity at flowering, including dry Zululand, Valley Bushveld and Lowveld. The text distinguishes grafted stock from seed-raised Peach/Sabre; do not apply this age to an unrecorded seedling or infer yearly kg. Actual timing depends on variety and site care.

Grafted mango establishment - regional guide: Grafted stock generally first bears around 3-4 years and may remain productive 20 years or longer under the guide conditions. Seed-raised trees are a separate planting-material case. Frost, wet spring flowering conditions, cultivar and field establishment must be checked; longevity is not an age-yield curve.

Check frost, drainage and flowering-season disease pressure. Heat and sunburn sensitivity differ between cultivars.

Assumed planting variety: Tommy Atkins, subject to local flowering and frost checks. Confirm its local picking months; the calendar uses a labelled regional reference where sourced.

Buying advice: ARC identifies Tommy Atkins and Kent among more heat/sunburn-tolerant options; Sensation and Keitt are more susceptible. Confirm local harvest dates before choosing a succession.

Regional reference: Mango. South Africa (general; Limpopo + Mpumalanga = 96% of SA mango area). Regional industry season, not a Tommy Atkins-only harvest window. Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: ARC Climate-Smart Agriculture, module 6, printed pp. 239-242

Source: National Agricultural Marketing Council (NAMC), South African Fruit Trade Flow Report, Issue 56, December 2024 (citing FPEF 2024), p. 3

Source: KZN DARD, Fruit and Nut Production in KZN, Mango bearing age; printed p.16, p. 20

Moringa

0 existing; 2 proposed. Products: Leaves and green pods; keep their records separate.

First crop reference: 6-12 months. The 6-12-month reference concerns the first leaf harvest after planting. Flowers and pods are normally produced during the second year of growth in the same North West guide. Leaf readiness does not establish pod production or picking months. Actual timing depends on variety and site care.

Pod development, not a young green-pod picking calendar: This botanical description discusses pods that become brown at maturity. The separate harvesting section requires young, tender, green pods for food and supplies no calendar for that stage. March-April is retained as pod-development phenology, not a confirmed young-pod picking season or leaf calendar.

Choose a sunny, well-drained growing position, protect establishment and plan pruning for leaf harvest.

Buying advice: Use correctly identified local planting material. No locally validated named cultivar shortlist is sourced; this plan does not claim nitrogen fixation or medicinal effects.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: North West DARD: Moringa production for food security (2021)

Source: North West Department of Agriculture and Rural Development, Moringa (Moringa oleifera) production: A possible solution to South African food security challenges, p. 33

Source: North West Department of Agriculture and Rural Development, Moringa (Moringa oleifera) production: A possible solution to South African food security challenges, p. 7

UBHEJANE CRÈCHE

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Num-num

0 existing; 1 proposed. Products: Ripe fruit, fresh or preserved.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Coastal plant; allow for thorns near children and paths.

Buying advice: Ask for a locally suitable fruiting plant; no named food cultivar is established by this source.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: carissa-macrocarpa

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and ground.

Garlic

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Onions

Recorded variety: not chosen yet.

Research shortlist for similar climates: Texas Grano 502 PPR; Granex 33; Pyramid.

Source: Onion chapter, KZN DARD vegetable production guide, chapter 27, p. 1

Source: Onion chapter, KZN DARD vegetable production guide, chapter 27, p. 2

Source: Onion chapter, KZN DARD vegetable production guide, chapter 27, p. 3

Butternut

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Swiss chard (spinach)

Recorded variety: not chosen yet.

Research shortlist for similar climates: Fordhook Giant; Bright Lights.

Source: Fordhook Giant, Sakata product page

Source: Bright Lights, Starke Ayres product page

Tomatoes

Recorded variety: not chosen yet.

Research shortlist for similar climates: Rodade.

Source: Rodade, Sakata product page

Maize (mielies)

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

UBHEJANE CRÈCHE

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Crop varieties to check locally (continued)

Green beans

Recorded variety: not chosen yet.

Research shortlist for similar climates: Contender; Star 2000.

Source: Contender bush bean, Starke Ayres product page

Source: Star 2000 bush bean, Starke Ayres product page

Lettuce

Recorded variety: not chosen yet.

Research shortlist for similar climates: Aviram; Tropical Emperor; Summer Gold; Brigadier.

Source: LETTUCE INTRODUCTION, KZN DARD, p. 2

Source: Brigadier Lettuce Seeds, Starke Ayres product page

Source: LETTUCE INTRODUCTION, KZN DARD, p. 1

Broccoli

Recorded variety: not chosen yet.

Research shortlist for similar climates: Green Valiant; Liberty; Pirate; Star 2201; Viking.

Source: COLE CROPS, KZN DARD, p. 3

Source: COLE CROPS, KZN DARD (Table 14, Sowing times, reading the Broccoli row's three columns: Cold/Warm/Hot areas), p. 3

Sweet potato

Recorded variety: not chosen yet.

Research shortlist for similar climates: Blesbok; Khumo; Impala; Bosbok.

Source: History of Sweet Potato Breeding in South Africa: 1952-2024, ARC-VIMP Newsletter 19, March 2024, p. 1

Source: History of Sweet Potato Breeding in South Africa: 1952-2024, ARC-VIMP Newsletter 19, March 2024, p. 2

Source: History of Sweet Potato Breeding in South Africa: 1952-2024, ARC-VIMP Newsletter 19, March 2024, p. 3

Cabbage

Recorded variety: not chosen yet.

Research shortlist for similar climates: Conquistador.

Source: Conquistador F1 Hybrid Fresh Market Cabbage, Sakata product page

Carrots

Recorded variety: not chosen yet.

Research shortlist for similar climates: Chantenay Karoo; Kuroda.

Source: Chantenay Karoo, Sakata product page

Source: Kuroda, Sakata product page

Groundnuts (peanuts)

Recorded variety: not chosen yet.

Research shortlist for similar climates: Akwa; Anel; Kwarts.

Source: Groundnut Production: A Concise Guide, A.J. Cilliers, ARC Grain Crops Institute, p. 6

Source: South African Registered Groundnut Cultivars Overview, opot.co.za, August 2018, p. 6

Source: South African Registered Groundnut Cultivars Overview, opot.co.za, August 2018, p. 13

Source: Groundnut Production: A Concise Guide, A.J. Cilliers, ARC Grain Crops Institute, p. 1

UBHEJANE CRÈCHE

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Crop varieties to check locally (continued)

Beetroot

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Parsley

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Cucumber

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Brinjal (eggplant)

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

Sweetcorn

Recorded variety: not chosen yet.

The research has no matching regional shortlist. Ask a local grower or extension officer which named variety suits this season.

INPUTS

Seed and seedling buying schedule

Find the month, then the crop and bed. Check the seed quantity and sowing method before buying.

The buying rule

Source direct-sown or tray seed before the named sowing month; the source does not provide a universal procurement lead time. Ready-grown seedlings are listed at the start of the field-readiness window; buy them only when the bed and seedlings are ready. Living corms, slips, cloves and seed potatoes are listed close to planting. For a tray crop, choose either packet seed for the nursery or ready-grown seedlings - not both.

A published field seed rate gives a weight range for the mapped area. Confirm the sowing method with the supplier; no germination or loss allowance is added. For crops without a published seed rate, field spacing supports only an approximate FINAL stand. It does not prove a seed-buying quantity: use the packet's crop-specific sowing rate and germination guidance. Living-material ranges are approximate field positions from mapped area and published spacing, not guaranteed order quantities or loss allowances; supplier and crop-specific guidance may change what to purchase.

Source/check	Crop	Quantity	Method	For	Planting timeline
November 2026	Maize (mielies)	Packet rate needed	Direct sow; ~92-93 final positions	Plot 3	Sow Nov
January 2027	Sweetcorn	3.6-4.5 g seed	Published field seed rate; confirm sowing method	Bed 1	Sow Jan
February 2027	Green beans	Packet rate needed	Direct sow; ~134-313 final positions	Beds 1, 5	Sow Feb
March 2027	Broccoli	~11-21 seedlings positions	Ready seedlings; own seed before Feb nursery	Bed 6	Source own seed before Feb; nursery Feb; check/transplant Mar-May
	Butternut	Packet rate needed	Direct sow; ~10 final positions	Bed 3	Sow Mar
April 2027	Carrots	Packet rate needed	Direct sow; ~316-1 585 final positions	Beds 2, 4	Sow Apr

Seed and seedling buying schedule (continued)

Source/check	Crop	Quantity	Method	For	Planting timeline
September 2027	Carrots	Packet rate needed	Direct sow; ~99-500 final positions	Bed 3	Sow Sep
	Beetroot	Packet rate needed	Direct sow; ~95-200 final positions	Bed 3	Sow Sep
	Sweet potato	~56-101 slips positions	Living pieces; confirm loss allowance	Bed 4; Plot 4	Sow Sep
	Green beans	Packet rate needed	Direct sow; ~47-112 final positions	Bed 2	Sow Sep
	Butternut	Packet rate needed	Direct sow; ~35 final positions	Plot 1	Sow Sep
	Lettuce	~21-37 seedlings positions	Ready seedlings; own seed before Aug nursery	Bed 2	Source own seed before Aug; nursery Aug; check/transplant Sep-Nov
	Brinjal (eggplant)	~5-6 seedlings positions	Ready seedlings; own seed before Aug nursery	Bed 3	Source own seed before Aug; nursery Aug; check/transplant Sep-Nov
	Cucumber	Packet rate needed	Direct sow; ~2-4 final positions	Bed 5	Sow Sep
	Tomatoes	~2-6 seedlings positions	Ready seedlings; own seed before Aug nursery	Bed 5	Source own seed before Aug; nursery Aug; check/transplant Sep-Nov

Seed and seedling buying schedule (continued)

Source/check	Crop	Quantity	Method	For	Planting timeline
September 2027	Parsley	0.6 g seed	Published field seed rate; confirm sowing method	Bed 5	Sow Sep
October 2027	Groundnuts (peanuts)	Packet rate needed	Direct sow; ~622-1 400 final positions	Plot 2	Sow Oct
	Maize (mielies)	Packet rate needed	Direct sow; ~21-22 final positions	Bed 7	Sow Oct
	Tomatoes	~8-23 seedlings positions	Ready seedlings; own seed before Sep nursery	Beds 1, 4	Source own seed before Sep; nursery Sep; check/transplant Oct-Dec
	Sweet potato	~9-17 slips positions	Living pieces; confirm loss allowance	Bed 6	Sow Oct

Sweetcorn - check the sowing method

Published field rate: 12-15 kg/ha. The buying range uses the mapped area; no germination allowance is added.

Grown and eaten as a fresh vegetable, unlike field maize - the dossier's own food-group is fruiting vegetable, not staple grain, so it is kept there rather than lumped with maize despite the shared species. Tender to frost like maize.

Parsley - check the sowing method

Published field rate: 3 kg/ha. The buying range uses the mapped area; no germination allowance is added.

A biennial herb grown as an annual for its leaves. Apiaceae family, same as carrots - avoid repeating that family in the same bed too often.

ImbewuField

FIELD SHEET

November 2026 field sheet

Ubhejane Crèche - tick each job off as it is done.

7 jobs - 1 sowing / planting - 1 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before cultivating or adding amendments.	
☐	**Plot 3**	Prepare the ground for maize (mielies) (whole plot; 21 m²).	
Direct sowing and planting			
☐	**Plot 3**	Sow maize (mielies) (whole plot; 21 m²; rows 91 cm apart, 25 cm in the row, 5-10 cm deep).	
Harvest and record			
☐	**Bed 1**	Harvest onions (61% of the bed; 3.67 m²). Record kilograms and where it went.	
Care and establishment - dates to confirm			
		Undated reminders, not scheduled harvests. Check local care and readiness before establishment; record dates when known.	
☐	**Mapped plants**	Avocado: Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.	
☐	**Mapped plants**	Mango: Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.	
☐	**Mapped plants**	Moringa: Plants on map: 0 existing; 2 proposed. Picking months and fruit-bearing plants need local confirmation.	
☐	**Mapped plants**	Num-num: Plants on map: 0 existing; 1 proposed. Picking months and fruit-bearing plants need local confirmation.	

ImbewuField

FIELD SHEET

December 2026 field sheet

Ubhejane Crèche - tick each job off as it is done.

2 jobs - 0 sowing / planting - 1 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before adding amendments.	
☐	**Bed 1**	Prepare the ground for sweetcorn (half the bed; 3 m²).	
Harvest and record			
☐	**Bed 6**	Harvest garlic (half the bed; 4 m²). Record kilograms and where it went.	

ImbewuField

FIELD SHEET

January 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

3 jobs - 1 sowing / planting - 0 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before adding amendments.	
☐	**Bed 1**	Prepare the ground for green beans (a quarter of the bed; 1.5 m²).	
☐	**Bed 5**	Prepare the ground for green beans (69% of the bed; 4.13 m²).	
Direct sowing and planting			
☐	**Bed 1**	Sow sweetcorn (half the bed; 3 m²; rows 70-80 cm apart, 25-35 cm in the row, 2.5-4 cm deep).	

ImbewuField

FIELD SHEET

February 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

5 jobs - 3 sowing / planting - 0 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before adding amendments.	
☐	**Bed 3**	Prepare the ground for butternut (whole bed; 6 m²).	
☐	**Bed 6**	Prepare the ground for broccoli (47% of the bed; 3.75 m²).	
Nursery - raise seedlings			
☐	**Nursery**	Raise and label trays for broccoli (47% of the bed; 3.75 m²; sow 1.5-2 cm deep) for Bed 6. Start checking in Mar; transplant when ready, within the planning window through May.	
Direct sowing and planting			
☐	**Bed 1**	Sow green beans (a quarter of the bed; 1.5 m²; rows 45-60 cm apart, 4-7 cm in the row, 2-5 cm deep).	
☐	**Bed 5**	Sow green beans (69% of the bed; 4.13 m²; rows 45-60 cm apart, 4-7 cm in the row, 2-5 cm deep).	

March 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

4 jobs - 2 sowing / planting - 0 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before adding amendments.	
☐	**Bed 2**	Prepare the ground for carrots (86% of the bed; 5.13 m²).	
☐	**Bed 4**	Prepare the ground for carrots (20% of the bed; 1.21 m²).	
Direct sowing and planting			
☐	**Bed 3**	Sow butternut (whole bed; 6 m²; rows 150 cm apart, 40 cm in the row).	
☐	**Bed 6**	From this month, transplant broccoli (47% of the bed; 3.75 m²; rows 60-70 cm apart, 30-45 cm in the row) when seedlings and the bed are ready.	

April 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

5 jobs - 2 sowing / planting - 3 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Direct sowing and planting			
☐	**Bed 2**	Sow carrots (86% of the bed; 5.13 m²; rows 20-40 cm apart, 2-5 cm in the row, 1-2.5 cm deep).	
☐	**Bed 4**	Sow carrots (20% of the bed; 1.21 m²; rows 20-40 cm apart, 2-5 cm in the row, 1-2.5 cm deep).	
Harvest and record			
☐	**Bed 1**	Harvest sweetcorn (half the bed; 3 m²) and green beans (a quarter of the bed; 1.5 m²). Record kilograms and where it went.	
☐	**Bed 5**	Harvest green beans (69% of the bed; 4.13 m²). Record kilograms and where it went.	
☐	**Plot 3**	Harvest maize (mielies) (whole plot; 21 m²). Record kilograms and where it went.	

July 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

2 jobs - 0 sowing / planting - 2 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Harvest and record			
☐	**Bed 3**	Harvest butternut (whole bed; 6 m²). Record kilograms and where it went.	
☐	**Bed 6**	Harvest broccoli (47% of the bed; 3.75 m²). Record kilograms and where it went.	

ImbewuField

FIELD SHEET

August 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

11 jobs - 1 sowing / planting - 4 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before cultivating or adding amendments.	
☐	**Bed 2**	Prepare the ground for lettuce (62% of the bed; 3.7 m²) and green beans (a third of the bed; 2 m²).	
☐	**Bed 3**	Prepare the ground for carrots (a third of the bed; 2 m²), beetroot (a third of the bed; 2 m²) and brinjal (eggplant) (a third of the bed; 2 m²).	
☐	**Bed 4**	Prepare the ground for sweet potato (a quarter of the bed; 1.5 m²).	
☐	**Bed 5**	Prepare the ground for tomatoes (a quarter of the bed; 1.5 m²), parsley (a third of the bed; 2 m²) and cucumber (a quarter of the bed; 1.5 m²).	
☐	**Plot 1**	Prepare the ground for butternut (whole plot; 21 m²).	
☐	**Plot 4**	Prepare the ground for sweet potato (whole plot; 21 m²).	
Nursery - raise seedlings			
☐	**Nursery**	Raise and label trays for lettuce (62% of the bed; 3.7 m²; sow 1.5-2 cm deep) for Bed 2, brinjal (eggplant) (a third of the bed; 2 m²; sow 1.5-2 cm deep) for Bed 3 and tomatoes (a quarter of the bed; 1.5 m²; sow 1 cm deep) for Bed 5. Start checking in Sep; transplant when ready, within the planning window through Nov.	
Harvest and record			
☐	**Bed 2**	Harvest carrots (86% of the bed; 5.13 m²). Record kilograms and where it went.	
☐	**Bed 3**	Harvest butternut (whole bed; 6 m²). Record kilograms and where it went.	
☐	**Bed 4**	Harvest carrots (20% of the bed; 1.21 m²). Record kilograms and where it went.	
☐	**Bed 6**	Harvest broccoli (47% of the bed; 3.75 m²). Record kilograms and where it went.	

September 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

12 jobs - 7 sowing / planting - 0 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Prepare for the next planting			
		Where a bed follows another crop, confirm it is finished and cleared before preparing it. Assess soil and drainage; use a soil test or local advice before cultivating or adding amendments.	
☐	**Bed 1**	Prepare the ground for tomatoes (a quarter of the bed; 1.5 m²).	
☐	**Bed 4**	Prepare the ground for tomatoes (75% of the bed; 4.5 m²).	
☐	**Bed 6**	Prepare the ground for sweet potato (46% of the bed; 3.67 m²).	
☐	**Bed 7**	Prepare the ground for maize (mielies) (83% of the bed; 5 m²).	
☐	**Plot 2**	Prepare the ground for groundnuts (peanuts) (whole plot; 21 m²).	
Nursery - raise seedlings			
☐	**Nursery**	Raise and label trays for tomatoes (a quarter of the bed; 1.5 m²; sow 1 cm deep) for Bed 1 and tomatoes (75% of the bed; 4.5 m²; sow 1 cm deep) for Bed 4. Start checking in Oct; transplant when ready, within the planning window through Dec.	
Direct sowing and planting			
☐	**Bed 2**	Sow green beans (a third of the bed; 2 m²; rows 45-60 cm apart, 4-7 cm in the row, 2-5 cm deep). From this month, transplant lettuce (62% of the bed; 3.7 m²; rows 40-50 cm apart, 25-35 cm in the row) when seedlings and the bed are ready.	
☐	**Bed 3**	Sow carrots (a third of the bed; 2 m²; rows 20-40 cm apart, 2-5 cm in the row, 1-2.5 cm deep) and beetroot (a third of the bed; 2 m²; rows 20-30 cm apart, 5-7 cm in the row, 1.5-2 cm deep). From this month, transplant brinjal (eggplant) (a third of the bed; 2 m²; rows 70-80 cm apart, 50 cm in the row) when seedlings and the bed are ready.	
☐	**Bed 4**	Plant sweet potato slips (a quarter of the bed; 1.5 m²; rows 90-100 cm apart, 25-40 cm in the row).	

September 2027 field sheet (continued)

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Direct sowing and planting (continued)			
☐	**Bed 5**	Sow parsley (a third of the bed; 2 m²; rows 30 cm apart, 10 cm in the row, 1.5-2 cm deep) and cucumber (a quarter of the bed; 1.5 m²; rows 120-140 cm apart, 35-50 cm in the row, 2-3 cm deep). From this month, transplant tomatoes (a quarter of the bed; 1.5 m²; rows 90-120 cm apart, 30-60 cm in the row) when seedlings and the bed are ready.	
☐	**Plot 1**	Sow butternut (whole plot; 21 m²; rows 150 cm apart, 40 cm in the row).	
☐	**Plot 4**	Plant sweet potato slips (whole plot; 21 m²; rows 90-100 cm apart, 25-40 cm in the row).	

October 2027 field sheet

Ubhejane Crèche - tick each job off as it is done.

5 jobs - 5 sowing / planting - 0 harvest / recording

DONE	PLACE	WORK TO COMPLETE	DATE / NOTE
Direct sowing and planting			
☐	**Bed 1**	From this month, transplant tomatoes (a quarter of the bed; 1.5 m²; rows 90-120 cm apart, 30-60 cm in the row) when seedlings and the bed are ready.	
☐	**Bed 4**	From this month, transplant tomatoes (75% of the bed; 4.5 m²; rows 90-120 cm apart, 30-60 cm in the row) when seedlings and the bed are ready.	
☐	**Bed 6**	Plant sweet potato slips (46% of the bed; 3.67 m²; rows 90-100 cm apart, 25-40 cm in the row).	
☐	**Bed 7**	Sow maize (mielies) (83% of the bed; 5 m²; rows 91 cm apart, 25 cm in the row, 5-10 cm deep).	
☐	**Plot 2**	Sow groundnuts (peanuts) (whole plot; 21 m²; rows 30-45 cm apart, 5-7.5 cm in the row, 5-7.5 cm deep).	

PLAN VERSUS REALITY

Monthly harvest and field record

Write the month: _____ Team: _____

Write down what actually happened

Record the date, crop, bed and actual harvest weight. Write how much was used, kept, shared or lost. These records help you check the next plan.

Harvest record

Date	Crop / bed	Picked kg	Used kg	Stored kg	Sold/shared kg	Lost kg

PLAN VERSUS REALITY

Fruit, eggs, honey and other production

Write the month: _____ Team: _____

Use the unit you actually measured

Write the food and its source: tree, coop, hive or bed. Use kg for weighed food, eggs for counted eggs, or jars, bunches or items. Record jar size in the source column. Keep all amounts on one row in the same unit.

Date	Food / source	Collected	Unit	Used	Sold/shared	Lost

Weekly observation

Week	Rainfall	Irrigation	Pests / disease	What we observed	What we changed

Decision for next month

Observe. Adjust. Write it down.