SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Kriedoring

1 existing; 1 proposed. Products: food product not verified.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Exact identity; no confirmed household food part: The checked primary assessment establishes exact identity and South African distribution only. It does not establish edible parts, food-use preparation, picking dates or yield; claims for a different Lycium species are not transferred.

Record the exact identity and obtain local expert food-use advice before counting it as household food.

Buying advice: Confirm exact species with a qualified indigenous nursery. Do not substitute genus-level berry claims for identification.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI Red List: Lycium cinereum

Kriedoring

1 existing; 1 proposed. Products: berries - food use requires specialist review.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented berry-use caution: This explicit caution prevents routine household-food forecasting. Sporadic all-year fruiting is not promoted to an edible-food availability calendar.

Keep this as an identified hedge/inventory plant. Seek expert local food-use review before treating berries as household food.

Buying advice: Confirm the exact species and intended hedge use with a local indigenous nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Lycium ferocissimum

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Macadamia

Regional reference: Macadamia - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

1 existing; 1 proposed. Products: Nuts: record nut-in-shell separately from kernels.

First crop reference: 4-5 years. KZN establishment reference; this source also describes later bearing at higher elevations. Not a promise for all cultivars or every SA site. Actual timing depends on variety and site care.

Established production reference: 7-9 years. USDA FAS describes commercial maturity at 7-9 years while yields can continue increasing for a further 15-20 years. This does not establish a stable maximum-yield age.

Macadamia output at named ages - approximate WIS references: Year 4 Beaumont: approximately 0.6-0.8 kg/tree (the year-4 basis is not explicit); year 5: 2-10 kg/tree Wet In Shell (WIS); year 8: about 16 kg/tree; year 13-14: up to 35 kg/tree WIS. SAMAC says area differences are guaranteed. This is a source reference, not a validated local schedule; it cannot seed edible kernel kg or a universal interpolated curve.

Macadamia establishment depends on propagation: Grafted stock: first nuts around 3-5 years, depending on variety; seedlings usually 10-12 years and sometimes longer. SAMAC recommends removing flowers until year 3 during establishment. The existing KZN guide 4-5-year first-bearing reference remains qualified; nursery propagation must be recorded before applying it.

Match the cultivar to the microclimate; avoid hard frost and poorly drained ground.

Collect fallen nuts frequently, dehusk promptly and dry with clean ventilation. Keep poultry and manure away from harvesting nuts.

Buying advice: KZN coastal 90-300 m shortlist: Beaumont (695), Purvis (294), Makai (800), 660. Confirm site elevation and current pollination/maturity advice.

Pollination: partly self fertile. Check the chosen cultivar's requirements with the nursery.

Regional reference: Macadamia. South Africa (general - Mpumalanga, KwaZulu-Natal, Limpopo). Regional nut-in-shell season; individual cultivar maturity still needs checking. Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: KZN DARD, Fruit and Nut Production, PDF pp. 24-25 (regional reference; confirm current nursery stock)

Source: ARC Climate-Smart Agriculture, module 6, printed pp. 245-246

Source: SAMAC, Best Practices: Harvesting, Dehusking, Drying (2023), pp. 1-6

Source: USDA Foreign Agricultural Service (FAS), GAIN Report SF2025-0023, 'South Africa's Macadamia Industry Grows Despite Export Uncertainty', FAS Pretoria Post, 2025, p. 5

Source: SAMAC (South African Macadamia Growers' Association), 'What have we learned 2024' research summary newsletter

Source: KZN DARD, Fruit and Nut Production in KZN, Macadamia bearing age, p. 24

Source: SAMAC Information Hub, FAQs: potential yield at different ages

Source: SAMAC Information Hub, FAQs: time to produce nuts

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Apple

1 existing; 1 proposed. Products: fruit.

First crop reference: 3 years. KZN guide reference for light cropping at year 3, in temperate apple conditions best adapted there to Dry Highland Sourveld. Rootstock, cultivar and nursery starting age are not specified. A named low-chill cultivar still needs adequate local winter chilling and establishment; this is not a warm-coast guarantee. Actual timing depends on variety and site care.

Established production reference: 7 years. Full production at year 7 in this general KZN guide. No per-tree output at that age is stated, and rootstock, cultivar and management can change bearing. The four Langkloof budgets are separate regional planning assumptions, not measured confirmation of this generic age.

Apple year 3 - Langkloof enterprise-budget output: Langkloof, Little Karoo, sandy loam: budgeted year 3 product quantity 10,000 kg/ha. This is a commercial enterprise-budget assumption, not a field measurement. The budget gives no matched tree density, rootstock or cultivar; do not convert to kg/tree, apply it to an isolated young tree or interpolate other ages. Citation uses table excerpts: the total is in the Quantity column, and the apple product-income rows use Kg; the one-hectare land basis is stated in the heading.

Apple year 4-5 - Langkloof enterprise-budget output: Langkloof, Little Karoo, sandy loam: budgeted year 4-5 product quantity 25,000 kg/ha. This is a commercial enterprise-budget assumption, not a field measurement. The budget gives no matched tree density, rootstock or cultivar; do not convert to kg/tree, apply it to an isolated young tree or interpolate other ages. Citation uses table excerpts: the total is in the Quantity column, and the apple product-income rows use Kg; the one-hectare land basis is stated in the heading.

Apple year 6-7 - Langkloof enterprise-budget output: Langkloof, Little Karoo, sandy loam: budgeted year 6-7 product quantity 48,590 kg/ha. This is a commercial enterprise-budget assumption, not a field measurement. The budget gives no matched tree density, rootstock or cultivar; do not convert to kg/tree, apply it to an isolated young tree or interpolate other ages. Citation uses table excerpts: the total is in the Quantity column, and the apple product-income rows use Kg; the one-hectare land basis is stated in the heading.

Apple year 8 - Langkloof enterprise-budget output: Langkloof, Little Karoo, sandy loam: budgeted year 8 product quantity 64,000 kg/ha. This is a commercial enterprise-budget assumption, not a field measurement. The budget gives no matched tree density, rootstock or cultivar; do not convert to kg/tree, apply it to an isolated young tree or interpolate other ages. Citation uses table excerpts: the total is in the Quantity column, and the apple product-income rows use Kg; the one-hectare land basis is stated in the heading.

Check the actual cultivar, rootstock, winter chilling and compatible pollen source before buying. A warm or cold climate-zone label alone does not establish these conditions.

Buying advice: Afri-Star is an ARC-bred low-chill candidate with a stated 300-500 Infruitec Chilling Unit requirement; its December-January dates were obtained at Bien Donne, Simondium, Western Cape and need local checking elsewhere. Granny Smith has an Elgin first-pick reference in late March; SAPO names Golden Delicious or Hillieri as pollen partners. These are separate cultivar references, not a continuous picking season or a warm KZN recommendation. Confirm current nursery stock and local performance.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Bien Donne, Simondium, Western Cape - Afri-Star cultivar trial reference: Dec-Jan.

Reference only - Elgin, Western Cape - Granny Smith cultivar first pick: Mar.

Source: Culdevco / ARC: Afri-Star apple cultivar information (Bien Donne, Simondium trial context; checked 10 October 2026)

Source: SAPO Trust: Granny Smith apple cultivar information (Elgin first pick; checked 10 October 2026)

Source: Culdevco / ARC Infruitec-Nietvoorbij, Afri-Star apple cultivar information sheet

Source: SAPO Trust, Granny Smith apple cultivar information sheet

Source: KZN DARD, Fruit and Nut Production in KZN, Apple bearing age; printed p.28, p. 32

Source: Western Cape Department of Agriculture, Agricultural Economic Services, Apples (Yr 3), updated November 2024, p. 1

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Apple (continued)

Source: Western Cape Department of Agriculture, Agricultural Economic Services, Apples (Yr 4-5), updated November 2024, p. 1

Source: Western Cape Department of Agriculture, Agricultural Economic Services, Apples (Yr 6-7), updated November 2024, p. 1

Source: Western Cape Department of Agriculture, Agricultural Economic Services, Apples (Yr 8), updated November 2024, p. 1

Mango

Regional reference: Mango - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

1 existing; 1 proposed. Products: fruit.

First crop reference: 3-4 years. Grafted mango stock in the KZN guide generally begins bearing after 3-4 years. Context is frost-free tropical/warm subtropical production with low humidity at flowering, including dry Zululand, Valley Bushveld and Lowveld. The text distinguishes grafted stock from seed-raised Peach/Sabre; do not apply this age to an unrecorded seedling or infer yearly kg. Actual timing depends on variety and site care.

Grafted mango establishment - regional guide: Grafted stock generally first bears around 3-4 years and may remain productive 20 years or longer under the guide conditions. Seed-raised trees are a separate planting-material case. Frost, wet spring flowering conditions, cultivar and field establishment must be checked; longevity is not an age-yield curve.

Check frost, drainage and flowering-season disease pressure. Heat and sunburn sensitivity differ between cultivars.

Assumed planting variety: Tommy Atkins, subject to local flowering and frost checks. Confirm its local picking months; the calendar uses a labelled regional reference where sourced.

Buying advice: ARC identifies Tommy Atkins and Kent among more heat/sunburn-tolerant options; Sensation and Keitt are more susceptible. Confirm local harvest dates before choosing a succession.

Regional reference: Mango. South Africa (general; Limpopo + Mpumalanga = 96% of SA mango area). Regional industry season, not a Tommy Atkins-only harvest window. Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: ARC Climate-Smart Agriculture, module 6, printed pp. 239-242

Source: National Agricultural Marketing Council (NAMC), South African Fruit Trade Flow Report, Issue 56, December 2024 (citing FPEF 2024), p. 3

Source: KZN DARD, Fruit and Nut Production in KZN, Mango bearing age; printed p.16, p. 20

Ice plant

1 existing; 1 proposed. Products: leaves and fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Annual edible leaf and fruit crop: This is an annual, not a fruit tree. The account describes boiled leaves as a spinach substitute and pickling. It supplies no food harvest duration, kg/plant or exact leaf calendar.

Record planting batches and harvested leaves/fruit separately.

Match this frost-sensitive plant to dry saline/coastal conditions. Its accumulated salts can return to the soil as plants die.

Buying advice: Obtain exact-species seed or plants and assess the bed soil; no regional food cultivar or annual food yield is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Mesembryanthemum crystallinum

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Coastal red milkwood

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The source describes bright orange-red mature fruit. It does not give a picking calendar.

SANBI records this tree in east-coast dune forest from Eastern Cape through KwaZulu-Natal, with tolerance of salty sea spray.

Buying advice: Confirm Mimusops afra and lawful nursery provenance. Check current protected-tree collection requirements before taking plants or products from existing trees.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, mimusops-afra

Nkaka

1 existing; 1 proposed. Products: traditionally prepared leaves/fruit - specialist review needed.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible-part and safety review: SANBI reports culinary uses but also toxic constituents and poisoning concerns. A preparation report is not a complete safe household recipe.

Retain this climber in mapped inventory, but do not count it as normal household food until a qualified local specialist confirms identity, edible part and safe preparation.

SANBI records it across South Africa except Western Cape, in varied habitats.

Buying advice: Resolve food-use and preparation with an experienced local food-plant specialist before buying it as a household crop.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, momordica-balsamica

Waxberry

1 existing; 1 proposed. Products: berry fat - traditional processed product.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Historical processed fat, not a fresh-fruit forecast: Historical food use concerns the fatty berry coating, not a generic fresh sweet berry. No validated household extraction method, recovered food yield or storage life is established. Source coating/oil percentages are not assigned as processing conversion factors.

Treat edible-fat preparation as a specialist process requiring local validation.

Use well-drained coastal sand with sun or semi-shade; fruit production needs correctly sexed planting material.

Buying advice: Confirm the exact species and male/female stock with an indigenous nursery. Keep landscape use separate from an unverified household food process.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Morella cordifolia

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Moringa

1 existing; 1 proposed. Products: Leaves and green pods; keep their records separate.

First crop reference: 6-12 months. The 6-12-month reference concerns the first leaf harvest after planting. Flowers and pods are normally produced during the second year of growth in the same North West guide. Leaf readiness does not establish pod production or picking months. Actual timing depends on variety and site care.

Pod development, not a young green-pod picking calendar: This botanical description discusses pods that become brown at maturity. The separate harvesting section requires young, tender, green pods for food and supplies no calendar for that stage. March-April is retained as pod-development phenology, not a confirmed young-pod picking season or leaf calendar.

Choose a sunny, well-drained growing position, protect establishment and plan pruning for leaf harvest.

Buying advice: Use correctly identified local planting material. No locally validated named cultivar shortlist is sourced; this plan does not claim nitrogen fixation or medicinal effects.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: North West DARD: Moringa production for food security (2021)

Source: North West Department of Agriculture and Rural Development, Moringa (Moringa oleifera) production: A possible solution to South African food security challenges, p. 33

Source: North West Department of Agriculture and Rural Development, Moringa (Moringa oleifera) production: A possible solution to South African food security challenges, p. 7

Black mulberry

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Exact fruit identity and variability: The institutional species reference distinguishes fruiting forms and variable breeding systems. Ripe fruits are used as food; a black-fruited tree must not silently inherit this identity or the legal status of another Morus species.

South African taxonomic cultivation record: The museum lists M. nigra as cultivated in southern Africa and distinguishes naturalised Morus taxa. It gives no ripe-harvest months or quantified yield.

Fruit maturity and water context: Institutional management reference links dry roots to early fruit drop. It contains a non-local 15-year substantial-bearing statement; that is not a South African first-crop age and is not used as a forecast.

Confirm the exact mulberry species and that the chosen plant is a fruiting form. Fruit colour alone cannot identify it.

Watch water stress during fruit development and record the actual ripe picking months.

Buying advice: Ask for identified Morus nigra fruiting material with a locally observed season. Do not use a generic black-fruited mulberry or fruitless ornamental as proof of this product.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Orwa et al. (2009), World Agroforestry, Morus nigra profile, p. 1

Source: Iziko Museums, Biodiversity Explorer, Morus

Source: Orwa et al. (2009), World Agroforestry, Morus nigra profile, p. 4

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Banana

1 existing; 1 proposed. Products: fruit (bunch of bananas).

First crop reference: 13-20 months. KZN DARD explicitly states the planting-to-harvest and harvest-to-harvest range. Converted months to years (13/12 to 20/12). This is not a dated forecast or an annual all-month harvest window. Actual timing depends on variety and site care.

Annual plantation output: KZN plantation reference per year: conservative 20-25 t/ha, average 25-30 t/ha, good 40+ t/ha. This is annual area production; it is not the weight of one bunch or a measured age-yield curve for three plants in a banana circle.

Cultivar, planting and crop cycle: The KZN guide gives 13-20 months for both the first planting-to-harvest cycle and subsequent harvest-to-harvest cycles. It lists different commercial densities for Williams, Grand Nain and Chinese Cavendish. A banana circle counts as three planted bananas here; replacement followers are stages of those mats, not extra mapped trees or guaranteed year-round supply.

Warm, frost-free conditions and dependable water are needed. Keep a replacement follower after the fruiting stem is removed.

Assumed planting variety: Grand Nain for a warm site; use the wind/cold alternatives in the buying advice if those conditions apply. Local picking months remain unconfirmed.

Buying advice: Cooler KZN: Williams; warmer valleys: Grand Nain; windy coast: Chinese Cavendish. Buy clean planting material. These are climate choices, not early/middle/late seasons.

Edible bananas set fruit without pollination.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: KZN DARD, Bananas in KwaZulu-Natal, pp. 1-4

Source: ProMusa (Bioversity International-affiliated global banana knowledge platform), 'Domestication of the banana'

Source: KZN DARD, Fruit and Nut Production in KZN, Bearing age, p. 8

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Bananas: Yields, p. 9

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Bananas: Bearing age and Spacing, p. 8

Wild olive

1 existing; 1 proposed. Products: ripe wild-olive fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Wild olive is distinct from a named commercial olive cultivar. March-July describes fruit following flowers; no exact ripe-picking start/end is established.

SANBI records this slow-growing tree in summer and winter rainfall areas, with frost, drought and wind resistance.

Keep its roots away from walls, patios and pools; SANBI notes that they can be aggressive.

Buying advice: Ask for the wild-olive taxon in the catalogue, rather than substituting a commercial olive cultivar. Record actual local picking.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, olea-europaea-subsp-africana

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Olive

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Cultivar and product choice: Frantoio is described as an oil and cross-pollen candidate. Mission suits black table olives and oil; Kalamata black table, Manzanilla and Barouni green table. Climate fit and pollen needs depend on cultivar.

Regional orchard budget is not per-tree yield evidence: This is a historical one-hectare financial model; it does not provide matched per-tree kg or a measured biological maturity curve. No financial values or pesticide recipes are copied.

Match the cultivar to table olives or oil. Check drainage, irrigation and winter conditions; humid summer rainfall can bring disease problems.

Pick at the intended green, black or oil stage. Processing and safe kept-stock conditions must be recorded separately.

Buying advice: Mission is a source-backed black-table and oil candidate; Kalamata is for black table olives; Manzanilla and Barouni for green table olives; Frantoio for oil and compatible pollen. Confirm regional fit through SA Olive's registered nurseries.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Western Cape - cultivar and end-use dependent olive harvest: Feb-Jul.

Source: SA Olive Association, Olive Growing

Source: Western Cape Department of Agriculture, Oil Olives, Garden/Little Karoo budget, 13 March 2021, p. 1

Bietou

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Food is the ripe fruit. Main flowering in autumn/winter and fruit shortly afterwards do not establish named picking months.

SANBI describes a spreading pioneer shrub for sun and well-drained soil, including coastal and inland habitats. Leave room for its spread.

Buying advice: Confirm the exact species and local provenance; the checked source gives no named fruit cultivar.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, osteospermum-moniliferum

Jacket plum

1 existing; 1 proposed. Products: Ripe fruit pulp for fresh use or preserves.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Fruit appearance, not a dated ripe picking interval: SANBI describes green fruits that later split to expose edible red flesh. Its December-July fruit-production interval does not state when the fruit is ripe. Retain this as a general southern African phenology reference; local ripe picking months remain unknown.

Growth is slower in cold or dry conditions. Seed oil is not presented here as cooking oil.

Buying advice: Confirm local fruiting material; no named cultivar is sourced.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: pappea-capensis

Source: SANBI PlantZAfrica - Pappea capensis

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Crop varieties to check locally (continued)

ground.

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.