SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Avocado

Assumed variety: Hass - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

1 existing; 1 proposed. Products: fruit.

First crop reference: 2-3 years. Bruyns Hill, Natal Midlands bioclimatic group 2; irrigated Hass on clonal Duke 7 rootstock, 7 x 7 m (200 trees/ha), planted 1982-1984. The paper reports Hass starting in its second season and Fuerte in its third. The two-to-three-year reference describes those trial cultivars, not every avocado or nursery starting age. Actual timing depends on variety and site care.

Established production reference: 6-8 years. In this irrigated, intensively managed Natal Midlands trial, Hass reached about 20 t/ha at age 6 and orchards approached maximum sustainable average yield by age 8. This is a trial-specific bearing reference; there is no national plateau or automatically interpolated age-yield curve.

Production reference: 100-112.5 kg/plant. Same-trial arithmetic at the initial 200 trees/ha: 20,000 kg/ha / 200 = 100 kg/tree at age 6; sustainable average 22,500 kg/ha / 200 = 112.5 kg/tree. At age 3, 10,000 kg/ha / 200 = 50 kg/tree equivalent. These are average-equivalent reference values for the named Hass/Duke 7 orchard conditions, not measured individual-tree yields or default production for a different density, cultivar, climate or management. Intermediate ages are not inferred.

Hass age benchmarks - irrigated Natal Midlands trial: Hass on Duke 7, initially 200 trees/ha at Bruyns Hill (Natal Midlands bioclimatic group 2), with intensive irrigation and nutrition. Age 3: 10,000 kg/ha; age 6: 20,000 kg/ha; sustainable average about 22,500 kg/ha. Same-study density permits equivalents of 50 kg/tree at age 3 and 100 kg/tree at age 6; there is no reference for every intermediate age. These orchard benchmarks are conditional on good resources, comparable climate and timely orchard management, including Phytophthora control and thinning; do not auto-fill farm quantities.

Keep roots well drained; water matters during flowering.

Buying advice: KZN sequence: earlier Fuerte (May-Nov), middle Hass (Aug-Dec), later Ryan (Nov-Feb). These seasons overlap. Buy labelled grafted trees; confirm rootstock and pollination partners.

Pollination: partly self fertile. Check the chosen cultivar's requirements with the nursery.

Assumed variety: Hass. South Africa - warm subtropical regions (Hass cultivar). Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: KZN DARD, Fruit and Nut Production, PDF pp. 6-7 (regional reference; confirm current nursery stock)

Source: SA Avocado Industry Overview - South African Avocado Growers' Association (SAAGA)

Source: Directorate Communication: Cultivation of Avocados (South African Dept. of Agriculture / ARC-Institute for Tropical and Subtropical Crops), p. 5

Source: Wolstenholme, Kaiser & Palmer, SAAGA Yearbook 1991, 14:15-18

Source: Wolstenholme, Kaiser & Palmer, 'Yield Potential of Intensively Managed Avocados in the Natal Midlands - The Early Bearing Years', South African Avocado Growers' Association Yearbook 1991, 14:15-18, p. 6

Source: Wolstenholme, Kaiser & Palmer, SAAGA Yearbook 1991, 14:15-18, p. 7

Source: Wolstenholme, Kaiser & Palmer (1991), SAAGA Yearbook 14:15-18; reflowed PDF, p. 6

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Date palm

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Kakamas phenology is not a picking calendar: Table is fruit-set-to-maturation duration, not months of ripe harvest. Its separate table lists July-August flowering. Neither becomes a food-availability bar.

Southern Hemisphere reference, awaiting local confirmation: Broad hemispheric context, not a measured cultivar/site calendar. Kept outside the structured local harvest windows.

Western Cape Medjool ripening and spoilage trial: 2022 trial at Kleinplasie near Hermon concerns Medjool. Abstract treatment dates label ripening stages inconsistently; no harvest months are inferred. Full university handle returned HTTP 403 during checking.

Female palms need suitable male pollen. Record sex, cultivar and pollination management before forecasting fruit.

Check heat, water and rain during ripening. The Hermon trial found uneven ripening and spoilage, so regional success alone does not establish a local crop.

Buying advice: Medjool is documented in the Hermon study. Obtain known fruiting female material and compatible pollen advice; a seedling's sex and eventual fruit quality are not established by its appearance.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: FAO Date Palm Cultivation (2002), Chapter VIII

Source: FAO Date Palm Cultivation (2002), Chapter IV, Table 23

Source: FAO Date Palm Cultivation (2002), Chapter IX

Source: Peddie (2023), Stellenbosch MScAgric thesis, university-provided abstract in FAO AGRIS

Pistachio

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Producer's own cultivars: The same questionnaire confirms the nuts are grown on the producer's farm. Cultivar names are reproduced as written; spelling is not a taxonomy or catalogue change.

Prince Albert establishment and maturity expectation: An expectation for the newly established Platberg project, not a measured universal age curve. Its planted cultivars are Shufra, Ariyeh and Sirora; irrigated Karoo context retained.

Female trees bear nuts and need pollen from suitable male trees. The Prieska reference comes from a dry, irrigated Karoo farm.

Confirm male and female planting material, irrigation and local winter/summer conditions; a February producer reference does not validate a humid coast.

Buying advice: Discuss Shufrah and Ariyeh from the Prieska producer evidence, or Shufra, Ariyeh and Sirora from the Prince Albert project, with a specialist nursery. Select a compatible male and confirm local fit.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Orange River farm near Prieska, Northern Cape - Karoo Pistachios producer observation: Feb.

Source: Karoo Pistachios, first-person producer questionnaire on FoodHub

Source: Prince Albert Municipality, Pistachio Pioneers, 20 October 2025

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Spekboom

1 existing; 1 proposed. Products: edible leaves.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The food is the tart leaf; flowers do not define leaf picking months.

Primary food-processing research: The study analyses processed spekboom leaves. Its laboratory method and composition are not a farm harvest rate, storage life or household food-safety recipe.

Protect young plants from heavy browsing. SANBI reports warm rocky and semi-arid habitats; ground-level goat browsing can prevent vegetative spread and reduce regeneration.

Buying advice: Ask for correctly identified spekboom and record whether it is planted for food, browse or restoration. No measured household leaf yield is supplied.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, portulacaria-afra

Source: Mahlanza et al., University of Johannesburg research record, International Journal of Food Science and Technology (2026)

Apricot

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Solitaire winter and pollen context: Solitaire is listed as self-compatible. Its chilling units are named-cultivar requirements and are not a numeric range for every apricot.

Irrigated Robertson trial context: The field experiment used managed organic orchards and found trade-offs in mulch effects; it does not justify a universal yield improvement from mulch.

Keep each cultivar's picking reference separate. The Robertson example used summer irrigation.

Confirm winter chilling, water and the actual planting material before using a reference as a local plan.

Buying advice: Solitaire has a sourced November first-pick reference and named chilling context at Simondium; discuss its local fit with the nursery.

Buying advice: Imperial has a December reference at Robertson. This is a local trial candidate, not advice for frost-free coastal KZN.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Bien Donne, Simondium, Western Cape - Solitaire first pick: Nov.

Reference only - Tierhoek Organic Farm, Robertson, Breede River Valley, Western Cape - Imperial trial: Dec.

Source: Culdevco / ARC Infruitec-Nietvoorbij, Solitaire apricot cultivar sheet

Source: Birkhofer et al. (2019), Effects of Ground Cover Management in Organic Deciduous Fruit Orchards in South Africa, Table 1

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Sweet cherry

1 existing; 1 proposed. Products: Ripe sweet cherries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Production at age six: The guide reports 4-6 kg/tree in year 6 and warns that yields are difficult to predict because of pollination and fruit set. The separate good result at year 8-9 is 9 t/ha. Do not connect them into a made-up growth curve or convert the later hectare result to arbitrary tree spacing.

Chilling and pollen partners: Conventional commercial cherry reference: more than 1,000 Positive Utah Chill Units, not 1,000 chill hours. Low-chill candidates need their own verified requirements. Most cultivars need compatible cross-pollination; the named self-fertile choices differ. The September flowering reference is not a picking-month array.

Observed Ceres season start: Ceres growers reported the start of the 2025 cherry harvest in late October, seven to ten days earlier than the prior year. This is an observed season in a named region, not a fixed future duration, KZN calendar or all-variety season.

Check measured winter chilling, spring frost, drainage and rain near picking. The older KZN guide regarded commercial suitability there as still under trial; a broad climate-zone label does not replace current local advice.

Pollination depends on the selected cultivar. Compatible flowering and pollen sources are separate from early or late fruit ripening.

Buying advice: Regional guide candidates include Bing, Van, Rainier, Burlat and Vista, which generally need compatible pollen partners. It explicitly lists Lapins, Stella and Sweetheart as self-fertile. Minnie Royal, Royal Lee, Royal Dawn and Early Sweet are lower-chill candidates. Confirm current stock, exact rootstock, compatibility and measured local chilling before purchase; this is not a warm-coast recommendation.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Yield, p. 31

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Climate and Cultivars, p. 30

Source: Hortgro / Tru-Cape growers, Cherry season begins: Ideal conditions, 29 October 2025

Almond

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Texas Mission pollen, chill and weight context: Texas Mission is classified high chill and names Nonpareil as pollinator. Its 5 kg/tree production entry does not state tree age, management, trial region or whether weight is kernel or in-shell; it is not a mature annual food forecast.

Texas Mission named production entry: The Texas Mission table's Production (kg/tree) row gives 5 kg/tree. It does not state tree age, reporting period, trial region or whether this means kernels or nuts in shell. Retained as a cultivar source claim to investigate; numeric mature annual kg remains null.

Check the named cultivar's winter needs and compatible pollen before planting. Texas Mission is self-incompatible.

Record whether nut weights mean kernels or nuts in shell; these are different quantities.

Buying advice: Texas Mission is a documented SAPO cultivar; its sheet names Nonpareil as pollen partner and describes a high chilling requirement. Confirm local suitability with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - South African SAPO cultivar sheet; trial location not stated - Texas Mission first pick: Feb.

Source: SAPO Trust, Texas Mission almond cultivar information

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Japanese plum

1 existing; 1 proposed. Products: fruit.

First crop reference: 3 years. Stated for African Delight® specifically, an ARC-bred Japanese plum cultivar; 'third growth season' interpreted as year 3. Actual timing depends on variety and site care.

African Delight yield by orchard age - native hectare unit: African Delight: age 3, 12,000-20,000 kg/ha; age 4, 20,000-35,000 kg/ha; mature orchards, 35,000-50,000 kg/ha. Full-bearing age and matched planting density are not stated; do not convert using Ruby Crisp spacing or interpolate a per-tree curve.

Ruby Crisp achievable yield - age unspecified: Ruby Crisp achievable output is 30,000 kg/ha. Table 1 on PDF p.5 lists Ruby Crisp trial spacing 4 x 1.25 m and 2,000 trees/ha; the derived equivalent is 15 kg/tree, but the achievable claim is not explicitly the trial measurement or a mature-age benchmark. Keep native kg/ha and the qualification.

African Delight cultivar chilling reference: Named requirement only: 200-400 Infruitec Chilling Units. The source data is from Bien Donne, Simondium, Western Cape and may not represent other areas. Confirm local chill, rootstock and pollen compatibility before buying.

Ruby Crisp cultivar chilling reference: Named requirement only: 400-600 Infruitec Chilling Units. The source data is from Bien Donne, Simondium, Western Cape and may not represent other areas. Confirm local chill, rootstock and pollen compatibility before buying.

Flavour Star cultivar chilling reference: Named requirement only: 500-600 Infruitec Chilling Units. The source data is from Bien Donne, Simondium, Western Cape and may not represent other areas. Confirm local chill, rootstock and pollen compatibility before buying.

African Delight named production - age unspecified: Direct named production reference: 25 kg/tree at 4 × 1.5 m, Bien Donne, Simondium, Western Cape. Tree age and reporting period are not stated, so it is retained as an unaged cultivar reference, not generic annual mature output.

Match chilling and pollination partners. A second variety must flower compatibly, not merely ripen at a different time.

Buying advice: Candidate: African Delight (200-400 Infruitec units). Pollinators listed by ARC/Culdevco: Pioneer or African Rose. Late-February first-pick data is from the Western Cape reference site, not a promise for warm KZN.

Pollination: needs cross pollinator. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Western Cape (Boland / Klein Karoo / Robertson stone-fruit areas): Feb.

Reference only - Bien Donne, Simondium, Western Cape - Ruby Crisp first pick: Jan.

Reference only - Bien Donne, Simondium, Western Cape - Flavour Star first pick: Jan.

Source: Culdevco / ARC: African Delight cultivar sheet

Source: Culdevco (ARC/Deciduous Fruit Industry joint-venture cultivar licensing body) - African Delight® plum cultivar information; cross-referenced with Hortgro's 'TOP CULDEVCO LICENSED PLUM CULTIVARS' article

Source: Culdevco / ARC, Ruby Crisp cultivar information sheet

Source: Culdevco / ARC, Flavour Star cultivar information sheet

Source: Hortgro / Culdevco, 'TOP CULDEVCO LICENSED PLUM CULTIVARS' (2020) - African Delight® section

Source: Hortgro / Culdevco, 'TOP CULDEVCO LICENSED PLUM CULTIVARS' (2020) - African Delight® section, p. 4

Source: Hortgro / Culdevco (2020), African Delight section, p. 4

Source: Hortgro / Culdevco (2020), Ruby Crisp section and Table 1, p. 2

Source: Culdevco / ARC, African Delight cultivar information sheet

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Pear

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Rosemarie Select pollen partners: These are cultivar-specific compatible-pollen candidates, not a guarantee that any two pear trees pollinate one another.

Packham's Triumph pollen partners: SAPO describes chilling qualitatively as low and production as high; neither supplies a numeric age or kg/tree.

Record the named cultivar and actual first and last picking months. January and February references are alternatives, not a mixed-orchard calendar for every tree.

Buying advice: Rosemarie Select is a source-backed Western Cape candidate to discuss with the nursery. Ask about December or Packham's Triumph as compatible pollen partners.

Buying advice: For Packham's Triumph, ask about Clapp's Favourite, December or Starkrimson and check local winter conditions.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Ceres, Koue Bokkeveld, Western Cape - Rosemarie Select first pick: Jan.

Reference only - Wolseley, Western Cape - Rosemarie Select first pick: Jan.

Reference only - South African cultivar sheet; test location not stated - Packham's Triumph first pick: Feb.

Source: Culdevco / ARC Infruitec-Nietvoorbij, Rosemarie Select pear cultivar sheet

Source: SAPO Trust, Packham's Triumph pear cultivar sheet

Dogwood

1 existing; 1 proposed. Products: leaf brewing flavour - specialist use.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December-June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Buying advice: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Source: SANBI PlantZAfrica: Rhamnus prinoides

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Baboon grape

1 existing; 1 proposed. Products: ripe berries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The source gives a ripe-berry interval. This does not establish stored jam months or safe shelf life.

Provide climbing support and control its spread where space is limited. SANBI records coastal dunes and forest margins from Western Cape to KwaZulu-Natal and Mpumalanga.

Buying advice: Confirm Rhoicissus digitata; other wild-grape species have different descriptions and cannot supply its dates.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Western/Eastern Cape, KwaZulu-Natal and Mpumalanga botanical account - ripe berries: Mar-Aug.

Source: SANBI PlantZAfrica, rhoicissus-digitata

Bushman's grape

1 existing; 1 proposed. Products: edible berries; roots excluded.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Food is the berry; the same SANBI account reports poisonous tuberous roots. September-June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Buying advice: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, rhoicissus-tridentata

Rosemary

1 existing; 1 proposed. Products: fresh or dried leaves for seasoning.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Buying advice: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: DAFF: Rosemary production brochure (2012), p. 2

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Karoo boer-bean

1 existing; 1 proposed. Products: edible seeds.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Buying advice: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Little Karoo and drier Cape botanical account - edible seed production reference: May-Jun.

Source: SANBI PlantZAfrica, schotia-afra-var-afra

Karee

1 existing; 1 proposed. Products: ripe berries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: September-January describes fruit production; exact ripe-picking dates are not stated. Do not count leaves used as livestock fodder as household food.

SANBI records this frost- and drought-resistant shade tree in Karoo and Kalahari conditions.

Large amounts of its leaves can taint dairy milk, according to SANBI; this is not a ration recommendation.

Buying advice: Ask about male and female plants if fruit is intended. Natural distribution in the source excludes KwaZulu-Natal; broad hardiness is not a local fruit guarantee.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, searsia-lancea

Glossy currant

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Currant fruit food use and phenology: The primary account establishes edible ripe fruit and October-June fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Pick ripe fruit and record drying separately.

This shrub suits exposed coastal gardens and dune stabilisation; confirm local provenance.

Buying advice: Ask an indigenous nursery for exact species and local material. No regional high-yield food cultivar is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Searsia lucida

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and ground.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Crop varieties to check locally (continued)

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.