SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Kriedoring

1 existing; 1 proposed. Products: food product not verified.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Exact identity; no confirmed household food part: The checked primary assessment establishes exact identity and South African distribution only. It does not establish edible parts, food-use preparation, picking dates or yield; claims for a different Lycium species are not transferred.

Record the exact identity and obtain local expert food-use advice before counting it as household food.

Buying advice: Confirm exact species with a qualified indigenous nursery. Do not substitute genus-level berry claims for identification.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI Red List: Lycium cinereum

Kriedoring

1 existing; 1 proposed. Products: berries - food use requires specialist review.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented berry-use caution: This explicit caution prevents routine household-food forecasting. Sporadic all-year fruiting is not promoted to an edible-food availability calendar.

Keep this as an identified hedge/inventory plant. Seek expert local food-use review before treating berries as household food.

Buying advice: Confirm the exact species and intended hedge use with a local indigenous nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Lycium ferocissimum

Ice plant

1 existing; 1 proposed. Products: leaves and fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Annual edible leaf and fruit crop: This is an annual, not a fruit tree. The account describes boiled leaves as a spinach substitute and pickling. It supplies no food harvest duration, kg/plant or exact leaf calendar.

Record planting batches and harvested leaves/fruit separately.

Match this frost-sensitive plant to dry saline/coastal conditions. Its accumulated salts can return to the soil as plants die.

Buying advice: Obtain exact-species seed or plants and assess the bed soil; no regional food cultivar or annual food yield is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Mesembryanthemum crystallinum

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Nkaka

1 existing; 1 proposed. Products: traditionally prepared leaves/fruit - specialist review needed.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible-part and safety review: SANBI reports culinary uses but also toxic constituents and poisoning concerns. A preparation report is not a complete safe household recipe.

Retain this climber in mapped inventory, but do not count it as normal household food until a qualified local specialist confirms identity, edible part and safe preparation.

SANBI records it across South Africa except Western Cape, in varied habitats.

Buying advice: Resolve food-use and preparation with an experienced local food-plant specialist before buying it as a household crop.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, momordica-balsamica

Waxberry

1 existing; 1 proposed. Products: berry fat - traditional processed product.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Historical processed fat, not a fresh-fruit forecast: Historical food use concerns the fatty berry coating, not a generic fresh sweet berry. No validated household extraction method, recovered food yield or storage life is established. Source coating/oil percentages are not assigned as processing conversion factors.

Treat edible-fat preparation as a specialist process requiring local validation.

Use well-drained coastal sand with sun or semi-shade; fruit production needs correctly sexed planting material.

Buying advice: Confirm the exact species and male/female stock with an indigenous nursery. Keep landscape use separate from an unverified household food process.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Morella cordifolia

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Banana

1 existing; 1 proposed. Products: fruit (bunch of bananas).

First crop reference: 13-20 months. KZN DARD explicitly states the planting-to-harvest and harvest-to-harvest range. Converted months to years (13/12 to 20/12). This is not a dated forecast or an annual all-month harvest window. Actual timing depends on variety and site care.

Annual plantation output: KZN plantation reference per year: conservative 20-25 t/ha, average 25-30 t/ha, good 40+ t/ha. This is annual area production; it is not the weight of one bunch or a measured age-yield curve for three plants in a banana circle.

Cultivar, planting and crop cycle: The KZN guide gives 13-20 months for both the first planting-to-harvest cycle and subsequent harvest-to-harvest cycles. It lists different commercial densities for Williams, Grand Nain and Chinese Cavendish. A banana circle counts as three planted bananas here; replacement followers are stages of those mats, not extra mapped trees or guaranteed year-round supply.

Warm, frost-free conditions and dependable water are needed. Keep a replacement follower after the fruiting stem is removed.

Assumed planting variety: Grand Nain for a warm site; use the wind/cold alternatives in the buying advice if those conditions apply. Local picking months remain unconfirmed.

Buying advice: Cooler KZN: Williams; warmer valleys: Grand Nain; windy coast: Chinese Cavendish. Buy clean planting material. These are climate choices, not early/middle/late seasons.

Edible bananas set fruit without pollination.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: KZN DARD, Bananas in KwaZulu-Natal, pp. 1-4

Source: ProMusa (Bioversity International-affiliated global banana knowledge platform), 'Domestication of the banana'

Source: KZN DARD, Fruit and Nut Production in KZN, Bearing age, p. 8

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Bananas: Yields, p. 9

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Bananas: Bearing age and Spacing, p. 8

Wild olive

1 existing; 1 proposed. Products: ripe wild-olive fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Wild olive is distinct from a named commercial olive cultivar. March-July describes fruit following flowers; no exact ripe-picking start/end is established.

SANBI records this slow-growing tree in summer and winter rainfall areas, with frost, drought and wind resistance.

Keep its roots away from walls, patios and pools; SANBI notes that they can be aggressive.

Buying advice: Ask for the wild-olive taxon in the catalogue, rather than substituting a commercial olive cultivar. Record actual local picking.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, olea-europaea-subsp-africana

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Pistachio

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Producer's own cultivars: The same questionnaire confirms the nuts are grown on the producer's farm. Cultivar names are reproduced as written; spelling is not a taxonomy or catalogue change.

Prince Albert establishment and maturity expectation: An expectation for the newly established Platberg project, not a measured universal age curve. Its planted cultivars are Shufra, Ariyeh and Sirora; irrigated Karoo context retained.

Female trees bear nuts and need pollen from suitable male trees. The Prieska reference comes from a dry, irrigated Karoo farm.

Confirm male and female planting material, irrigation and local winter/summer conditions; a February producer reference does not validate a humid coast.

Buying advice: Discuss Shufrah and Ariyeh from the Prieska producer evidence, or Shufra, Ariyeh and Sirora from the Prince Albert project, with a specialist nursery. Select a compatible male and confirm local fit.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Orange River farm near Prieska, Northern Cape - Karoo Pistachios producer observation: Feb.

Source: Karoo Pistachios, first-person producer questionnaire on FoodHub

Source: Prince Albert Municipality, Pistachio Pioneers, 20 October 2025

Spekboom

1 existing; 1 proposed. Products: edible leaves.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The food is the tart leaf; flowers do not define leaf picking months.

Primary food-processing research: The study analyses processed spekboom leaves. Its laboratory method and composition are not a farm harvest rate, storage life or household food-safety recipe.

Protect young plants from heavy browsing. SANBI reports warm rocky and semi-arid habitats; ground-level goat browsing can prevent vegetative spread and reduce regeneration.

Buying advice: Ask for correctly identified spekboom and record whether it is planted for food, browse or restoration. No measured household leaf yield is supplied.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, portulacaria-afra

Source: Mahlanza et al., University of Johannesburg research record, International Journal of Food Science and Technology (2026)

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Sweet cherry

1 existing; 1 proposed. Products: Ripe sweet cherries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Production at age six: The guide reports 4-6 kg/tree in year 6 and warns that yields are difficult to predict because of pollination and fruit set. The separate good result at year 8-9 is 9 t/ha. Do not connect them into a made-up growth curve or convert the later hectare result to arbitrary tree spacing.

Chilling and pollen partners: Conventional commercial cherry reference: more than 1,000 Positive Utah Chill Units, not 1,000 chill hours. Low-chill candidates need their own verified requirements. Most cultivars need compatible cross-pollination; the named self-fertile choices differ. The September flowering reference is not a picking-month array.

Observed Ceres season start: Ceres growers reported the start of the 2025 cherry harvest in late October, seven to ten days earlier than the prior year. This is an observed season in a named region, not a fixed future duration, KZN calendar or all-variety season.

Check measured winter chilling, spring frost, drainage and rain near picking. The older KZN guide regarded commercial suitability there as still under trial; a broad climate-zone label does not replace current local advice.

Pollination depends on the selected cultivar. Compatible flowering and pollen sources are separate from early or late fruit ripening.

Buying advice: Regional guide candidates include Bing, Van, Rainier, Burlat and Vista, which generally need compatible pollen partners. It explicitly lists Lapins, Stella and Sweetheart as self-fertile. Minnie Royal, Royal Lee, Royal Dawn and Early Sweet are lower-chill candidates. Confirm current stock, exact rootstock, compatibility and measured local chilling before purchase; this is not a warm-coast recommendation.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Yield, p. 31

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Climate and Cultivars, p. 30

Source: Hortgro / Tru-Cape growers, Cherry season begins: Ideal conditions, 29 October 2025

Almond

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Texas Mission pollen, chill and weight context: Texas Mission is classified high chill and names Nonpareil as pollinator. Its 5 kg/tree production entry does not state tree age, management, trial region or whether weight is kernel or in-shell; it is not a mature annual food forecast.

Texas Mission named production entry: The Texas Mission table's Production (kg/tree) row gives 5 kg/tree. It does not state tree age, reporting period, trial region or whether this means kernels or nuts in shell. Retained as a cultivar source claim to investigate; numeric mature annual kg remains null.

Check the named cultivar's winter needs and compatible pollen before planting. Texas Mission is self-incompatible.

Record whether nut weights mean kernels or nuts in shell; these are different quantities.

Buying advice: Texas Mission is a documented SAPO cultivar; its sheet names Nonpareil as pollen partner and describes a high chilling requirement. Confirm local suitability with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - South African SAPO cultivar sheet; trial location not stated - Texas Mission first pick: Feb.

Source: SAPO Trust, Texas Mission almond cultivar information

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Crop varieties to check locally (continued)

ground.

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.