SYNTHETIC GRAPH SOURCE {COUNT} $&

U nga si xava kumbe ku sungula

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Ntlawa wa maxelo wa ndzavisiso: Ndhawu yo byala a yi si tiveka. Ntlawa lowu a hi xitsundzuxo lexi tiyisisiweke xa purasi leri.

Leswi u swi vikeleke hi purasi leri

Mati hi nguva yo oma

A swi rhekhodiwanga. Hetisa vulavisisi bya ndhawu u tiyisisa mati u nga si hlawula kungu leri lavaka switirhisiwa swo tala.

Rharhama epurasini leri

Leswi voniweke hi rharhama endhawini a swi rhekhodiwanga. Miringaniso ya n’hweti ya sathelayiti a yi koti ku tiyisisa masiku yo sungula ni yo hetelela ya rharhama.

Dyambu ni ku huma ka mati emisaveni

Dyambu: a swi rhekhodiwanga.

Ku huma ka mati: a swi rhekhodiwanga. Mbuyelo wa ku kambela misava ni swilaveko swa xibyariwa xin’wana ni xin’wana swa ha fanele ku kamberiwa.

Tirhekhodo ta lembe ro vika - ku nga ri mvhumbo

Tinhlayo ni tin’hweti ta lembe leri hundzeke ro vika swi hlamusela leswi vikiweke. A swi tshembisi swakudya eka kungu leri taka.

Original eggs {year} $&

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 240 eggs.

Ku tirhisiwa ekaya: 0 eggs. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: Khot, Maw. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Fish

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: a swi rhekhodiwanga.

Ku tirhisiwa ekaya: a swi rhekhodiwanga. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: a swi rhekhodiwanga. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Miroho yin’wana

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 12 kg.

Ku tirhisiwa ekaya: 7 kg. Swo xavisiwa: 3 kg.

Tin’hweti ta ntshovelo leti vikiweke: Sung, N'we. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Mihandzu, timanga ni swakudya swa ndzhavuko

Dogwood

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: nantswo wa matluka loko ku swekiwa swakunwa — ku tirhisiwa hi mutivi.

Nkambisiso wa ku tirhisiwa ku ri swakudya: tiyisisa leswaku i ximilana xihi, xiphemu lexi dyiwaka ni malunghiselelo lama faneleke ni mutsundzuxi wa ndhawu loyi a nga ni vutivi u nga si xi katsa eka swakudya swa le kaya. A xi katsiwanga eka tinhlayo ta swakudya swa ntolovelo ni mintirho yo tshovela.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Switsundzuxo swo xava: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Xihlovo: SANBI PlantZAfrica: Rhamnus prinoides

Bushman's grape

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: mihandzu leyitsongo leyi dyiwaka; timitsu a ti katsiwi.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Xiphemu lexi dyiwaka ni ku tirhisiwa endhawini: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Switsundzuxo swo xava: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica, rhoicissus-tridentata

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Mihandzu, timanga ni swakudya swa ndzhavuko (swi ya mahlweni)

Rosemary

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: matluka lamantshwa kumbe lama omisiweke yo nandzihisa swakudya.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Switsundzuxo swo xava: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: DAFF: Rosemary production brochure (2012), p. 2

Karoo boer-bean

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: mbewu leyi dyiwaka.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Xiphemu lexi dyiwaka ni ku tirhisiwa endhawini: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Switsundzuxo swo xava: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Nguva yo tshovela leyi languteriweke: a ku si tiyisisiwa nkarhi wun’we wa muxaka wa ndhawu. Tinguva ta swikombo leswi nga laha hansi a hi mvhumbo.

Xikombo ntsena — xihlovo xa botaniki xa Little Karoo ni Cape leyi omeke ku tlurisa — xikombo xa vuhumelerisi bya mbewu leyi dyiwaka: Mudy–Khot.

Xihlovo: SANBI PlantZAfrica, schotia-afra-var-afra

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Mihandzu, timanga ni swakudya swa ndzhavuko (swi ya mahlweni)

Kuni bush

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: mihandzu leyi vupfeke leyi omisiweke.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Switsundzuxo swo xava: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Xihlovo: SANBI Red List: Searsia undulata

Dune spinach

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: swiphemu swa le henhla leswintshwa leswi nga ni matluka ni swirhabyana.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Switsundzuxo swo xava: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica: Tetragonia decumbens

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Mihandzu, timanga ni swakudya swa ndzhavuko (swi ya mahlweni)

Kinkelbossie

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: swihluke leswintshwa, swi swekiwile.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Switsundzuxo swo xava: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: swiluva leswi nga si pfuleka.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Switsundzuxo swo xava: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica: Trachyandra ciliata

Pigeonwood

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: matluka lamantshwa ya matsavu lama swekiweke.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Switsundzuxo swo xava: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin’hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: SANBI PlantZAfrica: Trema orientalis

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Mihandzu, timanga ni swakudya swa ndzhavuko (swi ya mahlweni)

Wild garlic

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: matluka ni swiluva leswi dyiwaka.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

Xiphemu lexi dyiwaka ni ku tirhisiwa endhawini: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Switsundzuxo swo xava: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Nguva yo tshovela leyi languteriweke: a ku si tiyisisiwa nkarhi wun'we wa muxaka wa ndhawu. Tinguva ta swikombo leswi nga laha hansi a hi mvhumbo.

Xikombo ntsena — xihlovo xa botaniki xa le vuxa bya South Africa — mabilomu lama dyiwaka ntsena: Sung–Dziv.

Xihlovo: SANBI PlantZAfrica, tulbaghia-violacea

Ginger

Leswi nga kona: 1; leswi pulaniweke: 1. Swibveledzwa: swirhabyana leswintshwa swa le hansi ka misava; ginger leyi omisiweke yi rhekhodiwa yi ri yoxe.

Nkarhi wa kotulo yo sungula: a wu kumekanga eka xihlovo; u nga titshegi hi swimilana leswi swa ha ku byariwaka leswaku u kuma swakudya hi ku hatlisa.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Switsundzuxo swo xava: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Nguva yo tshovela leyi languteriweke: a ku na xihlovo xa tin'hweti leti vekiweke ta xifundzha. Rhekhoda masiku yo byala ni yo tshovela endhawini.

Xihlovo: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Xihlovo: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Xihlovo: University of Pretoria author manuscript: Gatabazi et al. (2022)

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Tinxaka ta swibyariwa to kambela endhawini

Hlayisa vito ra muxaka lowu u wu hlawulaka hakunene. Minxaxamelo yo koma ya ndzavisiso yi lava ku tiyisisiwa eka nguva ya ndhawu, mati ni misava.

Tomatoes

Muxaka lowu rhekhodiweke: a wu si hlawuriwa.

Maxelo ya ndhawu leyi a ma si tiveka, kutani a ku vuriwi leswaku ku ni nxaxamelo wo koma wa ndhawu. Pfula khadi ra xibyariwa u vona swihlovo ni ku rhekhoda xihlawuriwa xa wena.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.