SYNTHETIC GRAPH SOURCE {COUNT} $&

Ngaphambi kokuthenga noma ukuqala

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Iqembu lesimo sezulu locwaningo: Indawo yokulima ayikacaci. Leli qembu alisona isincomo esiqinisekisiwe saleli pulazi.

Lokho okubikile ngaleli pulazi

Amanzi ngesikhathi esomile

Akurekhodiwe. Qedela inhlolovo yendawo bese uqinisekisa amanzi ngaphambi kokukhetha uhlelo oludinga izinsiza eziningi.

Isithwathwa kuleli pulazi

Okubonwe ngesithwathwa endaweni akurekhodiwe. Izilinganiso zanyanga zonke zesathelayithi azikwazi ukuthola izinsuku zokuqala nezokugcina zesithwathwa.

Ilanga nokuphuma kwamanzi

Ilanga: akurekhodiwe.

Ukuphuma kwamanzi: akurekhodiwe. Kusadingeka ukuhlola imiphumela yokuhlolwa komhlabathi nezidingo zesitshalo ngasinye.

Amarekhodi onyaka wokubika - hhayi isibikezelo

Izamba nezinyanga zonyaka wokubika odlule zichaza lokho okwabikwa. Azithembisi ukudla ohlelweni oluzayo.

Original eggs {year} $&

Isamba sonyaka esibikwe umlimi sango-2025: 240 eggs.

Okusetshenziswa ekhaya: 0 eggs. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: Jun, Jul. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Fish

Isamba sonyaka esibikwe umlimi sango-2025: akurekhodiwe.

Okusetshenziswa ekhaya: akurekhodiwe. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: akurekhodiwe. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Eminye imifino

Isamba sonyaka esibikwe umlimi sango-2025: 12 kg.

Okusetshenziswa ekhaya: 7 kg. Okuthengisiwe: 3 kg.

Izinyanga zokuvuna ezibikiwe: Jan, Dis. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izithelo, amantongomane nokudla komdabu

Dogwood

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amaqabunga okunandisa lapho kuphiswa — ukusetshenziswa nguchwepheshe.

Ukubuyekezwa kokusetshenziswa njengokudla: qinisekisa ubunikazi, ingxenye edliwayo nokulungiswa okufanele nomeluleki wendawo oqeqeshiwe ngaphambi kokufaka lesi sitshalo ekudleni kwasekhaya. Asibalwa ezambeni zokudla okuvamile nasemisebenzini yokuvuna.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Confirm exact identity and the intended food preparation with an experienced local adviser.

The South African account describes stream-bank/forest-slope growth and frost resistance; its berries are described as bird food.

Iseluleko sokuthenga: Buy exact-species stock for the verified intended use. Do not treat medicinal or wildlife-fruit evidence as a fresh household-food forecast.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Umthombo: SANBI PlantZAfrica: Rhamnus prinoides

Bushman's grape

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amajikijolo adliwayo; izimpande azifakiwe.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Ingxenye edliwayo nokusetshenziswa endaweni: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Provide a trellis or other support. Young plants need water until established; SANBI describes later drought tolerance and sun to semi-shade use.

Iseluleko sokuthenga: Ask for Rhoicissus tridentata and record local ripe-fruit months. Do not use medicinal or animal root use as food evidence.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica, rhoicissus-tridentata

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izithelo, amantongomane nokudla komdabu (kuyaqhubeka)

Rosemary

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amaqabunga amasha noma omisiwe okunandisa ukudla.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Use well-drained soil and adequate water during establishment; avoid waterlogging and root rot.

Record fresh leaves, dried seasoning and essential oil as different products.

Iseluleko sokuthenga: Obtain identified healthy rosemary material and local growing advice; no regional culinary-leaf yield variety is verified.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: DAFF: Rosemary production brochure (2012), p. 2

Karoo boer-bean

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: imbewu edliwayo.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Ingxenye edliwayo nokusetshenziswa endaweni: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Its natural setting includes dry stream banks in the Little Karoo and drier Eastern and southern Western Cape.

The pods disperse seeds when dry. Record your local seed collection and use; this is not a measured food yield.

Iseluleko sokuthenga: Ask for Schotia afra var. afra, with botanical identity confirmed. The source's five-year shade claim is not a first-food age.

Isikhathi sokuvuna esilindelwe: alukho uhlobo olulodwa lwendawo olunefasitela eliqinisekisiwe. Izikhathi eziyizinkomba ezingezansi azisona isibikezelo.

Inkomba kuphela — umbhalo webhothani wase-Little Karoo nasezindaweni ezomile kakhulu zase-Cape — inkomba yokukhiqizwa kwembewu edliwayo: Mey–Jun.

Umthombo: SANBI PlantZAfrica, schotia-afra-var-afra

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izithelo, amantongomane nokudla komdabu (kuyaqhubeka)

Kuni bush

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: izithelo ezivuthiwe ezomisiwe.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Iseluleko sokuthenga: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Umthombo: SANBI Red List: Searsia undulata

Dune spinach

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: iziqongo ezincane ezinamaqabunga neziqu.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Iseluleko sokuthenga: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica: Tetragonia decumbens

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izithelo, amantongomane nokudla komdabu (kuyaqhubeka)

Kinkelbossie

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amahlumela amancane, aphekiwe.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Iseluleko sokuthenga: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: imiqumbe yezimbali.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Iseluleko sokuthenga: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica: Trachyandra ciliata

Pigeonwood

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amaqabunga amancane okuba yimifino ephekiwe.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Iseluleko sokuthenga: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: SANBI PlantZAfrica: Trema orientalis

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izithelo, amantongomane nokudla komdabu (kuyaqhubeka)

Wild garlic

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: amaqabunga nezimbali ezidliwayo.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

Ingxenye edliwayo nokusetshenziswa endaweni: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Iseluleko sokuthenga: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Isikhathi sokuvuna esilindelwe: alukho uhlobo olulodwa lwendawo olunefasitela eliqinisekisiwe. Izikhathi eziyizinkomba ezingezansi azisona isibikezelo.

Inkomba kuphela — umbhalo webhothani wasempumalanga ye-South Africa — izimbali ezidliwayo kuphela: Jan–Eph.

Umthombo: SANBI PlantZAfrica, tulbaghia-violacea

Ginger

Okukhona: 1; okuhleliwe: 1. Okukhiqizwayo: iziqu ezingaphansi komhlabathi ezisanda kuvunwa; ujinja omisiwe ubhalwa ngokwehlukana.

Isikhathi sesivuno sokuqala: asitholakali emthonjeni; ungathembeli ezitshalweni ezisanda kutshalwa ukuze uthole ukudla ngokushesha.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Iseluleko sokuthenga: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Isikhathi sokuvuna esilindelwe: azikho izinyanga ezimisiwe zesifunda ezinomthombo. Rekhoda izinsuku zokutshala nezokuvuna endaweni.

Umthombo: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Umthombo: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Umthombo: University of Pretoria author manuscript: Gatabazi et al. (2022)

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Izinhlobo zezitshalo okufanele zihlolwe endaweni

Gcina igama lohlobo olukhethayo ngempela. Uhlu olufushane locwaningo ludinga ukuqinisekiswa kwesikhathi sendawo, amanzi nomhlabathi.

Tomatoes

Uhlobo olurekhodiwe: akukakhethwa.

Isimo sezulu sale ndawo asikacaci, ngakho akukho hlu olufushane lwendawo olushiwo. Vula ikhadi lesitshalo ukuze ubone imithombo bese urekhoda okukhethile.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.