SYNTHETIC GRAPH SOURCE {COUNT} $&

U nga si xava kumbe ku sungula

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Ntlawa wa maxelo wa ndzavisiso: Ndhawu yo byala a yi si tiveka. Ntlawa lowu a hi xitsundzuxo lexi tiyisisiweke xa purasi leri.

Leswi u swi vikeleke hi purasi leri

Mati hi nguva yo oma

A swi rhekhodiwanga. Hetisa vulavisisi bya ndhawu u tiyisisa mati u nga si hlawula kungu leri lavaka switirhisiwa swo tala.

Rharhama epurasini leri

Leswi voniweke hi rharhama endhawini a swi rhekhodiwanga. Miringaniso ya n’hweti ya sathelayiti a yi koti ku tiyisisa masiku yo sungula ni yo hetelela ya rharhama.

Dyambu ni ku huma ka mati emisaveni

Dyambu: a swi rhekhodiwanga.

Ku huma ka mati: a swi rhekhodiwanga. Mbuyelo wa ku kambela misava ni swilaveko swa xibyariwa xin’wana ni xin’wana swa ha fanele ku kamberiwa.

Tirhekhodo ta lembe ro vika - ku nga ri mvhumbo

Tinhlayo ni tin’hweti ta lembe leri hundzeke ro vika swi hlamusela leswi vikiweke. A swi tshembisi swakudya eka kungu leri taka.

Original eggs {year} $&

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 240 eggs.

Ku tirhisiwa ekaya: 0 eggs. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: Khot, Maw. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Fish

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: a swi rhekhodiwanga.

Ku tirhisiwa ekaya: a swi rhekhodiwanga. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: a swi rhekhodiwanga. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Miroho yin’wana

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 12 kg.

Ku tirhisiwa ekaya: 7 kg. Swo xavisiwa: 3 kg.

Tin’hweti ta ntshovelo leti vikiweke: Sung, N'we. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso

Tinyoxi: Honeybee (managed hives)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: vulombe. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Ntshovelo wu titshege hi ntlawa wa tinyoxi, ku rhumbuka ni mpfula ya ndhawu. Kambela ku vupfa u siyela ntlawa swakudya; a ku na nguva ya vulombe leyi vekiweke ya tiko hinkwaro.

Nkarhi wa xikumiwa xo sungula: a wu na xihlovo eka bindzu leri.

Xikombo xa vuhumelerisi: 25 kg ya vulombe/hi yindlu ya tinyoxi (hive)/hi lembe. From a 2008 national postal survey of 112 SABIO-member beekeepers (22.4% response rate) managing an estimated 19,520 hives at 2,094 apiary sites; figure is self-reported and un-weighted. Table 7 (same doc, p.13-14) shows wide variance by region/operation-focus/size for the same year: e.g. 'All' region, honey-focused, all-size operations = 25.1-26.9 kg/hive/yr across 3 survey sites; Western Cape honey-focused = 23.6-27.9 kg/hive/yr; Western Cape pollination-focused, small operations = only 7.0-7.5 kg/hive/yr. The same document also quotes an earlier, more rigorously validated study: 'Their average reported yield is very low, at just 13.4 kilograms per hive' for Western Cape hives (Allsopp & Cherry 2004, cited p.2 of this working paper), and historical national bee-census data of 8.3 kg/hive/annum (1911-1988 average) rising to 24 kg/hive in the 1974/5 census (same doc, p.1). No single number can be called definitive; 25 kg/hive/yr is used as the document's own headline national estimate.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Veka tindlu ta tinyoxi ekule ni tindhawu to tlangela, nyika mati yo tenga naswona tirhisa ndhawu yo oma. Kombela mufuwi wa tinyoxi loyi a tsarisiweke ku kambela ku vekiwa loku hlayisekeke ka laha u nga si engetela yindlu ya tinyoxi.

Kambela ntlawa wun’wana ni wun’wana wa tinyoxi nkarhi na nkarhi ku kuma swikombiso swa vuvabyi bya tinyoxi leswi faneleke ku vikiwa ni ku tluleriwa hi Cape-bee (Capensis).

Veka ni ku sirhelela tindlu ta tinyoxi eka ku yiviwa ni ku onhiwa hi vomu, leswi nga xin’wana xa swiphiqo leswikulu leswi langutaneke ni vafuwi va tinyoxi va Afrika-Dzonga.

Register with the Department before beekeeping. Registration lasts 24 months from registration or renewal and must be renewed when it expires.

A beekeeper who fails to renew registration at expiry will be deregistered and face penalties under the Act.

Beekeeping activities and hired beekeeping services require a valid registration certificate. Check the certificate before using a beekeeper.

Every beehive must be clearly and legibly marked with the beekeeper's registration number.

Keep written records of colonies, apiary locations, diseases, disease-control measures and colony losses with their reasons.

Do not move honeybees across the Cape bee line, or keep Cape bees north of it, without written approval.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tinyoxi: Honeybee (managed hives) (swi ya mahlweni)

Report any suspected notifiable disease (American foulbrood is listed); officials can order an infected colony destroyed.

Xihlovo: FAO: Beekeeping in Africa, Site selection (general African apiary guidance)

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 6

Xihlovo: Gums & Bees: A roadmap for landowners in South Africa (South African National Biodiversity Institute, SANBI), p. 21

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(1) and 2(4), p. 4

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measure 2(5), p. 4

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(8) and 2(9), p. 5

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 5

Xihlovo: Survey of Beekeeping in South Africa, CSSR Working Paper No. 221 (Conradie & Nortje, 2008, UCT Centre for Social Science Research), p. 14

Xihlovo: FAO, Beekeeping in Africa, honey harvesting

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Nguni / crossbred beef cattle (communal and extensive veld)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: nyama. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Nkarhi wa xikumiwa xo sungula: a wu na xihlovo eka bindzu leri.

Xikombo xa vuhumelerisi: 114.5–121.2 kg ya ntiko loko ri tshikisiwa ku mama/hi rhole. Table 5 gives weaning weight (WW) across four agro-ecological zones for the same nine Nguni communal herds: arid 114.51 kg (lowest), semi-arid 119.98 kg, dry sub-humid 117.91 kg, humid 121.21 kg (highest). The range recorded here spans this one study's own arid-to-humid zone means; it is not merged with any other paper's weaning-weight figures (other SA studies report different ranges for different systems - e.g. research-station or stud herds - and were deliberately excluded to avoid mixing systems). 826 Nguni calves from nine herds that the paper itself says were 'kept under extensive grazing system' in Limpopo Province - a genuine communal/extensive veld system matching this enterprise.

Xikombo xa nkarhi wo humelerisa: malembe ya 6.4. 2354 days / 365.25 = 6.44 years. 'Failure time' is the length of productive herd life for the 580 uncensored (fully-observed) records in the combined dataset; the dissertation itself defines productive herd life as 'the time interval between first calving date to culling date'. Combined data from three extensively-grazed Nguni stud herds, animals born 1968-2015.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Masaka ya tihomu a ya fanelanga ku tala ku tlurisa: ku fanele ku va ni ndhawu leyi eneleke ehansi leswaku xifuwo xin’wana ni xin’wana xi etlela hi nkarhi wun’we.

Hi nkarhi wo fambisiwa, homu yin’wana ni yin’wana yi fanele ku va ni kwalomu ka 1.4 m^2 ya ndhawu ehansi leswaku ku layicha ku nga yi phusumeti ngopfu kumbe ku siya ndhawu yo anama ngopfu.

Tihomu a ti fanelanga ku fambisiwa nkarhi wo hundza 18 wa tiawara ti nga si rhelerisiwa ti nyikiwa swakudya, mati ni vutumbelo.

I nandzu ku dlaya tihomu hi ndlala, ku ti nyika swakudya leswi nga enelangiki, kumbe ku ti tsona mati kumbe swakudya.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, as part of the national bovine brucellosis control scheme under the Animal Diseases Act.

No animal may be slaughtered anywhere but a registered abattoir, with the exception below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Nguni / crossbred beef cattle (communal and extensive veld) (swi ya mahlweni)

Slaughter for the farmer's own household consumption, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.4.6.1, p. 7

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Xihlovo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Xihlovo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Xihlovo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Xihlovo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: Mpofu, Ginindza, Siwendu, Nephawe & Mtileni (2016). Effect of agro-ecological zone, season of birth and sex on pre-weaning performance of Nguni calves in Limpopo Province, South Africa. Tropical Animal Health and Production 49:187-194 (PMC5203849), Table 5

Xihlovo: Ngayo, M. (2017). Factors Affecting Productive Life and Fertility in Nguni Cows (MSc dissertation, University of the Free State), Table 5.1, p. 58

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Small-scale dairy cow (Jersey, Holstein or crossbred)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: masi. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Kunguhata hi ku ya hi ku tswariwa ka xiviri ka marhole/swimbutana ni ku humesa masi. Swifuwo leswi nga humesiki masi a swi nyiki masi.

Xikombo xa xikumiwa xo sungula: mavhiki ya 121.8–130.5. Table 1, AFC column, rows in the table's order Holstein-Friesland Reg./Grade, Jersey Reg./Grade, Ayrshire Reg./Grade: 28, 30, 25, 28, 29, 32 months. The Grade rows used here are Holstein-Friesland 30 and Jersey 28 (Grade Ayrshire 32 is not used: the enterprise is Jersey or Holstein type). 28 x 4.35 = 121.8 weeks; 30 x 4.35 = 130.5 weeks. Milk begins at first calving, so age at first calving is the time to first product.

Xikombo xa vuhumelerisi: 3840–5422 kg ya masi/hi homo ya xisati/hi nkarhi wo humesa masi. Table 1 ('Comparison of number of cows, production and fertility in the main dairy breeds in South Africa'), Milk column, Grade rows: Jersey 3840 kg and Holstein-Friesland 5422 kg per lactation (Registered rows: 4468 and 6426 kg). The document says these are National Milk Recording Scheme herds, which 'have been shown to be better managed than those outside the scheme, with their performance per lactation being 50 % greater than that of the national herd', so a smallholder herd should expect less.

Xikombo xa nkarhi wo humelerisa: malembe ya 1.9–2. Holstein 739.33 and Jersey 696.81 days of productive life (first calving to culling) in South African milk-recorded herds, 2000-2019: 696.81/365.25 = 1.91 and 739.33/365.25 = 2.02 years.

Xikombo xa vukhale/endlelo: In hotter, humid KZN, KZN DARD describes Nguni × Jersey dairy trials at Makhathini to combine milk potential with local adaptation. Hand milking with the calf present highlights competition between household milk and calf growth. Discuss adapted stock and feeding locally; commercial milk-recording totals do not forecast this system. Record calving, lactation stage and milk available after calf needs.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Masaka ya tihomu a ya fanelanga ku tala ku tlurisa: ku fanele ku va ni ndhawu leyi eneleke ehansi leswaku xifuwo xin’wana ni xin’wana xi etlela hi nkarhi wun’we.

Tihomu ta xisati a ti fanelanga ku layichiwa leswaku ti fambisiwa loko matswele ya tona ya pfimbile hi masi.

Tihomu a ti fanelanga ku fambisiwa nkarhi wo hundza 18 wa tiawara ti nga si rhelerisiwa ti nyikiwa swakudya, mati ni vutumbelo.

I nandzu ku dlaya homu ya masi hi ndlala, ku yi nyika swakudya leswi nga enelangiki, kumbe ku yi tsona mati kumbe swakudya.

Tihomu ta xisati emadyelweni kumbe etinkampeni leti pfulekeke ti lava ndzhuti: swivumbeko swa ndzhuti swi fanele ku nyika kwalomu ka 4 ku fika eka 5 m² ya ndzhuti hi homu.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Small-scale dairy cow (Jersey, Holstein or crossbred) (swi ya mahlweni)

Dairy farmers are legally required to test their herds for bovine TB, mainly because of export requirements, so most dairy cattle in South Africa are tested.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, under the national bovine brucellosis control scheme.

A cull cow may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.6.13, p. 9

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Xihlovo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 129

Xihlovo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Xihlovo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Xihlovo: Bovine tuberculosis / dairy herd testing study, PMC10369683

Xihlovo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: KZN DARD: Subsistence Dairying in KZN, Makhathini Research Station trials

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 10

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), Table 21.1, p. 122

Xihlovo: Gertenbach, W.D., Dairying in KwaZulu-Natal: Breeds of Dairy Cattle, Cedara Agricultural Development Institute, KZN Department of Agriculture and Rural Development, Table 1, p. 1

Xihlovo: Cow Longevity and Reasons and Risk Factors for Culling in South African Holstein and Jersey Dairy Herds, Animals 2025 (PMC12560872)

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Broilers (commercial)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: nyama. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: hlawula muxaka wa nyama ni kungu ra mintlawa ra muphakeri; tiyisisa ku kufumeta swinkukwana, swakudya, mati ni malunghiselelo yo tlhava.

Kunguhata swakudya leswi ringanisiweke, mati yo tenga, ndzhuti, ku nghena ka moya ni nsirhelelo eka swihadyana leswi dlayaka. Mphakelo wa matandza wu titshege hi tihuku leti tshikelaka, vukhale bya tona, rihanyo ni nhlayiso; hoko yi ri yoxe a yi nyiki matandza.

Xikombo xa xikumiwa xo sungula: mavhiki ya 4.6–4.7. 32 days / 7 = 4.57 weeks; 33 days / 7 = 4.71 weeks. This enterprise has a single grow-out cycle ending in slaughter, so 'time to first product' = age at slaughter. South Africa, commercial industry average.

Xikombo xa vuhumelerisi: 1.8–1.9 kg/hi xinyenyana loko xi tlhaviwa (masiku ya 32–33). National commercial broiler industry average, South Africa, 2022. A separate breed-trial table (Table 4b, same document, p.39) gives named-strain 'Cumulative FI (kg)' and 'Body weight (kg)' at exactly 35 days: Arbor Acres Plus 2.29kg, Cobb 500 2.52kg, Indian River 2.30kg, Ross 308 2.30kg (cumulative feed intake 3.21-3.64kg) - a slightly older age (35 vs 32-33 days) and higher weight than the industry-wide average quoted above; not merged into the primary figure since it is a different (breed-standard trial) dataset, but used below for the feed-intake figure.

Xikombo xa vukhale/endlelo: Newer South African commercial reference: SAPA 2024 reports average slaughter at 30–31 days with bird weight 1.8–1.85 kg. The primary range above remains the earlier 2022 reference at 32–33 days. These are commercial bird weights, not dressed meat kg or a village flock target. Record hatch or purchase age, feeding system, actual harvest date and weighed product basis.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Tihuku ta nyama leti tshamaka ehansi endlwini: ku nga vi ni ntiko wa tihuku wo hundza 40 kg hi mitara ya xikwere.

Swiciwana leswi nga ha ku tlhotlhoriwaka swi fanele ku nyikiwa swakudya ku nga si hundza 24 wa tiawara endzhaku ka ku tlhotlhoriwa.

Swiciwana leswi nga ha ku tlhotlhoriwaka swi fanele ku kota ku kuma mati ku nga si hundza 24 wa tiawara endzhaku ka ku tlhotlhoriwa, hi ku hatlisa ku tlurisa loko ku hisa.

Mati yo tenga ni swakudya leswi faneleke swi fanele ku va kona minkarhi hinkwayo, leswaku tihuku ti nga vi ni ndlala ni torha.

Tindlu ta tihuku ta nyama ti lava mpimo wa ku nghena ni ku famba ka moya wa kwalomu ka 5 wa timitara ta khubiki hi awara eka khilogiramu yin’wana ni yin’wana ya ntiko wa tihuku leti nga endlwini.

Eka ku fuwa tihuku ta nyama leti fambafambaka ehandle, nhlayo leyi tshamaka endzeni a yi fanelanga ku hundza 15 wa tihuku hi mitara ya xikwere ya ndhawu ya le hansi.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Broilers (commercial) (swi ya mahlweni)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), 2.2 Housing of broilers in Floor Systems, p. 6

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 11

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 12

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 7

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 8

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), 2.4 Free Range Broiler Production, p. 8

Xihlovo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Xihlovo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: SAPA Industry Profile 2024, section 7.9 Performance efficiency, p. 82

Xihlovo: SAPA Industry Profile 2022, section 6.9 Performance efficiency, p. 79

Xihlovo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Village chickens (indigenous, free-range)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: matandza. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Buying advice: Potchefstroom Koekoek is a dual-purpose option; Venda, Ovambo and Naked Neck are other ARC-conserved options. No province-specific winning breed is claimed.

Kunguhata swakudya leswi ringanisiweke, mati yo tenga, ndzhuti, ku nghena ka moya ni nsirhelelo eka swihadyana leswi dlayaka. Mphakelo wa matandza wu titshege hi tihuku leti tshikelaka, vukhale bya tona, rihanyo ni nhlayiso; hoko yi ri yoxe a yi nyiki matandza.

Hlengeleta matandza nkarhi ni nkarhi, hlayisa leswi andlariweke eswisakeni swi tengile naswona swi omile, khoma/hlayisa matandza hi vukheta leswaku ma tirhisiwa ekaya kumbe ku xavisiwa.

Xikombo xa xikumiwa xo sungula: mavhiki ya 15.7–27.8. Free-range village chickens, KwaZulu-Natal; dry-environment birds started laying earlier (15.74 weeks) than wet-environment birds (27.78 weeks), difference flagged significant ('*') in the source table. Wide range reflects genuine dry/wet-season environmental variation reported by the source, not measurement uncertainty.

Xikombo xa vuhumelerisi: 40.4–49.9 matandza/hi huku ya xisati/hi lembe. Free-range/scavenging village chickens, KwaZulu-Natal: dry (40.43) vs wet (49.91) environment, difference not significant ('NS'). A managed-system ARC Irene trial of named indigenous breeds (Grobbelaar, Sutherland & Molalakgotla 2010, Animal Genetic Resources 46: 25-32) reports far higher counts, now separately verified and retained in productionReferences; managed conditions cannot replace this village range.

Xikombo xa vukhale/endlelo: The ARC Irene trial compared Potchefstroom Koekoek, Venda, Ovambo and Naked Neck under commercial feed, water and controlled lighting. Mean eggs/hen in a 52-week cycle were 195.9, 153.7, 125.5 and 138.9 respectively; first-egg ages were 138.5, 139.0, 134.5 and 129.0 days from hatch, with no significant first-egg age difference among breeds. This is managed breed context, not the yield of scavenging hens in KZN. Match feeding and housing before using a benchmark.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Eka maendlelo ya le migangeni yo fuwa endhawini yo anama leyi pfulekeke, tihuku ti fanele ku tihlotela mati ni swakudya.

Tihuku ta le migangeni leti tihlotelaka swakudya ti kula hi ku nonoka ku tlurisa leti fuyiwaka hi mabindzu naswona ti le khombyeni lerikulu ra mavabyi, swivumbiwa leswi hanyaka hi ku tswonga miri wa tona (parasites) ni swiharhi leswi ti hlaselaka.

Eka ndzavisiso wa Limpopo na KwaZulu-Natal, 94.9% ya varimi a va nghenisa tihuku ta vona etindlwini nivusiku.

Leswi tihuku ta xisati leti tifambafambelaka a ti swi kuma hi ku tihlotela swakudya a swi enela ku hlayisa miri ni ku kula kambe swi nga eneli ku tshikela: phurotheyini eka tinguva hinkwato, calcium na phosphorus hi xixika.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Village chickens (indigenous, free-range) (swi ya mahlweni)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 471

Xihlovo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 468

Xihlovo: Malatji, Tsotetsi, van Marle-Koster & Muchadeyi (2016), 'A description of village chicken production systems and prevalence of gastrointestinal parasites: Case studies in Limpopo and KwaZulu-Natal provinces of South Africa', Onderstepoort Journal of Veterinary Research 83(1)

Xihlovo: Raphulu, Jansen van Rensburg & van Ryssen (2015), 'Nutrient content of crop contents of free ranging indigenous chickens raised under village conditions', South African Journal of Animal Science 45(2), p. 1

Xihlovo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Xihlovo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: Grobbelaar, Sutherland & Molalakgotla (2010), Egg production potentials of certain indigenous chicken breeds from South Africa: methods pp27–28, first-egg p29, Table 4 p31, p. 31

Xihlovo: Tenza, Mhlongo & Chimonyo (2023), 'Village chicken production and egg quality in dry and wet, resource-limited environments in KwaZulu-Natal, South Africa', South African Journal of Animal Science 53(6), p. 5

Xihlovo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Xihlovo: FAO: Small-scale poultry production, egg marketing

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Laying hens (commercial)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: matandza. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: leswaku u humelerisa matandza, kombela tihuku letitsongo leti tlhaveriweke leti nga ekusuhi ni ku sungula ku tshikela eka muphakeri wa tihuku to tshikela, ni tirhekhodo ta vukhale, swakudya ni ku tlhaveriwa.

Kunguhata swakudya leswi ringanisiweke, mati yo tenga, ndzhuti, ku nghena ka moya ni nsirhelelo eka swihadyana leswi dlayaka. Mphakelo wa matandza wu titshege hi tihuku leti tshikelaka, vukhale bya tona, rihanyo ni nhlayiso; hoko yi ri yoxe a yi nyiki matandza.

Hlengeleta matandza nkarhi ni nkarhi, hlayisa leswi andlariweke eswisakeni swi tengile naswona swi omile, khoma/hlayisa matandza hi vukheta leswaku ma tirhisiwa ekaya kumbe ku xavisiwa.

Xikombo xa nguva: With a constant 14–16 hour light day, commercial hens keep laying at a high rate all year. Xikombo lexi xa mafambiselo ya bindzu a xi vuli mphakelo wa matandza wa le kaya lowu ringanaka hi ntumbuluko.

Xikombo xa xikumiwa xo sungula: mavhiki ya 21. Vukhale bya huku ku sukela yi tlhotlhorhiwa; tiyisisa muxaka ni ku kurisiwa ni muphakeri. Tihuku leti nga ekusuhi ni ku tshikela ti sungula endzhaku eka ndzhendzeleko lowu.

Xikombo xa vuhumelerisi: 407.1–428.6 matandza/hi huku ya xisati leyi nghenisiweke endlwini (hen housed), ndzhendzeleko wo tshikela ku fika eka mavhiki ya 90. Table 4a's cumulative 'Eggs/hen housed' for five commercial strains; the same table gives 'Age at depop. weeks 90.0' for every strain. Commercial barn/cage average, not a free-range smallholder trial. Kept as the source gives it: a per-year rate would need a point-of-lay age from another document.

Xikombo xa vukhale/endlelo: SAPA 2024 commercial strain standards cumulative to 90 weeks: Dekalb Amberlink 429 eggs/hen housed, Dekalb Brown 428, Hy-Line Silver Brown 435, Hy-Line Brown 434 and Lohmann Brown Lite 413. These are full-cycle objectives, not eggs per calendar year or a smallholder forecast. Purchase age, actual onset of lay, surviving laying hens, feed, water and lighting need separate records.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Eka maendlelo yo fuwa lama pfumelelaka ku fambafamba ehandle, tihuku ta xisati ti fanele ku kota ku ya endhawini ya le handle ku ringana kwalomu ka 8 wa tiawara hi siku ku ri nhlikanhi.

Laha ku tirhisiwaka mabokisi ya swisaka swa xin’we-xin’we, ku fanele ku va ni kwalomu ka xisaka xin’we eka 8 wa tihuku ta xisati.

Tihuku ta xisati ti lava kwalomu ka 8 wa tiawara ta munyama lowu nga kavanyetiwiki eka ndzhendzeleko wun’wana ni wun’wana wa 24 wa tiawara leswaku ti wisa kahle.

Ku tshamisa ehansi endlwini swi lava swo andlala eka kwalomu ka xiphemu xin’we xa swinharhu (33%) xa ndhawu ya le hansi.

Ndhawu ya le handle yo fambafamba a yi fanelanga ku tala ku tlurisa: ku nga vi ni tihuku to hundza 5 hi mitara ya xikwere ya ndhawu yoleyo.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Laying hens (commercial) (swi ya mahlweni)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: SAPA Abridged Code of Practice: Commercial Layers, p. 1

Xihlovo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Xihlovo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: SAPA Industry Profile 2024, Table 4: commercial standards cumulative to 90 weeks, p. 39

Xihlovo: SAPA Layer Notes, Commercial Layers (training module, November 2013), p. 13

Xihlovo: SAPA Industry Profile 2022, Table 4a (Breed standards in laying hens), p. 39

Xihlovo: ARC Climate-Smart Agriculture Training Manual, Poultry production, p. 8

Xihlovo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Xihlovo: FAO: Small-scale poultry production, egg marketing

Xihlovo: SAPA Layer Notes, Commercial Layers (training module, November 2013)

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Masekwa: Meat duck (Pekin / Muscovy, backyard)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: nyama. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Xikombo xa xikumiwa xo sungula: mavhiki ya 7–12. P44 = Pekin (7 weeks to slaughter); MR71 = Muscovy (10 weeks for ducks/hens, 12 weeks for drakes). Treated as 'time to first product' = age to slaughter, since this is a single-cycle meat enterprise (no repeat product like eggs). Non-South-African trial by Warsaw researchers, published in SAJAS; journal location does not establish local applicability.

Xikombo xa vuhumelerisi: 2.3–2.9 kg ya ntiko wa xinyenyana lexi hanyaka/hi xinyenyana loko xi tlhaviwa. Non-SA (Guyana study reporting a citation to Steklenev 1990); no SA-specific absolute slaughter live-weight figure was found (see gaps). 2.93 kg = Pekin at 8-10 weeks, 2.29 kg = Muscovy at 8-10 weeks - matches this enterprise's two breeds. The same LRRD paper's own experiment (Table 2, intensive system, Guyana) measured live weight at slaughter (8 weeks) as: Pekin drakes 2.53 kg, Pekin hens 2.29 kg, Muscovy drakes 2.47 kg, Muscovy hens 1.51 kg - broadly consistent with, but not merged into, the cited range above since it is a separate dataset/table in the same source. No sourced eggs/duck/year figure was found for either breed (see gaps); the LRRD and SAJAS sources used here studied ducks reared only for meat.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

I nandzu ku tsona sekwa swakudya kumbe mati, kumbe ku xi hlayisa laha ku nga riki na ndhawu leyi eneleke, moya lowu eneleke, ku vonakala kumbe vutumbelo eka ku hisa ni xirhami.

Languta mitlhambi ku vona ku fa ka xitshuketa, tinsiva leti phaphamaleke, ku pfimba ka swiphemu swa nyama leswi lengalengaka ehansi ka nomo kumbe swa le nhlokweni, kumbe ku hunguteka ka ku tshikela hi xitshuketa - leswi i swikombiso swa mukhuhlwana wa tinyenyana naswona swi fanele ku vikiwa hi ku hatlisa.

Hlayisa masekwa ekule ni tinyenyana ta nhova ni mati kumbe swakudya leswi nga funengetiwangiki leswi ti kokaka, naswona lawula ku nghena ka vaendzi etindlwini ta masekwa, ku hunguta khombo ra mukhuhlwana wa tinyenyana.

Ducks are one of the domesticated animal species listed in the Meat Safety Act, so its rules on registered abattoirs and hygiene apply to them.

Ducks slaughtered to be sold must be slaughtered at a registered abattoir; slaughter for the farmer's own consumption is exempt but that meat may not be sold.

At the abattoir, ducks must be handled humanely during loading, transport, offloading, housing, immobilising and killing, under the Animals Protection Act.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Masekwa: Meat duck (Pekin / Muscovy, backyard) (swi ya mahlweni)

Ducks are birds/poultry, so an outbreak of avian influenza in a duck flock is a controlled animal disease under the Animal Diseases Act and must be reported to the state vet immediately.

Xihlovo: Animals Protection Act 71 of 1962 (consolidated text, s.2(1)(b)-(c)), p. 2

Xihlovo: Western Cape Department of Agriculture, Avian Influenza (Bird Flu) factsheet, p. 2

Xihlovo: Meat Safety Act 40 of 2000, Schedule 1 (Animals to which this Act applies), p. 16

Xihlovo: Meat Safety Act 40 of 2000, s.7, p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.9, p. 7

Xihlovo: Damaziak et al. (2014), Influence of housing system on the growth and histological structure of duck muscles, S. Afr. J. Anim. Sci. 44(2): Poland trial

Xihlovo: Livestock Research for Rural Development 18(11) 2006 - Solomon et al. (Guyana; not an SA figure), citing Steklenev 1990

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Xidziva: Tilapia (Mozambique tilapia, small pond or tank)

Miako ya 1 leyi nga kona; ya 2 leyi ringanyetiweke. Bindzu leri hlawuriweke: nhlampfi. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Xikombo xa xikumiwa xo sungula: mavhiki ya 43.5–52.2. 10 months x 4.35 = 43.5 weeks; 12 months x 4.35 = 52.2 weeks. This is Mozambique Tilapia (O. mossambicus) grow-out time to harvestable size; the source separately states Nile Tilapia (O. niloticus) reaches harvestable maturity faster, in 6-10 months (= 26.1-43.5 weeks) — not used here as it is a different species with a different legal status (see 'legal').

Xikombo xa vuhumelerisi: 800 g ya ntiko loko yi tshoveriwa/hi nhlampfi. Table 8-3 gives the Mozambique Tilapia's harvestable size as an approximate single figure ('+- 800g'), not a range, so [800, 800] is used rather than inventing a spread. The same table cell wraps '10-12' across a line break in the PDF ('10-\n12'), kept verbatim. The table's Nile Tilapia row gives the same approximate weight (+-800g) reached faster, in 6-10 months — not used here because this dossier is framed around the indigenous, legally unrestricted Mozambique Tilapia per task instruction; see 'legal' for why Nile Tilapia is treated separately. No SA source was found giving a feed-conversion-based or per-hectare yield figure specific to O. mossambicus that could be converted to this per-fish unit without stitching numbers across documents, so outputUnit follows the source's own unit (grams harvest weight per fish) rather than being converted.

Xikombo xa vukhale/endlelo: ARC makes tilapia suitability depend on water temperature: summer grow-out can be suitable anywhere in South Africa when water exceeds 20°C; growth is poor below 18°C and winter water below 13°C can kill farmed tilapia. The Highveld reference is a summer crop; tunnels can widen the range. Measure pond water and confirm stocking species and dates. Air climate or a mapped pond does not prove harvest months, fish numbers or food kg.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

The indigenous Mozambique Tilapia (O. mossambicus) is not a listed invasive species and may be farmed without an alien-species permit; the faster-growing Nile Tilapia (O. niloticus) is.

Nile Tilapia (Oreochromis niloticus) is a nationally listed invasive fish species: Category 1b inside National Parks, Provincial Reserves, Mountain Catchment Areas and Forestry Reserves, and Category 3 in all other areas.

Outside protected areas Nile tilapia is Category 3: it may remain in prescribed areas or provinces, but further propagation or trade is prohibited.

Nile Tilapia has been listed under the NEMBA Alien and Invasive Species Regulations as requiring a permit to produce.

Producing an alien or invasive species (such as Nile Tilapia) is prohibited in areas designated under the Protected Areas Act and the Fish Sanctuary Act; a permit is required to produce it elsewhere.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Xidziva: Tilapia (Mozambique tilapia, small pond or tank) (swi ya mahlweni)

Producing tilapia in South Africa requires a water use licence under the National Water Act.

Xihlovo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), Table 8-3 (Mozambique vs. Nile Tilapia), p. 69

Xihlovo: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), List 7 (National List of Invasive Fresh-water Fish Species), p. 32

Xihlovo: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), p. 3

Xihlovo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 69

Xihlovo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 74

Xihlovo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 75

Xihlovo: ARC Aquaculture Production App: General FAQ, Can I grow tilapia on the Highveld?

Tinxaka ta swibyariwa to kambela endhawini

Hlayisa vito ra muxaka lowu u wu hlawulaka hakunene. Minxaxamelo yo koma ya ndzavisiso yi lava ku tiyisisiwa eka nguva ya ndhawu, mati ni misava.

Tomatoes

Muxaka lowu rhekhodiweke: a wu si hlawuriwa.

Maxelo ya ndhawu leyi a ma si tiveka, kutani a ku vuriwi leswaku ku ni nxaxamelo wo koma wa ndhawu. Pfula khadi ra xibyariwa u vona swihlovo ni ku rhekhoda xihlawuriwa xa wena.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.