SYNTHETIC GRAPH SOURCE {COUNT} $&

Ngaphambi kokuthenga noma ukuqala

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Iqembu lesimo sezulu locwaningo: Indawo yokulima ayikacaci. Leli qembu alisona isincomo esiqinisekisiwe saleli pulazi.

Lokho okubikile ngaleli pulazi

Amanzi ngesikhathi esomile

Akurekhodiwe. Qedela inhlolovo yendawo bese uqinisekisa amanzi ngaphambi kokukhetha uhlelo oludinga izinsiza eziningi.

Isithwathwa kuleli pulazi

Okubonwe ngesithwathwa endaweni akurekhodiwe. Izilinganiso zanyanga zonke zesathelayithi azikwazi ukuthola izinsuku zokuqala nezokugcina zesithwathwa.

Ilanga nokuphuma kwamanzi

Ilanga: akurekhodiwe.

Ukuphuma kwamanzi: akurekhodiwe. Kusadingeka ukuhlola imiphumela yokuhlolwa komhlabathi nezidingo zesitshalo ngasinye.

Amarekhodi onyaka wokubika - hhayi isibikezelo

Izamba nezinyanga zonyaka wokubika odlule zichaza lokho okwabikwa. Azithembisi ukudla ohlelweni oluzayo.

Original eggs {year} $&

Isamba sonyaka esibikwe umlimi sango-2025: 240 eggs.

Okusetshenziswa ekhaya: 0 eggs. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: Jun, Jul. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Fish

Isamba sonyaka esibikwe umlimi sango-2025: akurekhodiwe.

Okusetshenziswa ekhaya: akurekhodiwe. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: akurekhodiwe. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Eminye imifino

Isamba sonyaka esibikwe umlimi sango-2025: 12 kg.

Okusetshenziswa ekhaya: 7 kg. Okuthengisiwe: 3 kg.

Izinyanga zokuvuna ezibikiwe: Jan, Dis. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela

Izinyosi: Honeybee (managed hives)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: uju. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Ukuvuna kuncike eqoqweni lezinyosi, ukuqhakaza nemvula endaweni. Hlola ukuvuthwa ushiyele iqoqo ukudla; asikho isikhathi sezwe lonke esinqunyiwe sokuvuna uju.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 25 kg woju/ngesidleke sezinyosi (hive)/ngonyaka. From a 2008 national postal survey of 112 SABIO-member beekeepers (22.4% response rate) managing an estimated 19,520 hives at 2,094 apiary sites; figure is self-reported and un-weighted. Table 7 (same doc, p.13-14) shows wide variance by region/operation-focus/size for the same year: e.g. 'All' region, honey-focused, all-size operations = 25.1-26.9 kg/hive/yr across 3 survey sites; Western Cape honey-focused = 23.6-27.9 kg/hive/yr; Western Cape pollination-focused, small operations = only 7.0-7.5 kg/hive/yr. The same document also quotes an earlier, more rigorously validated study: 'Their average reported yield is very low, at just 13.4 kilograms per hive' for Western Cape hives (Allsopp & Cherry 2004, cited p.2 of this working paper), and historical national bee-census data of 8.3 kg/hive/annum (1911-1988 average) rising to 24 kg/hive in the 1974/5 census (same doc, p.1). No single number can be called definitive; 25 kg/hive/yr is used as the document's own headline national estimate.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Beka izidleke zezinyosi kude nezindawo zokudlala, unikeze amanzi ahlanzekile futhi usebenzise indawo eyomile. Cela umfuyi wezinyosi obhalisiwe ahlole ukubekwa okuphephile kwendawo ngaphambi kokwengeza isidleke.

Hlola iqoqo ngalinye lezinyosi njalo ukuze uthole izimpawu zesifo sezinyosi okufanele sibikwe kanye nokutheleleka nge-Cape-bee (Capensis).

Beka futhi uvikele izidleke ekwebiweni nasekucekelweni phansi, okungenye yezinselelo ezinkulu kakhulu kubafuyi bezinyosi baseNingizimu Afrika.

Register with the Department before beekeeping. Registration lasts 24 months from registration or renewal and must be renewed when it expires.

A beekeeper who fails to renew registration at expiry will be deregistered and face penalties under the Act.

Beekeeping activities and hired beekeeping services require a valid registration certificate. Check the certificate before using a beekeeper.

Every beehive must be clearly and legibly marked with the beekeeper's registration number.

Keep written records of colonies, apiary locations, diseases, disease-control measures and colony losses with their reasons.

Do not move honeybees across the Cape bee line, or keep Cape bees north of it, without written approval.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinyosi: Honeybee (managed hives) (kuyaqhubeka)

Report any suspected notifiable disease (American foulbrood is listed); officials can order an infected colony destroyed.

Umthombo: FAO: Beekeeping in Africa, Site selection (general African apiary guidance)

Umthombo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 6

Umthombo: Gums & Bees: A roadmap for landowners in South Africa (South African National Biodiversity Institute, SANBI), p. 21

Umthombo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(1) and 2(4), p. 4

Umthombo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measure 2(5), p. 4

Umthombo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(8) and 2(9), p. 5

Umthombo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 5

Umthombo: Survey of Beekeeping in South Africa, CSSR Working Paper No. 221 (Conradie & Nortje, 2008, UCT Centre for Social Science Research), p. 14

Umthombo: FAO, Beekeeping in Africa, honey harvesting

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Nguni / crossbred beef cattle (communal and extensive veld)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 114.5–121.2 kg yesisindo ngesikhathi sokulumula/ngethole. Table 5 gives weaning weight (WW) across four agro-ecological zones for the same nine Nguni communal herds: arid 114.51 kg (lowest), semi-arid 119.98 kg, dry sub-humid 117.91 kg, humid 121.21 kg (highest). The range recorded here spans this one study's own arid-to-humid zone means; it is not merged with any other paper's weaning-weight figures (other SA studies report different ranges for different systems - e.g. research-station or stud herds - and were deliberately excluded to avoid mixing systems). 826 Nguni calves from nine herds that the paper itself says were 'kept under extensive grazing system' in Limpopo Province - a genuine communal/extensive veld system matching this enterprise.

Inkomba yesikhathi sokukhiqiza: iminyaka engu-6.4. 2354 days / 365.25 = 6.44 years. 'Failure time' is the length of productive herd life for the 580 uncensored (fully-observed) records in the combined dataset; the dissertation itself defines productive herd life as 'the time interval between first calving date to culling date'. Combined data from three extensively-grazed Nguni stud herds, animals born 1968-2015.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Izibaya zezinkomo akufanele zigcwale ngokweqile: kufanele kube nendawo eyanele yaphansi ukuze zonke izilwane zilale ngesikhathi esisodwa.

Ngesikhathi zithuthwa, inkomo ngayinye kufanele ibe nendawo yaphansi okungenani engu-1.4 m^2 ukuze ukulayishwa kungabi kuminyene kakhulu noma kube nesikhala esikhulu kakhulu.

Izinkomo akufanele zithuthwe amahora angaphezu kwangu-18 zingakehliswa zinikezwe ukudla, amanzi nendawo yokukhosela.

Kuyicala ukulambisa, ukunikeza ukudla okunganele, noma ukunqabela izinkomo amanzi noma ukudla.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, as part of the national bovine brucellosis control scheme under the Animal Diseases Act.

No animal may be slaughtered anywhere but a registered abattoir, with the exception below.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Nguni / crossbred beef cattle (communal and extensive veld) (kuyaqhubeka)

Slaughter for the farmer's own household consumption, or for cultural or religious purposes, is exempt from the abattoir rule.

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.4.6.1, p. 7

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Umthombo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Umthombo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: Mpofu, Ginindza, Siwendu, Nephawe & Mtileni (2016). Effect of agro-ecological zone, season of birth and sex on pre-weaning performance of Nguni calves in Limpopo Province, South Africa. Tropical Animal Health and Production 49:187-194 (PMC5203849), Table 5

Umthombo: Ngayo, M. (2017). Factors Affecting Productive Life and Fertility in Nguni Cows (MSc dissertation, University of the Free State), Table 5.1, p. 58

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Small-scale dairy cow (Jersey, Holstein or crossbred)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: ubisi. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Hlela ngokuzala kwangempela kwamathole/amazinyane nokuncelisa. Izilwane ezingancelisi azinikezi ubisi.

Inkomba yomkhiqizo wokuqala: amasonto angu-121.8–130.5. Table 1, AFC column, rows in the table's order Holstein-Friesland Reg./Grade, Jersey Reg./Grade, Ayrshire Reg./Grade: 28, 30, 25, 28, 29, 32 months. The Grade rows used here are Holstein-Friesland 30 and Jersey 28 (Grade Ayrshire 32 is not used: the enterprise is Jersey or Holstein type). 28 x 4.35 = 121.8 weeks; 30 x 4.35 = 130.5 weeks. Milk begins at first calving, so age at first calving is the time to first product.

Inkomba yokukhiqiza: 3840–5422 kg wobisi/ngenkomazi/ngesikhathi sokuncelisa. Table 1 ('Comparison of number of cows, production and fertility in the main dairy breeds in South Africa'), Milk column, Grade rows: Jersey 3840 kg and Holstein-Friesland 5422 kg per lactation (Registered rows: 4468 and 6426 kg). The document says these are National Milk Recording Scheme herds, which 'have been shown to be better managed than those outside the scheme, with their performance per lactation being 50 % greater than that of the national herd', so a smallholder herd should expect less.

Inkomba yesikhathi sokukhiqiza: iminyaka engu-1.9–2. Holstein 739.33 and Jersey 696.81 days of productive life (first calving to culling) in South African milk-recorded herds, 2000-2019: 696.81/365.25 = 1.91 and 739.33/365.25 = 2.02 years.

Inkomba yobudala/uhlelo: In hotter, humid KZN, KZN DARD describes Nguni × Jersey dairy trials at Makhathini to combine milk potential with local adaptation. Hand milking with the calf present highlights competition between household milk and calf growth. Discuss adapted stock and feeding locally; commercial milk-recording totals do not forecast this system. Record calving, lactation stage and milk available after calf needs.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Izibaya zezinkomo akufanele zigcwale ngokweqile: kufanele kube nendawo eyanele yaphansi ukuze zonke izilwane zilale ngesikhathi esisodwa.

Izinkomazi akufanele zilayishwe ukuze zithuthwe uma amabele azo egcwele ubisi avuvukala.

Izinkomo akufanele zithuthwe amahora angaphezu kwangu-18 zingakehliswa zinikezwe ukudla, amanzi nendawo yokukhosela.

Kuyicala ukulambisa, ukunikeza ukudla okunganele, noma ukunqabela inkomazi yobisi amanzi noma ukudla.

Izinkomazi ezisemadlelweni noma ezinkanjini ezivulekile zidinga umthunzi: izakhiwo zomthunzi kufanele zinikeze okungenani u-4 kuya ku-5 m^2 womthunzi ngenkomazi ngayinye.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Dairy farmers are legally required to test their herds for bovine TB, mainly because of export requirements, so most dairy cattle in South Africa are tested.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, under the national bovine brucellosis control scheme.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Small-scale dairy cow (Jersey, Holstein or crossbred) (kuyaqhubeka)

A cull cow may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.6.13, p. 9

Umthombo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Umthombo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 129

Umthombo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: Bovine tuberculosis / dairy herd testing study, PMC10369683

Umthombo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: KZN DARD: Subsistence Dairying in KZN, Makhathini Research Station trials

Umthombo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 10

Umthombo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), Table 21.1, p. 122

Umthombo: Gertenbach, W.D., Dairying in KwaZulu-Natal: Breeds of Dairy Cattle, Cedara Agricultural Development Institute, KZN Department of Agriculture and Rural Development, Table 1, p. 1

Umthombo: Cow Longevity and Reasons and Risk Factors for Culling in South African Holstein and Jersey Dairy Herds, Animals 2025 (PMC12560872)

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Broilers (commercial)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: khetha uhlobo lwenyama nohlelo lwamaqoqo lomhlinzeki; qinisekisa ukufudumeza amachwane, ukudla, amanzi nezinhlelo zokuhlaba.

Hlela ukudla okunomsoco olinganiselwe, amanzi ahlanzekile, umthunzi, umoya nokuvikelwa ezilwaneni ezidlayo. Ukutholakala kwamaqanda kuncike ezinkukhwini ezizalelayo, ubudala bazo, impilo nokunakekelwa; ihhoko lodwa alinikezi amaqanda.

Inkomba yomkhiqizo wokuqala: amasonto angu-4.6–4.7. 32 days / 7 = 4.57 weeks; 33 days / 7 = 4.71 weeks. This enterprise has a single grow-out cycle ending in slaughter, so 'time to first product' = age at slaughter. South Africa, commercial industry average.

Inkomba yokukhiqiza: 1.8–1.9 kg/ngenyoni ngesikhathi sokuhlatshwa (izinsuku ezingu-32–33). National commercial broiler industry average, South Africa, 2022. A separate breed-trial table (Table 4b, same document, p.39) gives named-strain 'Cumulative FI (kg)' and 'Body weight (kg)' at exactly 35 days: Arbor Acres Plus 2.29kg, Cobb 500 2.52kg, Indian River 2.30kg, Ross 308 2.30kg (cumulative feed intake 3.21-3.64kg) - a slightly older age (35 vs 32-33 days) and higher weight than the industry-wide average quoted above; not merged into the primary figure since it is a different (breed-standard trial) dataset, but used below for the feed-intake figure.

Inkomba yobudala/uhlelo: Newer South African commercial reference: SAPA 2024 reports average slaughter at 30–31 days with bird weight 1.8–1.85 kg. The primary range above remains the earlier 2022 reference at 32–33 days. These are commercial bird weights, not dressed meat kg or a village flock target. Record hatch or purchase age, feeding system, actual harvest date and weighed product basis.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Izinkukhu zenyama ezigcinwa phansi endlini: kungabi ngaphezu kuka-40 kg wesisindo sezinkukhu ngemitha eyisikwele ngayinye.

Amatshwele asanda kuchanyuselwa kufanele anikwe ukudla engakapheli amahora angu-24 echanyuselwe.

Amatshwele asanda kuchanyuselwa kufanele akwazi ukuthola amanzi engakapheli amahora angu-24 echanyuselwe, futhi ngokushesha kakhulu uma kushisa.

Amanzi ahlanzekile nokudla okufanele kuhlale kutholakala, ukuze izinkukhu zingalambi futhi zingomi.

Izindlu zezinkukhu zenyama zidinga ukungena nokugeleza komoya okungenani okungamamitha ayi-cubic angu-5 ngehora ngekhilogremu ngayinye yesisindo sezinkukhu ezisendlini.

Ekufuyeni izinkukhu zenyama ezihamba ngaphandle, inani eligcinwa ngaphakathi akufanele lidlule izinkukhu ezingu-15 ngemitha eyisikwele yendawo yaphansi.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Broilers (commercial) (kuyaqhubeka)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), 2.2 Housing of broilers in Floor Systems, p. 6

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), p. 11

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), p. 12

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), p. 7

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), p. 8

Umthombo: SAPA Code of Practice for Broiler Production (May 2012), 2.4 Free Range Broiler Production, p. 8

Umthombo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Umthombo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Umthombo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: SAPA Industry Profile 2024, section 7.9 Performance efficiency, p. 82

Umthombo: SAPA Industry Profile 2022, section 6.9 Performance efficiency, p. 79

Umthombo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Village chickens (indigenous, free-range)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: amaqanda. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Buying advice: Potchefstroom Koekoek is a dual-purpose option; Venda, Ovambo and Naked Neck are other ARC-conserved options. No province-specific winning breed is claimed.

Hlela ukudla okunomsoco olinganiselwe, amanzi ahlanzekile, umthunzi, umoya nokuvikelwa ezilwaneni ezidlayo. Ukutholakala kwamaqanda kuncike ezinkukhwini ezizalelayo, ubudala bazo, impilo nokunakekelwa; ihhoko lodwa alinikezi amaqanda.

Qoqa amaqanda njalo, ugcine okwandlalwe ezidlekeni kuhlanzekile futhi komile, uphathe/ugcine amaqanda ngokucophelela ukuze asetshenziswe ekhaya noma athengiswe.

Inkomba yomkhiqizo wokuqala: amasonto angu-15.7–27.8. Free-range village chickens, KwaZulu-Natal; dry-environment birds started laying earlier (15.74 weeks) than wet-environment birds (27.78 weeks), difference flagged significant ('*') in the source table. Wide range reflects genuine dry/wet-season environmental variation reported by the source, not measurement uncertainty.

Inkomba yokukhiqiza: 40.4–49.9 amaqanda/ngenkukhukazi/ngonyaka. Free-range/scavenging village chickens, KwaZulu-Natal: dry (40.43) vs wet (49.91) environment, difference not significant ('NS'). A managed-system ARC Irene trial of named indigenous breeds (Grobbelaar, Sutherland & Molalakgotla 2010, Animal Genetic Resources 46: 25-32) reports far higher counts, now separately verified and retained in productionReferences; managed conditions cannot replace this village range.

Inkomba yobudala/uhlelo: The ARC Irene trial compared Potchefstroom Koekoek, Venda, Ovambo and Naked Neck under commercial feed, water and controlled lighting. Mean eggs/hen in a 52-week cycle were 195.9, 153.7, 125.5 and 138.9 respectively; first-egg ages were 138.5, 139.0, 134.5 and 129.0 days from hatch, with no significant first-egg age difference among breeds. This is managed breed context, not the yield of scavenging hens in KZN. Match feeding and housing before using a benchmark.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Ezinhlelweni zasemakhaya zokufuya endaweni evulekile, izinkukhu kufanele zizifunele amanzi nokudla.

Izinkukhu zasemakhaya ezizifunela ukudla zikhula kancane kunezinkukhu ezifuywa ngokuhweba futhi zisengozini enkulu yezifo, izilokazane eziphila kuzo nezilwane ezizizingelayo.

Ocwaningweni lwaseLimpopo naseKwaZulu-Natal, abalimi abangu-94.9% babefaka izinkukhu zabo ezindlini ebusuku.

Lokho izikhukhukazi ezizihambelayo ezazikuthola ngokuzifunela ukudla kwakwanele ukugcina umzimba nokukhula, kodwa kunganele ukuzalela amaqanda: amaprotheni kuzo zonke izinkathi zonyaka, i-calcium ne-phosphorus ebusika.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Village chickens (indigenous, free-range) (kuyaqhubeka)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Umthombo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 471

Umthombo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 468

Umthombo: Malatji, Tsotetsi, van Marle-Koster & Muchadeyi (2016), 'A description of village chicken production systems and prevalence of gastrointestinal parasites: Case studies in Limpopo and KwaZulu-Natal provinces of South Africa', Onderstepoort Journal of Veterinary Research 83(1)

Umthombo: Raphulu, Jansen van Rensburg & van Ryssen (2015), 'Nutrient content of crop contents of free ranging indigenous chickens raised under village conditions', South African Journal of Animal Science 45(2), p. 1

Umthombo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Umthombo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Umthombo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: Grobbelaar, Sutherland & Molalakgotla (2010), Egg production potentials of certain indigenous chicken breeds from South Africa: methods pp27–28, first-egg p29, Table 4 p31, p. 31

Umthombo: Tenza, Mhlongo & Chimonyo (2023), 'Village chicken production and egg quality in dry and wet, resource-limited environments in KwaZulu-Natal, South Africa', South African Journal of Animal Science 53(6), p. 5

Umthombo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Umthombo: FAO: Small-scale poultry production, egg marketing

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Laying hens (commercial)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: amaqanda. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: ukukhiqiza amaqanda, cela izikhukhukazi ezincane ezigonyiwe esezizozalela kumhlinzeki wezinkukhu ezizalelayo, namarekhodi eminyaka, ukudla nokugonywa.

Hlela ukudla okunomsoco olinganiselwe, amanzi ahlanzekile, umthunzi, umoya nokuvikelwa ezilwaneni ezidlayo. Ukutholakala kwamaqanda kuncike ezinkukhwini ezizalelayo, ubudala bazo, impilo nokunakekelwa; ihhoko lodwa alinikezi amaqanda.

Qoqa amaqanda njalo, ugcine okwandlalwe ezidlekeni kuhlanzekile futhi komile, uphathe/ugcine amaqanda ngokucophelela ukuze asetshenziswe ekhaya noma athengiswe.

Inkomba yesikhathi: With a constant 14–16 hour light day, commercial hens keep laying at a high rate all year. Le nkomba yokuphatha kwezohwebo ayisho ukutholakala kwamaqanda kwasekhaya okulinganayo ngokwemvelo.

Inkomba yomkhiqizo wokuqala: amasonto angu-21. Ubudala benkukhu kusukela ekuchanyuselweni; qinisekisa uhlobo nokukhuliswa nomhlinzeki. Izinkukhu esezizozalela ziqala kamuva kulo mjikelezo.

Inkomba yokukhiqiza: 407.1–428.6 amaqanda/ngenkukhukazi efakwe endlini (hen housed), umjikelezo wokubekela kuze kube amasonto angu-90. Table 4a's cumulative 'Eggs/hen housed' for five commercial strains; the same table gives 'Age at depop. weeks 90.0' for every strain. Commercial barn/cage average, not a free-range smallholder trial. Kept as the source gives it: a per-year rate would need a point-of-lay age from another document.

Inkomba yobudala/uhlelo: SAPA 2024 commercial strain standards cumulative to 90 weeks: Dekalb Amberlink 429 eggs/hen housed, Dekalb Brown 428, Hy-Line Silver Brown 435, Hy-Line Brown 434 and Lohmann Brown Lite 413. These are full-cycle objectives, not eggs per calendar year or a smallholder forecast. Purchase age, actual onset of lay, surviving laying hens, feed, water and lighting need separate records.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Ezinhlelweni zokufuya okuvumela ukuhamba ngaphandle, izikhukhukazi kufanele zithole indawo yangaphandle okungenani amahora angu-8 ngosuku emini.

Lapho kusetshenziswa amabhokisi ezidleke ahlukene, kube okungenani nesidleke esisodwa ngezikhukhukazi ezingu-8.

Izikhukhukazi zidinga okungenani amahora angu-8 obumnyama obungaphazanyiswa kuwo wonke umjikelezo wamahora angu-24 ukuze ziphumule kahle.

Ukugcinwa phansi endlini kudinga okokwandlalela okungenani engxenyeni eyodwa kwezintathu (33%) yendawo yaphansi.

Indawo yangaphandle yokuzihambela akufanele igcwale ngokweqile: kungabi ngaphezu kwezinkukhu ezingu-5 ngemitha eyisikwele yendawo.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izinkukhu: Laying hens (commercial) (kuyaqhubeka)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Umthombo: SAPA Abridged Code of Practice: Commercial Layers, p. 1

Umthombo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Umthombo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Umthombo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: SAPA Industry Profile 2024, Table 4: commercial standards cumulative to 90 weeks, p. 39

Umthombo: SAPA Layer Notes, Commercial Layers (training module, November 2013), p. 13

Umthombo: SAPA Industry Profile 2022, Table 4a (Breed standards in laying hens), p. 39

Umthombo: ARC Climate-Smart Agriculture Training Manual, Poultry production, p. 8

Umthombo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Umthombo: FAO: Small-scale poultry production, egg marketing

Umthombo: SAPA Layer Notes, Commercial Layers (training module, November 2013)

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Amadada: Meat duck (Pekin / Muscovy, backyard)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Inkomba yomkhiqizo wokuqala: amasonto angu-7–12. P44 = Pekin (7 weeks to slaughter); MR71 = Muscovy (10 weeks for ducks/hens, 12 weeks for drakes). Treated as 'time to first product' = age to slaughter, since this is a single-cycle meat enterprise (no repeat product like eggs). Non-South-African trial by Warsaw researchers, published in SAJAS; journal location does not establish local applicability.

Inkomba yokukhiqiza: 2.3–2.9 kg yesisindo esiphilayo/ngenyoni ngesikhathi sokuhlatshwa. Non-SA (Guyana study reporting a citation to Steklenev 1990); no SA-specific absolute slaughter live-weight figure was found (see gaps). 2.93 kg = Pekin at 8-10 weeks, 2.29 kg = Muscovy at 8-10 weeks - matches this enterprise's two breeds. The same LRRD paper's own experiment (Table 2, intensive system, Guyana) measured live weight at slaughter (8 weeks) as: Pekin drakes 2.53 kg, Pekin hens 2.29 kg, Muscovy drakes 2.47 kg, Muscovy hens 1.51 kg - broadly consistent with, but not merged into, the cited range above since it is a separate dataset/table in the same source. No sourced eggs/duck/year figure was found for either breed (see gaps); the LRRD and SAJAS sources used here studied ducks reared only for meat.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Kuyicala ukunqabela idada ukudla noma amanzi, noma ukuligcina endaweni engenayo indawo eyanele, umoya owanele, ukukhanya noma indawo yokukhosela ekushiseni nasemakhazeni.

Bheka emihlambini ukufa okungazelelwe, izimpaphe ezimahliphihliphi, ukuvuvukala kwezicubu ezilenga ngaphansi koqhwaku noma amagekle, noma ukwehla okungazelelwe kokuzalela amaqanda - lezi yizimpawu zomkhuhlane wezinyoni futhi kufanele zibikwe ngokushesha.

Gcina amadada kude nezinyoni zasendle namanzi noma ukudla okuvulekile okuzidonsayo, futhi ulawule ukungena kwezivakashi ezindlini zamadada, ukuze unciphise ingozi yomkhuhlane wezinyoni.

Ducks are one of the domesticated animal species listed in the Meat Safety Act, so its rules on registered abattoirs and hygiene apply to them.

Ducks slaughtered to be sold must be slaughtered at a registered abattoir; slaughter for the farmer's own consumption is exempt but that meat may not be sold.

At the abattoir, ducks must be handled humanely during loading, transport, offloading, housing, immobilising and killing, under the Animals Protection Act.

Ducks are birds/poultry, so an outbreak of avian influenza in a duck flock is a controlled animal disease under the Animal Diseases Act and must be reported to the state vet immediately.

Umthombo: Animals Protection Act 71 of 1962 (consolidated text, s.2(1)(b)-(c)), p. 2

Umthombo: Western Cape Department of Agriculture, Avian Influenza (Bird Flu) factsheet, p. 2

Umthombo: Meat Safety Act 40 of 2000, Schedule 1 (Animals to which this Act applies), p. 16

Umthombo: Meat Safety Act 40 of 2000, s.7, p. 6

Umthombo: Meat Safety Act 40 of 2000, s.9, p. 7

Umthombo: Damaziak et al. (2014), Influence of housing system on the growth and histological structure of duck muscles, S. Afr. J. Anim. Sci. 44(2): Poland trial

Umthombo: Livestock Research for Rural Development 18(11) 2006 - Solomon et al. (Guyana; not an SA figure), citing Steklenev 1990

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Ichibi elincane: Tilapia (Mozambique tilapia, small pond or tank)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-2. Uhlobo lokukhiqiza olukhethiwe: inhlanzi. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Inkomba yomkhiqizo wokuqala: amasonto angu-43.5–52.2. 10 months x 4.35 = 43.5 weeks; 12 months x 4.35 = 52.2 weeks. This is Mozambique Tilapia (O. mossambicus) grow-out time to harvestable size; the source separately states Nile Tilapia (O. niloticus) reaches harvestable maturity faster, in 6-10 months (= 26.1-43.5 weeks) — not used here as it is a different species with a different legal status (see 'legal').

Inkomba yokukhiqiza: 800 g wesisindo ngesikhathi sokuvuna/ngenhlanzi. Table 8-3 gives the Mozambique Tilapia's harvestable size as an approximate single figure ('+- 800g'), not a range, so [800, 800] is used rather than inventing a spread. The same table cell wraps '10-12' across a line break in the PDF ('10-\n12'), kept verbatim. The table's Nile Tilapia row gives the same approximate weight (+-800g) reached faster, in 6-10 months — not used here because this dossier is framed around the indigenous, legally unrestricted Mozambique Tilapia per task instruction; see 'legal' for why Nile Tilapia is treated separately. No SA source was found giving a feed-conversion-based or per-hectare yield figure specific to O. mossambicus that could be converted to this per-fish unit without stitching numbers across documents, so outputUnit follows the source's own unit (grams harvest weight per fish) rather than being converted.

Inkomba yobudala/uhlelo: ARC makes tilapia suitability depend on water temperature: summer grow-out can be suitable anywhere in South Africa when water exceeds 20°C; growth is poor below 18°C and winter water below 13°C can kill farmed tilapia. The Highveld reference is a summer crop; tunnels can widen the range. Measure pond water and confirm stocking species and dates. Air climate or a mapped pond does not prove harvest months, fish numbers or food kg.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

The indigenous Mozambique Tilapia (O. mossambicus) is not a listed invasive species and may be farmed without an alien-species permit; the faster-growing Nile Tilapia (O. niloticus) is.

Nile Tilapia (Oreochromis niloticus) is a nationally listed invasive fish species: Category 1b inside National Parks, Provincial Reserves, Mountain Catchment Areas and Forestry Reserves, and Category 3 in all other areas.

Outside protected areas Nile tilapia is Category 3: it may remain in prescribed areas or provinces, but further propagation or trade is prohibited.

Nile Tilapia has been listed under the NEMBA Alien and Invasive Species Regulations as requiring a permit to produce.

Producing an alien or invasive species (such as Nile Tilapia) is prohibited in areas designated under the Protected Areas Act and the Fish Sanctuary Act; a permit is required to produce it elsewhere.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Ichibi elincane: Tilapia (Mozambique tilapia, small pond or tank) (kuyaqhubeka)

Producing tilapia in South Africa requires a water use licence under the National Water Act.

Umthombo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), Table 8-3 (Mozambique vs. Nile Tilapia), p. 69

Umthombo: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), List 7 (National List of Invasive Fresh-water Fish Species), p. 32

Umthombo: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), p. 3

Umthombo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 69

Umthombo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 74

Umthombo: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 75

Umthombo: ARC Aquaculture Production App: General FAQ, Can I grow tilapia on the Highveld?

Izinhlobo zezitshalo okufanele zihlolwe endaweni

Gcina igama lohlobo olukhethayo ngempela. Uhlu olufushane locwaningo ludinga ukuqinisekiswa kwesikhathi sendawo, amanzi nomhlabathi.

Tomatoes

Uhlobo olurekhodiwe: akukakhethwa.

Isimo sezulu sale ndawo asikacaci, ngakho akukho hlu olufushane lwendawo olushiwo. Vula ikhadi lesitshalo ukuze ubone imithombo bese urekhoda okukhethile.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.