SYNTHETIC GRAPH SOURCE {COUNT} $&

U nga si xava kumbe ku sungula

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Ntlawa wa maxelo wa ndzavisiso: Highveld. Ntlawa lowu a hi xitsundzuxo lexi tiyisisiweke xa purasi leri.

Leswi u swi vikeleke hi purasi leri

Mati hi nguva yo oma

A swi rhekhodiwanga. Hetisa vulavisisi bya ndhawu u tiyisisa mati u nga si hlawula kungu leri lavaka switirhisiwa swo tala.

Rharhama epurasini leri

Leswi voniweke hi rharhama endhawini a swi rhekhodiwanga. Miringaniso ya n’hweti ya sathelayiti a yi koti ku tiyisisa masiku yo sungula ni yo hetelela ya rharhama.

Dyambu ni ku huma ka mati emisaveni

Dyambu: a swi rhekhodiwanga.

Ku huma ka mati: a swi rhekhodiwanga. Mbuyelo wa ku kambela misava ni swilaveko swa xibyariwa xin’wana ni xin’wana swa ha fanele ku kamberiwa.

Tirhekhodo ta lembe ro vika - ku nga ri mvhumbo

Tinhlayo ni tin’hweti ta lembe leri hundzeke ro vika swi hlamusela leswi vikiweke. A swi tshembisi swakudya eka kungu leri taka.

Original eggs {year} $&

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 240 eggs.

Ku tirhisiwa ekaya: 0 eggs. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: Khot, Maw. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Fish

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: a swi rhekhodiwanga.

Ku tirhisiwa ekaya: a swi rhekhodiwanga. Swo xavisiwa: a swi rhekhodiwanga.

Tin’hweti ta ntshovelo leti vikiweke: a swi rhekhodiwanga. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

Miroho yin’wana

Nhlayo ya lembe leyi vikiweke hi murimi ya 2025: 12 kg.

Ku tirhisiwa ekaya: 7 kg. Swo xavisiwa: 3 kg.

Tin’hweti ta ntshovelo leti vikiweke: Sung, N'we. Leswi i leswi voniweke eka vulavisisi, ku nga ri mvhumbo wuntshwa wa vuhumelerisi.

Kambela yuniti ni tinhlayo leti rhekhodiweke u nga si tirhisa layini leyi.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso

Tinyoxi: Honeybee (managed hives)

Miako ya 1 leyi nga kona; ya 0 leyi ringanyetiweke. Bindzu leri hlawuriweke: vulombe. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Ntshovelo wu titshege hi ntlawa wa tinyoxi, ku rhumbuka ni mpfula ya ndhawu. Kambela ku vupfa u siyela ntlawa swakudya; a ku na nguva ya vulombe leyi vekiweke ya tiko hinkwaro.

Nkarhi wa xikumiwa xo sungula: a wu na xihlovo eka bindzu leri.

Xikombo xa vuhumelerisi: 25 kg ya vulombe/hi yindlu ya tinyoxi (hive)/hi lembe. From a 2008 national postal survey of 112 SABIO-member beekeepers (22.4% response rate) managing an estimated 19,520 hives at 2,094 apiary sites; figure is self-reported and un-weighted. Table 7 (same doc, p.13-14) shows wide variance by region/operation-focus/size for the same year: e.g. 'All' region, honey-focused, all-size operations = 25.1-26.9 kg/hive/yr across 3 survey sites; Western Cape honey-focused = 23.6-27.9 kg/hive/yr; Western Cape pollination-focused, small operations = only 7.0-7.5 kg/hive/yr. The same document also quotes an earlier, more rigorously validated study: 'Their average reported yield is very low, at just 13.4 kilograms per hive' for Western Cape hives (Allsopp & Cherry 2004, cited p.2 of this working paper), and historical national bee-census data of 8.3 kg/hive/annum (1911-1988 average) rising to 24 kg/hive in the 1974/5 census (same doc, p.1). No single number can be called definitive; 25 kg/hive/yr is used as the document's own headline national estimate.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Keep hives away from playgrounds, provide fresh water and use a dry site. Ask a registered beekeeper to check safe local placement before adding a hive.

Inspect every colony regularly for signs of notifiable bee disease and Cape-bee (Capensis) infection.

Site and secure hives against theft and vandalism, which is one of the biggest challenges facing South African beekeepers.

Register with the Department before beekeeping. Registration lasts 24 months from registration or renewal and must be renewed when it expires.

A beekeeper who fails to renew registration at expiry will be deregistered and face penalties under the Act.

Beekeeping activities and hired beekeeping services require a valid registration certificate. Check the certificate before using a beekeeper.

Every beehive must be clearly and legibly marked with the beekeeper's registration number.

Keep written records of colonies, apiary locations, diseases, disease-control measures and colony losses with their reasons.

Do not move honeybees across the Cape bee line, or keep Cape bees north of it, without written approval.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tinyoxi: Honeybee (managed hives) (swi ya mahlweni)

Report any suspected notifiable disease (American foulbrood is listed); officials can order an infected colony destroyed.

Xihlovo: FAO: Beekeeping in Africa, Site selection (general African apiary guidance)

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 6

Xihlovo: Gums & Bees: A roadmap for landowners in South Africa (South African National Biodiversity Institute, SANBI), p. 21

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(1) and 2(4), p. 4

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measure 2(5), p. 4

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(8) and 2(9), p. 5

Xihlovo: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 5

Xihlovo: Survey of Beekeeping in South Africa, CSSR Working Paper No. 221 (Conradie & Nortje, 2008, UCT Centre for Social Science Research), p. 14

Xihlovo: FAO, Beekeeping in Africa, honey harvesting

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Nguni / crossbred beef cattle (communal and extensive veld)

Miako ya 1 leyi nga kona; ya 0 leyi ringanyetiweke. Bindzu leri hlawuriweke: nyama. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Nkarhi wa xikumiwa xo sungula: a wu na xihlovo eka bindzu leri.

Xikombo xa vuhumelerisi: 114.5–121.2 kg ya ntiko loko ri tshikisiwa ku mama/hi rhole. Table 5 gives weaning weight (WW) across four agro-ecological zones for the same nine Nguni communal herds: arid 114.51 kg (lowest), semi-arid 119.98 kg, dry sub-humid 117.91 kg, humid 121.21 kg (highest). The range recorded here spans this one study's own arid-to-humid zone means; it is not merged with any other paper's weaning-weight figures (other SA studies report different ranges for different systems - e.g. research-station or stud herds - and were deliberately excluded to avoid mixing systems). 826 Nguni calves from nine herds that the paper itself says were 'kept under extensive grazing system' in Limpopo Province - a genuine communal/extensive veld system matching this enterprise.

Xikombo xa nkarhi wo humelerisa: malembe ya 6.4. 2354 days / 365.25 = 6.44 years. 'Failure time' is the length of productive herd life for the 580 uncensored (fully-observed) records in the combined dataset; the dissertation itself defines productive herd life as 'the time interval between first calving date to culling date'. Combined data from three extensively-grazed Nguni stud herds, animals born 1968-2015.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Cattle pens must not be overcrowded: there must be enough floor space for every animal to lie down at the same time.

During transport, cattle must have at least 1.4 m^2 of floor space each so loading is neither too tight nor too loose.

Cattle may not be transported for more than 18 hours without being off-loaded and given feed, water and shelter.

It is an offence to starve, under-feed, or deny water or food to cattle.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, as part of the national bovine brucellosis control scheme under the Animal Diseases Act.

No animal may be slaughtered anywhere but a registered abattoir, with the exception below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Nguni / crossbred beef cattle (communal and extensive veld) (swi ya mahlweni)

Slaughter for the farmer's own household consumption, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.4.6.1, p. 7

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Xihlovo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Xihlovo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Xihlovo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Xihlovo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: Mpofu, Ginindza, Siwendu, Nephawe & Mtileni (2016). Effect of agro-ecological zone, season of birth and sex on pre-weaning performance of Nguni calves in Limpopo Province, South Africa. Tropical Animal Health and Production 49:187-194 (PMC5203849), Table 5

Xihlovo: Ngayo, M. (2017). Factors Affecting Productive Life and Fertility in Nguni Cows (MSc dissertation, University of the Free State), Table 5.1, p. 58

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Small-scale dairy cow (Jersey, Holstein or crossbred)

Miako ya 1 leyi nga kona; ya 0 leyi ringanyetiweke. Bindzu leri hlawuriweke: masi. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: fananisa muxaka/swifuwo ni xikumiwa, swakudya swa ndhawu, maxelo ni vuswikoti byo hlayisa; tiyisisa ni mutsundzuxi wa ndhawu wa swifuwo kumbe ku fuwa tinyoxi.

Kunguhata hi ku ya hi ku tswariwa ka xiviri ka marhole/swimbutana ni ku humesa masi. Swifuwo leswi nga humesiki masi a swi nyiki masi.

Xikombo xa xikumiwa xo sungula: mavhiki ya 121.8–130.5. Table 1, AFC column, rows in the table's order Holstein-Friesland Reg./Grade, Jersey Reg./Grade, Ayrshire Reg./Grade: 28, 30, 25, 28, 29, 32 months. The Grade rows used here are Holstein-Friesland 30 and Jersey 28 (Grade Ayrshire 32 is not used: the enterprise is Jersey or Holstein type). 28 x 4.35 = 121.8 weeks; 30 x 4.35 = 130.5 weeks. Milk begins at first calving, so age at first calving is the time to first product.

Xikombo xa vuhumelerisi: 3840–5422 kg ya masi/hi homo ya xisati/hi nkarhi wo humesa masi. Table 1 ('Comparison of number of cows, production and fertility in the main dairy breeds in South Africa'), Milk column, Grade rows: Jersey 3840 kg and Holstein-Friesland 5422 kg per lactation (Registered rows: 4468 and 6426 kg). The document says these are National Milk Recording Scheme herds, which 'have been shown to be better managed than those outside the scheme, with their performance per lactation being 50 % greater than that of the national herd', so a smallholder herd should expect less.

Xikombo xa nkarhi wo humelerisa: malembe ya 1.9–2. Holstein 739.33 and Jersey 696.81 days of productive life (first calving to culling) in South African milk-recorded herds, 2000-2019: 696.81/365.25 = 1.91 and 739.33/365.25 = 2.02 years.

Xikombo xa vukhale/endlelo: In hotter, humid KZN, KZN DARD describes Nguni × Jersey dairy trials at Makhathini to combine milk potential with local adaptation. Hand milking with the calf present highlights competition between household milk and calf growth. Discuss adapted stock and feeding locally; commercial milk-recording totals do not forecast this system. Record calving, lactation stage and milk available after calf needs.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Cattle pens must not be overcrowded: there must be enough floor space for every animal to lie down at the same time.

Cows must not be loaded for transport if their udders are distended with milk.

Cattle may not be transported for more than 18 hours without being off-loaded and given feed, water and shelter.

It is an offence to starve, under-feed, or deny water or food to a dairy cow.

Cows on pasture or in open camps need shade: shade structures should give at least 4 to 5 m^2 of shade per cow.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Dairy farmers are legally required to test their herds for bovine TB, mainly because of export requirements, so most dairy cattle in South Africa are tested.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, under the national bovine brucellosis control scheme.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tshanga: Small-scale dairy cow (Jersey, Holstein or crossbred) (swi ya mahlweni)

A cull cow may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.6.13, p. 9

Xihlovo: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Xihlovo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 129

Xihlovo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Xihlovo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Xihlovo: Bovine tuberculosis / dairy herd testing study, PMC10369683

Xihlovo: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: KZN DARD: Subsistence Dairying in KZN, Makhathini Research Station trials

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 10

Xihlovo: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), Table 21.1, p. 122

Xihlovo: Gertenbach, W.D., Dairying in KwaZulu-Natal: Breeds of Dairy Cattle, Cedara Agricultural Development Institute, KZN Department of Agriculture and Rural Development, Table 1, p. 1

Xihlovo: Cow Longevity and Reasons and Risk Factors for Culling in South African Holstein and Jersey Dairy Herds, Animals 2025 (PMC12560872)

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Broilers (commercial)

Miako ya 1 leyi nga kona; ya 0 leyi ringanyetiweke. Bindzu leri hlawuriweke: nyama. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Xitsundzuxo xo xava: hlawula muxaka wa nyama ni kungu ra mintlawa ra muphakeri; tiyisisa ku kufumeta swinkukwana, swakudya, mati ni malunghiselelo yo tlhava.

Kunguhata swakudya leswi ringanisiweke, mati yo tenga, ndzhuti, ku nghena ka moya ni nsirhelelo eka swihadyana leswi dlayaka. Mphakelo wa matandza wu titshege hi tihuku leti tshikelaka, vukhale bya tona, rihanyo ni nhlayiso; hoko yi ri yoxe a yi nyiki matandza.

Xikombo xa xikumiwa xo sungula: mavhiki ya 4.6–4.7. 32 days / 7 = 4.57 weeks; 33 days / 7 = 4.71 weeks. This enterprise has a single grow-out cycle ending in slaughter, so 'time to first product' = age at slaughter. South Africa, commercial industry average.

Xikombo xa vuhumelerisi: 1.8–1.9 kg/hi xinyenyana loko xi tlhaviwa (masiku ya 32–33). National commercial broiler industry average, South Africa, 2022. A separate breed-trial table (Table 4b, same document, p.39) gives named-strain 'Cumulative FI (kg)' and 'Body weight (kg)' at exactly 35 days: Arbor Acres Plus 2.29kg, Cobb 500 2.52kg, Indian River 2.30kg, Ross 308 2.30kg (cumulative feed intake 3.21-3.64kg) - a slightly older age (35 vs 32-33 days) and higher weight than the industry-wide average quoted above; not merged into the primary figure since it is a different (breed-standard trial) dataset, but used below for the feed-intake figure.

Xikombo xa vukhale/endlelo: Newer South African commercial reference: SAPA 2024 reports average slaughter at 30–31 days with bird weight 1.8–1.85 kg. The primary range above remains the earlier 2022 reference at 32–33 days. These are commercial bird weights, not dressed meat kg or a village flock target. Record hatch or purchase age, feeding system, actual harvest date and weighed product basis.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

Floor-housed broilers: no more than 40 kg of bird per square metre.

Newly hatched chicks must be given feed within 24 hours of hatching.

Newly hatched chicks must be given access to water within 24 hours of hatching, sooner in hot weather.

Fresh water and the right feed always available, so birds are free from hunger and thirst.

Broiler houses need a minimum ventilation/airflow rate of 5 cubic metres per hour for every kilogram of bird mass in the house.

In free-range broiler production, indoor stocking must not exceed 15 birds per square metre of floor space.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Broilers (commercial) (swi ya mahlweni)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), 2.2 Housing of broilers in Floor Systems, p. 6

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 11

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 12

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 7

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), p. 8

Xihlovo: SAPA Code of Practice for Broiler Production (May 2012), 2.4 Free Range Broiler Production, p. 8

Xihlovo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Xihlovo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: SAPA Industry Profile 2024, section 7.9 Performance efficiency, p. 82

Xihlovo: SAPA Industry Profile 2022, section 6.9 Performance efficiency, p. 79

Xihlovo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Village chickens (indigenous, free-range)

Miako ya 1 leyi nga kona; ya 0 leyi ringanyetiweke. Bindzu leri hlawuriweke: matandza. A ku ehleketiwi nhlayo ya swifuwo kumbe vuhumelerisi.

Buying advice: Potchefstroom Koekoek is a dual-purpose option; Venda, Ovambo and Naked Neck are other ARC-conserved options. No province-specific winning breed is claimed.

Kunguhata swakudya leswi ringanisiweke, mati yo tenga, ndzhuti, ku nghena ka moya ni nsirhelelo eka swihadyana leswi dlayaka. Mphakelo wa matandza wu titshege hi tihuku leti tshikelaka, vukhale bya tona, rihanyo ni nhlayiso; hoko yi ri yoxe a yi nyiki matandza.

Hlengeleta matandza nkarhi ni nkarhi, hlayisa leswi andlariweke eswisakeni swi tengile naswona swi omile, khoma/hlayisa matandza hi vukheta leswaku ma tirhisiwa ekaya kumbe ku xavisiwa.

Xikombo xa xikumiwa xo sungula: mavhiki ya 15.7–27.8. Free-range village chickens, KwaZulu-Natal; dry-environment birds started laying earlier (15.74 weeks) than wet-environment birds (27.78 weeks), difference flagged significant ('*') in the source table. Wide range reflects genuine dry/wet-season environmental variation reported by the source, not measurement uncertainty.

Xikombo xa vuhumelerisi: 40.4–49.9 matandza/hi huku ya xisati/hi lembe. Free-range/scavenging village chickens, KwaZulu-Natal: dry (40.43) vs wet (49.91) environment, difference not significant ('NS'). A managed-system ARC Irene trial of named indigenous breeds (Grobbelaar, Sutherland & Molalakgotla 2010, Animal Genetic Resources 46: 25-32) reports far higher counts, now separately verified and retained in productionReferences; managed conditions cannot replace this village range.

Xikombo xa vukhale/endlelo: The ARC Irene trial compared Potchefstroom Koekoek, Venda, Ovambo and Naked Neck under commercial feed, water and controlled lighting. Mean eggs/hen in a 52-week cycle were 195.9, 153.7, 125.5 and 138.9 respectively; first-egg ages were 138.5, 139.0, 134.5 and 129.0 days from hatch, with no significant first-egg age difference among breeds. This is managed breed context, not the yield of scavenging hens in KZN. Match feeding and housing before using a benchmark.

Kambela swikombo leswi nga ni vuxokoxoko swa bindzu swa swakudya ni vutshamo. Fananisa mpimo wa swakudya ni ndhawu ni xiyimo xa xifuwo ni endlelo ra vutshamo; nyika mati yo tenga lama kumekaka nkarhi ni nkarhi.

In extensive village systems the chickens have to find their own water and food.

Scavenging village chickens grow more slowly than commercially-farmed birds and are more vulnerable to disease, parasites and predators.

In the Limpopo and KwaZulu-Natal survey, 94.9% of farmers housed their chickens at night.

What free-ranging hens forage met their needs for maintenance and growth but not for laying: protein in every season, calcium and phosphorus in winter.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Swihlawuriwa swa purasi - swi ya mahlweni

Tirhisa leswi u swi voneke epurasini ni minxaxamelo yo koma ya ndzavisiso. Tiyisisa swihlawuriwa endhawini u nga si xava.

Swikumiwa swa swifuwo ni nhlayiso (swi ya mahlweni)

Tihuku: Village chickens (indigenous, free-range) (swi ya mahlweni)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Xihlovo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 471

Xihlovo: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 468

Xihlovo: Malatji, Tsotetsi, van Marle-Koster & Muchadeyi (2016), 'A description of village chicken production systems and prevalence of gastrointestinal parasites: Case studies in Limpopo and KwaZulu-Natal provinces of South Africa', Onderstepoort Journal of Veterinary Research 83(1)

Xihlovo: Raphulu, Jansen van Rensburg & van Ryssen (2015), 'Nutrient content of crop contents of free ranging indigenous chickens raised under village conditions', South African Journal of Animal Science 45(2), p. 1

Xihlovo: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Xihlovo: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Xihlovo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Xihlovo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Xihlovo: Grobbelaar, Sutherland & Molalakgotla (2010), Egg production potentials of certain indigenous chicken breeds from South Africa: methods pp27–28, first-egg p29, Table 4 p31, p. 31

Xihlovo: Tenza, Mhlongo & Chimonyo (2023), 'Village chicken production and egg quality in dry and wet, resource-limited environments in KwaZulu-Natal, South Africa', South African Journal of Animal Science 53(6), p. 5

Xihlovo: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Xihlovo: FAO: Small-scale poultry production, egg marketing

Tinxaka ta swibyariwa to kambela endhawini

Hlayisa vito ra muxaka lowu u wu hlawulaka hakunene. Minxaxamelo yo koma ya ndzavisiso yi lava ku tiyisisiwa eka nguva ya ndhawu, mati ni misava.

Tomatoes

Muxaka lowu rhekhodiweke: a wu si hlawuriwa.

Nxaxamelo wo koma wa ndzavisiso wa maxelo yo fana: Rodade.

Xihlovo: Rodade, Sakata product page

A yi si kamberiwa hi mutivi wa vurimi — i nkongomiso wo pulana, a hi xileriso. Yi kombe mutsundzuxi wa wena wa vurimi, kutani u kambela misava u nga si tirhisa mali eka yona.
Vuhundzuluxi bya xitsalwana: a byi si kambisisiwa hi munhu la tivaka ririmi kahle. Kambisisa xihlovo xa Xinghezi u nga si endla leswi vuriwaka.