SYNTHETIC GRAPH SOURCE {COUNT} $&

Ngaphambi kokuthenga noma ukuqala

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Iqembu lesimo sezulu locwaningo: Highveld. Leli qembu alisona isincomo esiqinisekisiwe saleli pulazi.

Lokho okubikile ngaleli pulazi

Amanzi ngesikhathi esomile

Akurekhodiwe. Qedela inhlolovo yendawo bese uqinisekisa amanzi ngaphambi kokukhetha uhlelo oludinga izinsiza eziningi.

Isithwathwa kuleli pulazi

Okubonwe ngesithwathwa endaweni akurekhodiwe. Izilinganiso zanyanga zonke zesathelayithi azikwazi ukuthola izinsuku zokuqala nezokugcina zesithwathwa.

Ilanga nokuphuma kwamanzi

Ilanga: akurekhodiwe.

Ukuphuma kwamanzi: akurekhodiwe. Kusadingeka ukuhlola imiphumela yokuhlolwa komhlabathi nezidingo zesitshalo ngasinye.

Amarekhodi onyaka wokubika - hhayi isibikezelo

Izamba nezinyanga zonyaka wokubika odlule zichaza lokho okwabikwa. Azithembisi ukudla ohlelweni oluzayo.

Original eggs {year} $&

Isamba sonyaka esibikwe umlimi sango-2025: 240 eggs.

Okusetshenziswa ekhaya: 0 eggs. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: Jun, Jul. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Fish

Isamba sonyaka esibikwe umlimi sango-2025: akurekhodiwe.

Okusetshenziswa ekhaya: akurekhodiwe. Okuthengisiwe: akurekhodiwe.

Izinyanga zokuvuna ezibikiwe: akurekhodiwe. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Eminye imifino

Isamba sonyaka esibikwe umlimi sango-2025: 12 kg.

Okusetshenziswa ekhaya: 7 kg. Okuthengisiwe: 3 kg.

Izinyanga zokuvuna ezibikiwe: Jan, Dis. Lokhu ukubonwe kwinhlolovo, akusona isibikezelo esisha sokukhiqiza.

Hlola iyunithi nezamba ezirekhodiwe ngaphambi kokusebenzisa lo mugqa.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela

Izimbuzi: Boer goat (meat enterprise)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-0. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 1.5 amazinyane ezimbuzi alumuliwe/ngembuzikazi ekhwelwe (doe mated). The reported weaning rate is 149%, or 1.49 kids weaned per doe mated. The source describes observations over twenty years for extensive Boer does at 295 mm rainfall; it does not state kids weaned per doe per calendar year. This is a reproductive outcome, not kg of live animals or dressed meat. A separate mean litter size of 1.93 kids born per parturition has a different denominator and stage and is not merged into this figure.

Inkomba yobudala/uhlelo: The 2016 government goat manual distinguishes indigenous, Boer, Savanna and Kalahari Red meat-purpose stock from milk/fibre breeds, with breeding objectives and individual birth, weaning, weight and sale records. Uncontrolled communal mating can produce births at any time; that does not establish meat or weaned-kid availability all year. Choose stock for the local system and record survival and product stage before forecasting quantities.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

It is an offence to keep a goat somewhere with inadequate space, ventilation, light or shelter from heat, cold or weather, or to confine/tether it in a way that causes unnecessary suffering.

It is an offence to starve, under-feed, or deny water or food to a goat.

An owner must not let a goat stay in a dirty or parasite-infested condition, and must get veterinary or other treatment for a sick, injured or diseased animal when able to.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: veterinary services enforce isolation and movement restrictions, and a suspected case must be reported.

Inside a declared FMD-affected zone, goats and their products cannot be moved into, out of, or within the zone without a state veterinary movement permit and compliance with its conditions.

Every goat owner must register an official identification mark and mark their goats with it, under the Animal Identification Act.

A goat may only be slaughtered for sale at a registered abattoir under the Meat Safety Act; slaughtering it for the farmer's own consumption, or for cultural/religious purposes, is exempt from that rule, but that meat may not then be sold.

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(b), p. 1

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(e), p. 1

Umthombo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: DAFF (2016), Goat Training Manual: breeds p2, records pp4–5, uncontrolled mating section 2.4.6 p7, p. 7

Umthombo: King, F.J.M. (2009). Production Parameters for Boer Goats in South Africa (MSc dissertation, University of the Free State), citing National Dept of Agriculture data via Malan, S.W. (2000)

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izingulube: Pig (small-scale / emerging-farmer sow herd)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-0. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yobudala/uhlelo: ARC Table 1 lists pregnant sows at 10–18 L drinking water/day and lactating sows at 18–23 L/day. These are breeding-sow stages, not one amount for every pig. Farrowing care separates sow cooling from piglet warmth; outdoor systems need locally adapted hardy stock, shade and shelter. Confirm stage, reliable clean water, feed and housing locally. The manual supplies no verified edible kg per breeding sow for this plan.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

When pigs are trucked, the SA code of practice for handling and transporting animals recommends 0.3 m^2 per porker, 0.4 m^2 per baconer and 0.8 m^2 per adult pig.

It is an offence to keep a pig somewhere with inadequate space, ventilation, light or shelter from heat, cold or weather, or to confine/tether it in a way that causes unnecessary suffering.

It is an offence to starve, under-feed, or deny water or food to a pig.

In heat waves, pig houses need better ventilation/airflow and lower stocking density, and outdoor pigs need shade and shelter.

African swine fever (ASF) is listed as a controlled animal disease pertaining to pigs under the Animal Diseases Act; warthogs and Ornithodorus ticks are carriers/vectors.

Foot-and-mouth disease (FMD) is also listed as a controlled animal disease pertaining to pigs under the Animal Diseases Act.

Swill (food waste) may not be fed to pigs unless it has been boiled for at least 60 minutes or sterilised by another equally effective method.

Pigs may only be moved into a biosecure compartment from another compartment that is officially certified/approved for that purpose.

A pig may not be slaughtered anywhere but a registered abattoir, except for own household consumption or cultural/religious purposes.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Izingulube: Pig (small-scale / emerging-farmer sow herd) (kuyaqhubeka)

A pig owner must apply to register an official identification mark and mark their pigs with it, under the Animal Identification Act.

Umthombo: Muchenje, V. & Ndou, S.P., How pig pre-slaughter welfare affects pork quality and the pig industry (Univ. of Fort Hare, hosted by SAPPO), quoting Tomlinson (2000) Code of Practice for the Handling and Transport of Livestock, p. 7

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(b), p. 1

Umthombo: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Umthombo: ARC Climate-Smart Agriculture Training Manual, Module 10: Pig Production, p. 9

Umthombo: ARC Climate-Smart Agriculture Training Manual, Module 10: Pig Production, Table 2, p. 15

Umthombo: Munzhelele et al., Investigation of African swine fever outbreaks in pigs outside the controlled areas of South Africa, 2012-2017 (Journal of the South African Veterinary Association, SciELO SA mirror)

Umthombo: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: ARC Climate-Smart Agriculture: Pig production, farrowing/adaptation pp442–445, Table 1 p446, p. 446

Umthombo: ARC Climate-Smart Agriculture Training Manual, Module 10: Pig Production, Table 1 (Water requirements of pigs, litres/pig/day), p. 10

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Onogwaja: Meat rabbit (New Zealand White / Californian, backyard hutches)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-0. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 30–35 onogwaja abancane abafinyelela ebudaleni bokuhlatshwa/bokuzala/ngensikazi yonogwaja/ngonyaka. FAO general backyard target, not a South African observed rate. The 30–35 young per doe annually have reached slaughter or reproduction age after allowance for post-weaning loss; they are not merely weaned and are not kg of meat. The source models six litters and 5.5 surviving young per litter, describing a 30-rabbit objective; an extra litter raises its stated objective to 35. Do not combine this backyard target with grower-house carcass weight from a different trial.

Inkomba yesikhathi sokukhiqiza: iminyaka engu-0.8. 279 days / 365 = 0.76 years. This figure is from French INTENSIVE commercial rabbitries (average of 1992 national recording-scheme data), NOT backyard/extensive hutches - the source gives no separate productive-life figure for backyard systems (see gaps). Treat as a lower-bound / industrial-system reference only.

Inkomba yobudala/uhlelo: South African grower reference: New Zealand White and Phendula rabbits in the 2017 Stellenbosch trial were slaughtered at about 11 weeks from birth. Overall means were 2.62 kg live slaughter weight, 1.55 kg hot carcass, 1.48 kg chilled carcass and 1.26 kg reference carcass with head and red offal removed, still including bones. Balanced pellets and water were freely available in a grower house. These weights are not boneless meat or annual doe output; do not multiply them by the unrelated FAO backyard target.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

It is an offence to deny a rabbit food or water, or keep it somewhere lacking adequate space, ventilation, light or shelter from heat and cold.

Give breeding rabbits roughly 0.48-0.70 m2 of hutch floor space each, 50-60 cm high, not a cramped cage.

Rabbits are one of the domesticated animal species listed in the Meat Safety Act, so the Act's rules on abattoirs and hygiene apply to them, as they do to ducks and other listed species.

Rabbits slaughtered to be sold must be slaughtered at a registered abattoir; slaughter for the farmer's own consumption is exempt from this rule but the meat from it may not be sold.

At the abattoir, rabbits (like other listed animals) must be handled humanely during loading, transport, offloading, housing, immobilising and killing, under the Animals Protection Act.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Onogwaja: Meat rabbit (New Zealand White / Californian, backyard hutches) (kuyaqhubeka)

The wild European rabbit (the same species as farmed New Zealand White/Californian rabbits) is only listed as an invasive species under NEMBA on South Africa's off-shore islands - it is not a listed/restricted species on the mainland.

Umthombo: Animals Protection Act 71 of 1962 (consolidated text, s.2(1)(b)-(c)), p. 2

Umthombo: FAO Animal Production and Health Series No.21 - The Rabbit: Husbandry, Health and Production (Lebas et al.), p. 172

Umthombo: Meat Safety Act 40 of 2000, Schedule 1 (Animals to which this Act applies), p. 16

Umthombo: Meat Safety Act 40 of 2000, s.7, p. 6

Umthombo: Meat Safety Act 40 of 2000, s.9, p. 7

Umthombo: South Africa's National Listed Invasive Species booklet (NEMBA Alien and Invasive Species Lists), p. 27

Umthombo: North et al. (2019), Growth, carcass and meat quality traits of two South African meat rabbit breeds: methods p816, Table 2 p819, p. 819

Umthombo: FAO Animal Production and Health Series No.21 - The Rabbit: Husbandry, Health and Production (Lebas et al.), p. 107

Umthombo: FAO Animal Production and Health Series No.21 - The Rabbit: Husbandry, Health and Production (Lebas et al.), p. 201

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Meat sheep (Dorper, Damara, indigenous fat-tailed and crossbred)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-0. Uhlobo lokukhiqiza olukhethiwe: inyama. Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 0.6–1.4 amawundlu azelwe ngemvukazi. outputUnit is 'lambs born/ewe/year' rather than the suggested 'kg live weight/lamb at weaning or slaughter' or 'lambs weaned/ewe/year': the source explains 'Measuring accurate lambing percentages for smallholder sheep farmers is challenging as these farmers mostly farm extensively and do not keep records', and instead calculates 'Lambs born per ewe (as recorded from the surveys based on interviews with individual respondents)' - a lambing (born), not weaning, figure, self-reported by smallholder farmers rather than measured/weighed. 139%=1.39, 83%=0.83, 61%=0.61 lambs born/ewe/year; the range [0.61, 1.39] spans the three Western Cape smallholder districts surveyed (Eden, Central Karoo, West Coast), all from this one study/quote - not merged with any other document. The same paper separately states a COMMERCIAL extensive benchmark of 110-156% lambing and 94-132% weaning (Snyman & Herselman, 2005), explicitly 'expressed over ewes joined' (a different denominator/system) - not merged into this smallholder range.

Inkomba yobudala/uhlelo: The government sheep manual describes controlled breeding at Mara Research Station, Limpopo: June–July mating and November–December lambing to match rain and grass. This is one station breeding calendar, not mutton harvest months. Match adapted meat-purpose stock and feed locally, then record birth, survival, weaning and actual slaughter or sale; lambs born per ewe are not edible kg.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Sheep must not be penned overcrowded: at least 0.56 square metres of floor space per sheep so every animal can lie down.

Electric prodders, sticks or goads may not be used on sheep.

Sheep driven on the hoof may not be moved more than 20 km in a single day (25 km is regarded as excessive).

Livestock must be given water and fed as soon as they reach their night camp or final destination.

Injured, disabled or blind sheep must be given priority slaughter, or emergency slaughter, to prevent further pain or distress.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act; Veterinary Services enforce isolation and movement restrictions.

Inside a declared FMD zone, sheep and their products cannot be moved into, out of, or within it without a state veterinary permit.

Sheep scab is a controlled disease by law: a suspected or positive case must be reported to the State Veterinarian immediately.

Against sheep scab, all small stock on the farm must be treated with a remedy registered under Act 36 of 1947, and infected sheep treated twice, eight to ten days apart.

Every sheep owner must register an official identification mark and mark their sheep with it, under the Animal Identification Act.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Meat sheep (Dorper, Damara, indigenous fat-tailed and crossbred) (kuyaqhubeka)

A sheep may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.1.2, p. 4

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.2.7, p. 5

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.3.4.1, p. 6

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.3.5, p. 6

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.2.13, p. 5

Umthombo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Umthombo: KZN Department of Agriculture and Rural Development, Research & Technology Bulletin 2017-18/01: Sheep Scab, p. 1

Umthombo: KZN Department of Agriculture and Rural Development, Research & Technology Bulletin 2017-18/01: Sheep Scab, p. 3

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: DAFF: Sheep General Information Manual, Final Draft, section 3.4.6, p. 12

Umthombo: Factors influencing off-take rates of smallholder sheep farming systems in the Western Cape Province of South Africa (South African Journal of Agricultural Extension, 47(3), 2019)

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Wool sheep (Merino / Dohne Merino, incl. communal wool growers)

Izakhiwo ezikhona ezingu-1; ezihlongozwayo ezingu-0. Uhlobo lokukhiqiza olukhethiwe: uboya (akukona ukudla). Akucatshangwa isibalo sezilwane noma okukhiqizwayo.

Iseluleko sokuthenga: fanisa uhlobo/izilwane nomkhiqizo, ukudla kwendawo, isimo sezulu namandla okunakekela; qinisekisa nomeluleki wendawo wemfuyo noma wokufuywa kwezinyosi.

Isikhathi somkhiqizo wokuqala: asinamthombo walolu hlobo lokukhiqiza.

Inkomba yokukhiqiza: 4–5 kg woboya obunamafutha/ngemvu/ngonyaka. Semi-arid/extensive-veld range (4-5 kg greasy wool/sheep/year) used as the primary value; the same sentence's 8 kg/year figure is for a different system (sheep grazing CULTIVATED pastures) and is deliberately excluded from the range rather than averaged in. A Western Cape Government (Elsenburg) Dohne Merino Extensive enterprise budget (Ceres-Karoo, 2024) separately reports total flock wool sold as 4439.64 kg from a flock of '1 519' Smallstock Units, i.e. 4439.64/1519 = 2.92 kg per Smallstock Unit/year - not used as the primary value because the source does not state that 1 Smallstock Unit = 1 sheep (SSU is a standardised stock-class weighting, not a literal head count), so converting it to kg/sheep would require an unstated conversion factor. A separate NWGA/Mandela Initiative paper on the Eastern Cape communal Wool Sheep Development Program reports a Grootfontein ADI progeny-test 'Greasy fleece weight (kg) 2.8 [commercial rams] 2.4 [communal rams]' for communal Eastern Cape flocks - not merged into the primary value because that table is explicitly for '6-month old lambs' (a different life stage from a mature sheep's full annual clip), consistent with the rule against mixing figures across life stages; kept here only as EC-communal context: commercial-ram-sired lambs at 6 months averaged 2.8 kg, communal-ram-sired lambs 2.4 kg, a +16% difference.

Hlola izinkomba ezinemininingwane zohlobo lokukhiqiza mayelana nokudla nezindawo zokuhlala. Fanisa isabelo sokudla nendawo nesigaba sesilwane nohlelo lokuhlalisa; nikeza amanzi ahlanzekile athembekile.

Sheep must not be penned overcrowded: at least 0.56 square metres of floor space per sheep so every animal can lie down.

Electric prodders, sticks or goads may not be used on sheep.

Sheep driven on the hoof may not be moved more than 20 km in a single day (25 km is regarded as excessive).

Injured, disabled or blind sheep must be given priority slaughter, or emergency slaughter, to prevent further pain or distress.

A sheep-scab treatment plan should be in place: the disease carries a high economic and welfare cost.

Despite foot-and-mouth disease (FMD) in South Africa, wool exports continue under the inactivation clause of the OIE (World Organisation for Animal Health) rules.

Inside a declared FMD zone, sheep and their products cannot be moved into, out of, or within it without a state veterinary permit.

Sheep scab is a controlled disease by law: a suspected or positive case must be reported to the State Veterinarian immediately.

Wool from scab-infected sheep must be destroyed rather than sent to the buyer, where it would be marked as scab wool.

Every sheep owner must register an official identification mark and mark their sheep with it, under the Animal Identification Act.

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Izinketho zepulazi - kuyaqhubeka

Sebenzisa lokho okubonile epulazini nohlu olufushane locwaningo. Qinisekisa izinketho endaweni ngaphambi kokuthenga.

Imikhiqizo yezilwane nokunakekela (kuyaqhubeka)

Isibaya: Wool sheep (Merino / Dohne Merino, incl. communal wool growers) (kuyaqhubeka)

A sheep may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.1.2, p. 4

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.2.7, p. 5

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.3.4.1, p. 6

Umthombo: Livestock Welfare Co-ordinating Committee, Code of Practice for the Handling and Transport of Livestock (2018), s.2.13, p. 5

Umthombo: KZN Department of Agriculture and Rural Development, Research & Technology Bulletin 2017-18/01: Sheep Scab, p. 3

Umthombo: Cape Wools SA Annual Review 2020/2021, p. 8

Umthombo: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Umthombo: KZN Department of Agriculture and Rural Development, Research & Technology Bulletin 2017-18/01: Sheep Scab, p. 1

Umthombo: Animal Identification Act 6 of 2002, s.7(1), p. 4

Umthombo: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Umthombo: Assessing Socio-economic Factors Influencing Wool Production in Kolomana Villages of Eastern Cape, South Africa (South African Journal of Agricultural Extension), citing Bot et al., 2004

Izinhlobo zezitshalo okufanele zihlolwe endaweni

Gcina igama lohlobo olukhethayo ngempela. Uhlu olufushane locwaningo ludinga ukuqinisekiswa kwesikhathi sendawo, amanzi nomhlabathi.

Tomatoes

Uhlobo olurekhodiwe: akukakhethwa.

Uhlu olufushane locwaningo lwezimo zezulu ezifanayo: Rodade.

Umthombo: Rodade, Sakata product page

Awukahlolwa uchwepheshe wezolimo — umhlahlandlela wokuhlela, hhayi umyalelo. Wubonise kumeluleki wakho wezolimo, futhi uhlole umhlabathi ngaphambi kokuchitha imali kuwo.
Ukuhumusha okusabuyekezwa: akukahlolwa ngumuntu olwazi kahle lolu limi. Hlola umbhalo wesiNgisi ngaphambi kokwenza okushiwoyo.