SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Natal currant

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible Natal currant fruit and local phenology: The primary account establishes edible ripe fruit and February-July fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Allow fruit to ripen. This drought-resistant shrub can form a hedge; prune to suit the space.

Buying advice: Confirm the exact species with a local indigenous-plant nursery; no tested regional cultivar or quantified yield selection is established.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Searsia natalensis

Kuni bush

1 existing; 1 proposed. Products: ripe dried fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Record ripe fruit picking and drying separately. No storage duration or fermentation recipe is assigned.

Buying advice: Confirm the exact species and local provenance; the SANBI assessment places it on stony fynbos/karroid slopes of the Northern and Western Cape.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Source: SANBI Red List: Searsia undulata

White milkwood

1 existing; 1 proposed. Products: ripe berries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: SANBI qualifies the food statement; it is about ripe berries, not latex, bark or roots. Fruit presence February-September does not establish ripe picking throughout.

Species identification in KwaZulu-Natal: The primary study records confusion with Bridelia micrantha. It does not supply a safe harvest calendar or kg/tree.

Shelter this coastal tree from direct salt-laden sea wind; SANBI reports salt burn despite its coastal habitat.

Buying advice: Ask a nursery to confirm the species and lawful provenance. Check current collection requirements before taking plants or products from existing trees.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, sideroxylon-inerme

Source: University of KwaZulu-Natal, Phytochemical and elemental studies of two indigenous medicinal plants

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Spiny monkey orange

1 existing; 1 proposed. Products: Ripe fruit pulp only.

First crop reference: 3-5 years. The review attributes this 3-5-year age to Avakoudjo et al.'s study in Benin, West Africa, rather than a South African bearing trial. Propagation method and local conditions must be confirmed before using it at a South African site; it is not a locally validated age or a yield-by-age curve. Actual timing depends on variety and site care.

Production reference: 40-100 kg/plant. The review reports 40-100 kg per tree stand without an annual period, mature age, matched cultivar or South African field conditions. Retain the original per-plant weight as context only: do not relabel it kg/year, spread it across months or treat it as a farm forecast.

Harvest interval with unverified South African geography: The review states August-December but does not identify a locality for that interval. Related monkey-orange literature discusses this interval in Zimbabwe and more than one Strychnos species; it does not establish a South African spinosa picking calendar. The former South Africa (general) window is withheld. Record the farm's own ripe-pulp months; seeds and unripe pulp are not ordinary food.

Geography of the review's bearing-age citation: The 2021 review cites this Benin study for its 3-5-year bearing-age statement. The cited source's geography is West Africa, not a South African orchard. Retain the review's numeric statement with this condition; no local annual yield curve is supplied.

Seeds and unripe fruit are toxic. Keep children away from unsupervised harvesting and preparation.

Buying advice: Confirm identity and appropriate handling before including it in a food plan; no named cultivar is sourced.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: FAO Ecocrop / ICRAF: Strychnos spinosa

Source: Undervalued Spiny Monkey Orange (Strychnos spinosa Lam.): An Indigenous Fruit for Sustainable Food-Nutrition and Economic Prosperity (peer-reviewed review, PMC8703348)

Source: Omotayo & Aremu (2021), Undervalued Spiny Monkey Orange, section 3

Source: Avakoudjo et al. (2020), Local Knowledge, Uses, and Factors Determining the Use of Strychnos spinosa Organs in Benin (West Africa), abstract

Waterberry

1 existing; 1 proposed. Products: Ripe berries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Flowering and fruit appearance, not a ripe picking calendar: The original PROTA account dates flowering and fruiting but does not date ripening. SANBI describes red to dark-purple ripe berries without ripe picking months. The November-March observation is retained as southern African phenology; local food-ready picking months remain unknown.

Water-loving: a dry site needs a dependable moisture source. Flowering for bees is not a honey-harvest forecast.

Buying advice: Choose a moist position and local provenance; no named cultivar is sourced.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: syzygium-cordatum

Source: Brink, M. (2008), PROTA4U: Syzygium cordatum, Growth and development

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Dune spinach

1 existing; 1 proposed. Products: young leafy tops and stems.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August-November flowering is not leaf availability.

Harvest young leafy tops; wash off sand before cooking.

Use well-drained coastal sand and keep divided plants moist until established.

Buying advice: Obtain correctly identified coastal planting material; record local growth and leaf-picking months because rainy seasons differ across South Africa.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Tetragonia decumbens

Kinkelbossie

1 existing; 1 proposed. Products: young shoots, cooked.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September-January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Harvest young shoots for cooking. Record food harvest separately from animal fodder.

Use a sunny, well-drained coastal, fynbos or Karoo position suited to this water-wise pioneer.

Buying advice: Obtain correctly identified local planting material; no named food cultivar or annual kg/plant benchmark is sourced.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

1 existing; 1 proposed. Products: flower buds.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July-September flowering interval does not identify exact unopened-bud picking months.

Record picking of young flower buds separately from open flowers.

Match the plant to well-drained coastal sandy conditions and its seasonal growth.

Buying advice: Use exact-species nursery identification; no first bud harvest age or food kg/plant benchmark is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Trachyandra ciliata

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Pigeonwood

1 existing; 1 proposed. Products: young leaves for cooked greens.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Match this fast pioneer to warm, moist eastern/northern conditions; protect damp, shaded seedlings.

Record young-leaf harvest separately from fruit consumed by wildlife.

Buying advice: Use correctly identified local planting material; no named leaf-food cultivar or quantified annual food output is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Trema orientalis

Wild garlic

1 existing; 1 proposed. Products: leaves and edible flowers.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

It tolerates prolonged drought but grows better with regular water. SANBI records its eastern South African distribution.

Buying advice: Ask for Tulbaghia violacea, rather than a plant identified only as wild garlic. Record local leaf and flower picking separately.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Eastern South African botanical account - edible flowers only: Jan-Apr.

Source: SANBI PlantZAfrica, tulbaghia-violacea

Wild medlar

Regional reference: Wild medlar - expected when established

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec

1 existing; 1 proposed. Products: Ripe fruit, fresh or dried.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Wider fruiting observations, not a ripe picking interval: This account separately distinguishes green unripe and brown ripe fruit, but its November-April biology sentence does not date that ripening change. The wider interval remains southern African phenology. SANBI's January-April description of ripe edible fruit supplies the food-ready regional reference.

Confirm local establishment conditions before relying on the regional fruiting reference.

Buying advice: Request locally appropriate fruiting stock; no named cultivar is sourced.

Regional reference: Wild medlar. South Africa (general; SA distribution given as Eastern Cape, Free State, Gauteng, KwaZulu-Natal, Limpopo, Mpumalanga, North West, Northern Cape). Planning reference when established; confirm local picking dates. No food quantity or first-year crop promised.

Source: SANBI PlantZAfrica: vangueria-infausta

Source: Vangueria infausta | PlantZAfrica (SANBI)

Source: Agroforestree Database 4.0 (Orwa et al. 2009), Vangueria infausta fact sheet, BIOLOGY section, p. 1

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Grape vine

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Flame Seedless year 4 - Hex River Valley enterprise budget: Year 4 commercial budget: 4,400 cartons/ha at 4.5 kg/carton = 19,800 kg/ha. The source does not provide matched vines/ha or row/vine spacing, so no kg/vine is derived. This is a budget assumption for one cultivar and region, not a generic full-bearing age or a local mapped-vine production record.

Match the vineyard variety to its production region, water and support system. The national supply season combines several regions.

Buying advice: DAFF industry reference: earlier Prime Seedless, Sugraone and Flame Seedless; later Dauphine and Crimson Seedless. SATI identifies Orange River choices including Prime, Thompson Seedless and Sugraone. Confirm local dates and stock; the industry sequence is not a KZN farm calendar.

Pollination: self fertile. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - South Africa - Northern Province(s) region (SATI production region; mild sub-tropical, earliest SA harvest): Nov-Dec.

Reference only - South Africa (overall, all regions combined): Nov-May.

Source: DAFF, Table grape market value chain profile (2012), p. 1

Source: SATI: Production regions

Source: South African Table Grape Industry (SATI) - 'About' page, production regions section

Source: South African Table Grape Industry (SATI) - 'About' page

Source: Zou, Massonnet, Minio et al., 'Multiple independent recombinations led to hermaphroditism in grapevine', Proceedings of the National Academy of Sciences (PNAS), Vol. 118, No. 15, e2023548118 (2021)

Source: Western Cape Department of Agriculture (Elsenburg), Agricultural Economic Services - 'Table Grape-Flame SL (Yr 4)' Enterprise Budget, Hex River Valley, Cape Winelands (updated 12 March 2024), p. 1

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Ginger

1 existing; 1 proposed. Products: fresh rhizomes; dried ginger recorded separately.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Use the planting date to record the rhizome harvest cycle. The Pretoria trial reference is conditional, not a fixed calendar across climates.

Record fresh rhizome and dried spice weights separately; keep edible ginger distinct from ornamental or African ginger species.

Buying advice: Ask a local supplier for healthy, identified Zingiber officinale planting rhizomes; no South African cultivar-specific yield curve is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Source: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Source: University of Pretoria author manuscript: Gatabazi et al. (2022)

Buffalo thorn

1 existing; 1 proposed. Products: ripe berries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: SANBI describes dry, slightly sweet ripe pulp. Fruit remaining on leafless trees March-August is persistence, not a verified six-month picking season.

SANBI records a wide range from frost-free coast to winter-frost regions. Its hooked and straight thorns matter when choosing access routes.

Buying advice: Ask for an identified buffalo-thorn plant suited to local conditions; the checked source supplies no named food cultivar.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, ziziphus-mucronata

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and ground.

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.