SYNTHETIC GRAPH SOURCE {COUNT} $&

Pele o reka kapa o qala

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Sehlopha sa tlelaemete sa dipatlisiso: Highveld. Sehlopha sena ha se keletso e netefaditsweng bakeng sa polasi ena.

Seo o se tlalehileng ka polasi ena

Metsi nakong ya komello

Ha ho a rekotwa. Qetella phuputso ya sebaka mme o netefatse metsi pele o kgetha moralo o hlokang disebediswa tse ngata.

Serame polasing ena

Se bonweng ka serame sebakeng ha se a rekotwa. Dikarolelano tsa kgwedi tsa sathelaete di ke ke tsa tiisa matsatsi a pele le a ho qetela a serame.

Letsatsi le phallo ya metsi mobung

Letsatsi: ha ho a rekotwa.

Phallo ya metsi: ha ho a rekotwa. Diphetho tsa teko ya mobu le ditlhoko tsa sejalo ka seng di ntse di hloka ho hlahlojwa.

Direkoto tsa selemo sa tlaleho - eseng ponelopele

Dikakaretso le dikgwedi tsa selemo sa tlaleho se fetileng di hlalosa se tlalehilweng. Ha di tshepise dijo moralong o tlang.

Original eggs {year} $&

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 240 eggs.

Tshebediso ya lelapa: 0 eggs. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: Php, Phu. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Fish

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: ha ho a rekotwa.

Tshebediso ya lelapa: ha ho a rekotwa. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: ha ho a rekotwa. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Meroho e meng

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 12 kg.

Tshebediso ya lelapa: 7 kg. Ho rekisitsweng: 3 kg.

Dikgwedi tsa kotulo tse tlalehilweng: Phe, Tsh. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo

Dikgoho: Laying hens (commercial)

Meaho e 1 e teng; e 0 e sisintsweng. Kgwebo e kgethilweng: mahe. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bakeng sa tlhahiso ya mahe, kopa dikgoho tse nyane tse entilweng tse haufi le ho behela ho mofani wa dikgoho tse behelang, ka direkoto tsa dilemo, phepo le ente.

Rera dijo tse leka-lekaneng, metsi a hlwekileng, moriti, moya le tshireletso ho dibatana. Phepelo ya mahe e itshetlehile ka dinonyana tse behelang, dilemo tsa tsona, bophelo le tlhokomelo; hoko feela ha e fane ka mahe.

Bokella mahe kgafetsa, boloka moo dihlaha di alwang ho hlwekile ho omme, mme o tshware/boloke mahe ka hloko bakeng sa tshebediso ya lelapa kapa thekiso.

Tshupiso ya nako: With a constant 14–16 hour light day, commercial hens keep laying at a high rate all year. Tshupiso ena ya taolo ya kgwebo ha e bolele phepelo ya mahe ya lelapa e lekanang ka tlhaho.

Tshupiso ya sehlahiswa sa pele: dibeke tse 21. Dilemo tsa nonyana ho tloha ho qhotswa; netefatsa mofuta le kgodiso le mofani. Dinonyana tse haufi le ho behela di qala hamorao potolohong ena.

Tshupiso ya tlhahiso: 407.1–428.6 mahe/ka kgoho e tshehadi e kentsweng ntlong (hen housed), potoloho ya ho behela ho fihlela dibeke tse 90. Table 4a's cumulative 'Eggs/hen housed' for five commercial strains; the same table gives 'Age at depop. weeks 90.0' for every strain. Commercial barn/cage average, not a free-range smallholder trial. Kept as the source gives it: a per-year rate would need a point-of-lay age from another document.

Tshupiso ya dilemo/mokgwa: SAPA 2024 commercial strain standards cumulative to 90 weeks: Dekalb Amberlink 429 eggs/hen housed, Dekalb Brown 428, Hy-Line Silver Brown 435, Hy-Line Brown 434 and Lohmann Brown Lite 413. These are full-cycle objectives, not eggs per calendar year or a smallholder forecast. Purchase age, actual onset of lay, surviving laying hens, feed, water and lighting need separate records.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

In free-range systems, hens must have access to the outdoor range for at least 8 hours a day in daylight.

Where individual nest boxes are used, at least one nest for every 8 hens.

Hens need at least 8 hours of continuous darkness in every 24-hour cycle to rest properly.

Floor housing needs litter on at least a third (33%) of the floor.

The outdoor free-range area must not be overstocked: no more than 5 birds per square metre of range.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Laying hens (commercial) (e tswela pele)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Mohlodi: SAPA Abridged Code of Practice: Commercial Layers, p. 1

Mohlodi: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Mohlodi: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Mohlodi: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: SAPA Industry Profile 2024, Table 4: commercial standards cumulative to 90 weeks, p. 39

Mohlodi: SAPA Layer Notes, Commercial Layers (training module, November 2013), p. 13

Mohlodi: SAPA Industry Profile 2022, Table 4a (Breed standards in laying hens), p. 39

Mohlodi: ARC Climate-Smart Agriculture Training Manual, Poultry production, p. 8

Mohlodi: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Mohlodi: FAO: Small-scale poultry production, egg marketing

Mohlodi: SAPA Layer Notes, Commercial Layers (training module, November 2013)

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Matata: Meat duck (Pekin / Muscovy, backyard)

Meaho e 1 e teng; e 0 e sisintsweng. Kgwebo e kgethilweng: nama. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Tshupiso ya sehlahiswa sa pele: dibeke tse 7–12. P44 = Pekin (7 weeks to slaughter); MR71 = Muscovy (10 weeks for ducks/hens, 12 weeks for drakes). Treated as 'time to first product' = age to slaughter, since this is a single-cycle meat enterprise (no repeat product like eggs). Non-South-African trial by Warsaw researchers, published in SAJAS; journal location does not establish local applicability.

Tshupiso ya tlhahiso: 2.3–2.9 kg ya boima bo phelang/ka nonyana nakong ya ho hlajwa. Non-SA (Guyana study reporting a citation to Steklenev 1990); no SA-specific absolute slaughter live-weight figure was found (see gaps). 2.93 kg = Pekin at 8-10 weeks, 2.29 kg = Muscovy at 8-10 weeks - matches this enterprise's two breeds. The same LRRD paper's own experiment (Table 2, intensive system, Guyana) measured live weight at slaughter (8 weeks) as: Pekin drakes 2.53 kg, Pekin hens 2.29 kg, Muscovy drakes 2.47 kg, Muscovy hens 1.51 kg - broadly consistent with, but not merged into, the cited range above since it is a separate dataset/table in the same source. No sourced eggs/duck/year figure was found for either breed (see gaps); the LRRD and SAJAS sources used here studied ducks reared only for meat.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

It is an offence to deny a duck food or water, or keep it somewhere lacking adequate space, ventilation, light or shelter from heat and cold.

Watch flocks for sudden deaths, ruffled feathers, swollen wattles/combs or a sudden drop in egg production - these are signs of avian influenza and should be reported at once.

Keep ducks away from wild birds and open water/feed that attracts them, and control visitors' access to duck housing, to reduce avian-influenza risk.

Ducks are one of the domesticated animal species listed in the Meat Safety Act, so its rules on registered abattoirs and hygiene apply to them.

Ducks slaughtered to be sold must be slaughtered at a registered abattoir; slaughter for the farmer's own consumption is exempt but that meat may not be sold.

At the abattoir, ducks must be handled humanely during loading, transport, offloading, housing, immobilising and killing, under the Animals Protection Act.

Ducks are birds/poultry, so an outbreak of avian influenza in a duck flock is a controlled animal disease under the Animal Diseases Act and must be reported to the state vet immediately.

Mohlodi: Animals Protection Act 71 of 1962 (consolidated text, s.2(1)(b)-(c)), p. 2

Mohlodi: Western Cape Department of Agriculture, Avian Influenza (Bird Flu) factsheet, p. 2

Mohlodi: Meat Safety Act 40 of 2000, Schedule 1 (Animals to which this Act applies), p. 16

Mohlodi: Meat Safety Act 40 of 2000, s.7, p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.9, p. 7

Mohlodi: Damaziak et al. (2014), Influence of housing system on the growth and histological structure of duck muscles, S. Afr. J. Anim. Sci. 44(2): Poland trial

Mohlodi: Livestock Research for Rural Development 18(11) 2006 - Solomon et al. (Guyana; not an SA figure), citing Steklenev 1990

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Letangwana: Tilapia (Mozambique tilapia, small pond or tank)

Meaho e 1 e teng; e 0 e sisintsweng. Kgwebo e kgethilweng: tlhapi. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Tshupiso ya sehlahiswa sa pele: dibeke tse 43.5–52.2. 10 months x 4.35 = 43.5 weeks; 12 months x 4.35 = 52.2 weeks. This is Mozambique Tilapia (O. mossambicus) grow-out time to harvestable size; the source separately states Nile Tilapia (O. niloticus) reaches harvestable maturity faster, in 6-10 months (= 26.1-43.5 weeks) — not used here as it is a different species with a different legal status (see 'legal').

Tshupiso ya tlhahiso: 800 g ya boima nakong ya kotulo/ka tlhapi. Table 8-3 gives the Mozambique Tilapia's harvestable size as an approximate single figure ('+- 800g'), not a range, so [800, 800] is used rather than inventing a spread. The same table cell wraps '10-12' across a line break in the PDF ('10-\n12'), kept verbatim. The table's Nile Tilapia row gives the same approximate weight (+-800g) reached faster, in 6-10 months — not used here because this dossier is framed around the indigenous, legally unrestricted Mozambique Tilapia per task instruction; see 'legal' for why Nile Tilapia is treated separately. No SA source was found giving a feed-conversion-based or per-hectare yield figure specific to O. mossambicus that could be converted to this per-fish unit without stitching numbers across documents, so outputUnit follows the source's own unit (grams harvest weight per fish) rather than being converted.

Tshupiso ya dilemo/mokgwa: ARC makes tilapia suitability depend on water temperature: summer grow-out can be suitable anywhere in South Africa when water exceeds 20°C; growth is poor below 18°C and winter water below 13°C can kill farmed tilapia. The Highveld reference is a summer crop; tunnels can widen the range. Measure pond water and confirm stocking species and dates. Air climate or a mapped pond does not prove harvest months, fish numbers or food kg.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

The indigenous Mozambique Tilapia (O. mossambicus) is not a listed invasive species and may be farmed without an alien-species permit; the faster-growing Nile Tilapia (O. niloticus) is.

Nile Tilapia (Oreochromis niloticus) is a nationally listed invasive fish species: Category 1b inside National Parks, Provincial Reserves, Mountain Catchment Areas and Forestry Reserves, and Category 3 in all other areas.

Outside protected areas Nile tilapia is Category 3: it may remain in prescribed areas or provinces, but further propagation or trade is prohibited.

Nile Tilapia has been listed under the NEMBA Alien and Invasive Species Regulations as requiring a permit to produce.

Producing an alien or invasive species (such as Nile Tilapia) is prohibited in areas designated under the Protected Areas Act and the Fish Sanctuary Act; a permit is required to produce it elsewhere.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Letangwana: Tilapia (Mozambique tilapia, small pond or tank) (e tswela pele)

Producing tilapia in South Africa requires a water use licence under the National Water Act.

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), Table 8-3 (Mozambique vs. Nile Tilapia), p. 69

Mohlodi: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), List 7 (National List of Invasive Fresh-water Fish Species), p. 32

Mohlodi: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), p. 3

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 69

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 74

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 75

Mohlodi: ARC Aquaculture Production App: General FAQ, Can I grow tilapia on the Highveld?

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dipodi: Dairy goat (Saanen / Toggenburg / British Alpine)

Meaho e 1 e teng; e 0 e sisintsweng. Kgwebo e kgethilweng: lebese. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Rera ho ya ka ho tswala ha nnete ha manamane/madinyane le nako ya lebese. Diphoofolo tse sa hlahiseng lebese ha di fane ka lona.

Nako ya sehlahiswa sa pele: ha ho mohlodi bakeng sa kgwebo ena.

Tshupiso ya tlhahiso: 775.4–1073.4 kg ya lebese/ka podi e tshehadi/ka nako ya ho hlahisa lebese ya matsatsi a 305. Lowest (>=7th parity) and highest (3rd parity) group means for milk yield per 305-day standardised lactation in 16,407 analysed South African ARC Saanen lactation records, 1955–2018, mostly intensive/semi-intensive stud herds. Original PDF Table 3 labels the last group >=7th; HTML renders it >7th. Kept in kg, without litres conversion, annualisation or yield-by-age interpolation.

Tshupiso ya dilemo/mokgwa: South African Saanen reference by lactation number, not age in years: mean milk mass per standardised 305-day lactation was 947.3 kg at first parity, 1043.2 at second, 1073.4 at third, 1020.3 at fourth, 972.8 at fifth, 901.6 at sixth and 775.4 at >=7th. These ARC milk-recorded groups were mostly intensive/semi-intensive stud herds. Record actual kidding date, lactation number and collected milk; do not divide these totals evenly into months.

Tshupiso ya dilemo/mokgwa: At Putukwane near Polokwane, 179 lactating does under extensive feeding with irregular supplementation were observed for 210 days after kidding. Mean milk mass was 1.45 kg/day for Saanen, 0.75 for British Alpine and 0.56 for Toggenburg during observed lactation. Early lactation exceeded mid and late lactation; March–April kidding outperformed October–November in this cohort. These conditions differ from stud-herd records. Discuss adapted stock and feed locally; these are not litres/day or year-round promises.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

It is an offence to keep a goat somewhere with inadequate space, ventilation, light or shelter from heat, cold or weather, or to confine/tether it in a way that causes unnecessary suffering.

It is an offence to starve, under-feed, or deny water or food to a goat.

An owner must not let a goat stay in a dirty or parasite-infested condition, and must get veterinary or other treatment for a sick, injured or diseased animal when able to.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: veterinary services enforce isolation and movement restrictions, and a suspected case must be reported.

Inside a declared FMD-affected zone, goats and their products (including milk) cannot be moved into, out of, or within the zone without a state veterinary movement permit and compliance with its conditions.

Every goat owner must register an official identification mark and mark their goats with it, under the Animal Identification Act.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dipodi: Dairy goat (Saanen / Toggenburg / British Alpine) (e tswela pele)

A goat may only be slaughtered for sale at a registered abattoir under the Meat Safety Act (relevant when culled dairy does or male kids are sold for meat); slaughtering for the farmer's own consumption, or for cultural/religious purposes, is exempt but that meat may not then be sold.

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(b), p. 1

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(e), p. 1

Mohlodi: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Mohlodi: Animal Identification Act 6 of 2002, s.7(1), p. 4

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: Malemela, Nemutandani & Tada (2024), South African Saanen goats, Table 3, p. 625

Mohlodi: Norris et al. (2011), Milk Production of Three Exotic Dairy Goat Genotypes in Limpopo Province, South Africa: methods, results and Table 2

Mohlodi: Malemela, Nemutandani & Tada (2024), South African Saanen goats: Table 3, 305-day milk yield by parity, p. 625

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dipodi: Indigenous veld / communal goat (e.g. Nguni, Tankwa, Xhosa lob-ear)

Meaho e 1 e teng; e 0 e sisintsweng. Kgwebo e kgethilweng: nama. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Tshupiso ya sehlahiswa sa pele: dibeke tse 69.6–78.3. This is age at first KIDDING (birth of first kids), not first weaned kids. 16 months x 4.35 = 69.6 weeks; 18 months x 4.35 = 78.3 weeks, from doe birth to her giving birth for the first time. Adding the source's own ~150-day (~5-month, ~21.75-week) weaning assumption would put first WEANED kids at roughly 91-100 weeks of doe age, but that compound figure is not shown as the primary value since the field is documented as 'age/time from start (hatch, birth) to first product' and first kidding is the clearer, directly-stated milestone; the weaning offset is noted here for transparency.

Tshupiso ya dilemo/mokgwa: The 2016 government goat manual distinguishes indigenous, Boer, Savanna and Kalahari Red meat-purpose stock from milk/fibre breeds, with breeding objectives and individual birth, weaning, weight and sale records. Uncontrolled communal mating can produce births at any time; that does not establish meat or weaned-kid availability all year. Choose stock for the local system and record survival and product stage before forecasting quantities.

Tshupiso ya dilemo/mokgwa: In KZN Highland Sourveld smallholder systems, winter and early-spring feed shortage can restrict goats. Autumn-kidding does feed kids through winter; spring-kidding does face late-pregnancy feed shortage, especially with late rains. Plan local winter feed and monitor weights with extension support; this is regional care context, not kg or a universal kidding calendar.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

It is an offence to keep a goat somewhere with inadequate space, ventilation, light or shelter from heat, cold or weather, or to confine/tether it in a way that causes unnecessary suffering.

It is an offence to starve, under-feed, or deny water or food to a goat.

An owner must not let a goat stay in a dirty or parasite-infested condition, and must get veterinary or other treatment for a sick, injured or diseased animal when able to.

Winter feed shortage on communal sourveld is a known welfare/production risk: does that kid in autumn must feed kids through winter, and does that kid in spring suffer poor nutrition during late pregnancy, so winter feed supplementation matters.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: veterinary services enforce isolation and movement restrictions, and a suspected case must be reported.

Inside a declared FMD-affected zone, goats and their products cannot be moved into, out of, or within the zone without a state veterinary movement permit and compliance with its conditions.

Every goat owner, including communal/smallholder owners, must register an official identification mark and mark their goats with it, under the Animal Identification Act.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dipodi: Indigenous veld / communal goat (e.g. Nguni, Tankwa, Xhosa lob-ear) (e tswela pele)

A goat may only be slaughtered for sale at a registered abattoir under the Meat Safety Act; slaughtering it for the farmer's own consumption, or for cultural/religious purposes (including ceremonial slaughter, common with indigenous goats), is exempt from that rule, but that meat may not then be sold.

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(b), p. 1

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(e), p. 1

Mohlodi: Letty, B. & Madiba, S., KZN DARD. Constraints to Small-Scale Goat Production in the Highland Sourveld Area of KwaZulu-Natal, p. 3

Mohlodi: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Mohlodi: Animal Identification Act 6 of 2002, s.7(1), p. 4

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: DAFF (2016), Goat Training Manual: breeds p2, records pp4–5, uncontrolled mating section 2.4.6 p7, p. 7

Mohlodi: KZN DARD: Goat Production in KwaZulu-Natal, Highland Sourveld research and winter nutrition

Mohlodi: Webb, E.C. & Mamabolo, M.J. (2004). Production and reproduction characteristics of South African indigenous goats in communal farming systems. S. Afr. J. Anim. Sci. 34(5):236-239

Mefuta ya dijalo e lokelang ho hlahlojwa sebakeng seo

Boloka lebitso la mofuta oo o o kgethang ka nnete. Manane a makgutshwane a dipatlisiso a hloka netefatso bakeng sa nako ya sebaka, metsi le mobu.

Tomatoes

Mofuta o rekotilweng: ha ho eso kgethwe.

Lenane le lekgutshwane la dipatlisiso bakeng sa maemo a lehodimo a tshwanang: Rodade.

Mohlodi: Rodade, Sakata product page

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.