SYNTHETIC GRAPH SOURCE {COUNT} $&

Musi ni sa athu u renga kana u thoma

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Tshigwada tsha kilima tsha ṱhoḓisiso: Fhethu ha u lima a hu athu u ṱalutshedzea. Tshigwada itshi a si nyeletshedzo yo khwaṱhisedzwaho ya bulasi iḽi.

Zwe na vhiga nga bulasi iḽi

Maḓi nga khalaṅwaha yo oma

A zwo ngo rekhodwa. Fhedzisani ṱhoḓisiso ya fhethu ni khwaṱhisedze maḓi ni sa athu u nanga pulane ine ya ṱoḓa zwishumiswa zwinzhi.

Murotholo kha bulasi iḽi

Zwe zwa vhonwa nga murotholo wa henefho a zwo ngo rekhodwa. Zwikati zwa ṅwedzi zwa sathelaithi a zwi koni u khwaṱhisedza maḓuvha a murotholo wa u thoma na wa u fhedza.

Ḓuvha na u bva ha maḓi mavuni

Ḓuvha: a zwo ngo rekhodwa.

U bva ha maḓi: a zwo ngo rekhodwa. Mvelelo dza u ṱola mavu na ṱhoḓea dza tshiliṅwa tshiṅwe na tshiṅwe zwi kha ḓi ṱoḓa u ṱolwa.

Rekhodo dza ṅwaha wa u vhiga - hu si mbonelaphanḓa

Zwivhalo na miṅwedzi zwa ṅwaha wo fhiraho wa u vhiga zwi ṱalutshedza zwo vhigwaho. A zwi fulufhedzisi zwiḽiwa kha pulane i ḓaho.

Original eggs {year} $&

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: 240 eggs.

Zwo shumiswaho muṱani: 0 eggs. Zwo rengiswaho: a zwo ngo rekhodwa.

Miṅwedzi ya khaṇo yo vhigwaho: Fulw, Fwan. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

Fish

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: a zwo ngo rekhodwa.

Zwo shumiswaho muṱani: a zwo ngo rekhodwa. Zwo rengiswaho: a zwo ngo rekhodwa.

Miṅwedzi ya khaṇo yo vhigwaho: a zwo ngo rekhodwa. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

Miṅwe miroho

Tshivhalo tsha ṅwaha tsho vhigwaho nga mulimi tsha 2025: 12 kg.

Zwo shumiswaho muṱani: 7 kg. Zwo rengiswaho: 3 kg.

Miṅwedzi ya khaṇo yo vhigwaho: Phan, Nyen. Izwi ndi zwe zwa vhonwa kha ṱhoḓisiso, hu si mbonelaphanḓa ntswa ya vhubveledzi.

Ṱolani yuniti na zwivhalo zwo rekhodwaho ni sa athu u shumisa mutaladzi uyu.

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṇu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani

Dogwood

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: muthetshelo wa maṱari musi hu tshi khou ukwa — u shumiswa nga muḓivhi.

U sedzulusa tshumiso ya zwiḽiwa: khwaṱhisedzani vhuvha, tshipiḓa tshi ḽiwaho na u lugisela ho teaho na mueletshedzi wa henefho a re na ndalukanyo ni sa athu u katela tshimela itshi kha zwiḽiwa zwa muṱa. A tshi katelwi kha zwivhalo zwa zwiḽiwa zwo ḓoweleaho na mishumo ya u kaṋa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Khwaṱhisedzani vhuvha nga vhuḓalo na u lugisela zwiḽiwa ho dzudzanywaho na mueletshedzi wa henefho a re na tshenzhelo.

Mafhungo a South Africa a ṱalusa u aluwa mathungoni a milambwana na fhethu ho sendamaho ha maḓaka na u konḓelela murotholo wa gwanda; mitshelo yatsho miṱuku i ṱaluswa sa zwiḽiwa zwa zwiṋoni.

Nyeletshedzo ya u renga: Rengani zwimela zwa lushaka lwo teaho zwa tshumiso yo dzudzanywaho yo khwaṱhisedzwaho. Ni songo dzhia vhuṱanzi ha mushonga kana ha mitshelo ya zwipuka zwa ḓaka sa mbonelaphanḓa ya zwiḽiwa zwiswa zwa muṱa.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Tshiko: SANBI PlantZAfrica: Rhamnus prinoides

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Baboon grape

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo ya berry yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: The source gives a ripe-berry interval. This does not establish stored jam months or safe shelf life.

Ṋeani tshitsigiso tsha u gonyelela ni lange u phaḓalala hatsho hune tshikhala tsha vha tshiṱuku. SANBI i tshi rekhoda kha zwikwara zwa muṱavha zwa khunzi na mathungoni a maḓaka u bva Western Cape u ya KwaZulu-Natal na Mpumalanga.

Nyeletshedzo ya u renga: Khwaṱhisedzani Rhoicissus digitata; dziṅwe tshaka dza wild-grape dzi na ṱhalutshedzo dzo fhambanaho nahone a dzi koni u ṋea maḓuvha ayo.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu athu u khwaṱhisedzwa tshifhinga tshithihi tsha lushaka lwa henefho. Khalaṅwaha dza ndaedzi dzi re afho fhasi a si mbonelaphanḓa.

Ndaedzi fhedzi — tshiko tsha botaniki tsha Western/Eastern Cape, KwaZulu-Natal na Mpumalanga — mitshelo miṱuku yo vhibvaho: Ṯhaf–Ṯhan.

Tshiko: SANBI PlantZAfrica, rhoicissus-digitata

Bushman's grape

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo miṱuku i ḽiwaho; midzi a i katelwi.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Ṋeani fureme ya u gonyelela kana tshiṅwe tshitsigiso. Zwimela zwiṱuku zwi ṱoḓa maḓi u swika zwi tshi khwaṱha; SANBI i ṱalusa u konḓelela gomelelo nga murahu na u shumiswa ḓuvhani u ya murunzini u sa fhelelaho.

Nyeletshedzo ya u renga: Humbelani Rhoicissus tridentata ni rekhode miṅwedzi ya mitshelo yo vhibvaho ya henefho. Ni songo shumisa tshumiso ya midzi sa mushonga kana kha zwipuka sa vhuṱanzi ha zwiḽiwa.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica, rhoicissus-tridentata

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṯhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Rosemary

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari maswa kana o omiswaho a u naka zwiḽiwa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Shumisani mavu a bvisaho maḓi zwavhuḓi na maḓi o linganaho musi zwimela zwi tshi khou thoma u khwaṱha; iledzani u ḓala ha maḓi na u bola ha midzi.

Rekhodani maṱari maswa, tshinaki tsho omiswaho na mapfura a munukhelelo wa tshimela (essential oil) sa zwibveledzwa zwo fhambanaho.

Nyeletshedzo ya u renga: Wanani zwimela zwa rosemary zwi re na mutakalo zwo topolwaho na nyeletshedzo ya henefho ya u zwi alusa; a hu na lushaka lwa dzingu lwa khaṋo ya maṱari a u bika lwo khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: DAFF: Rosemary production brochure (2012), p. 2

Karoo boer-bean

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mbeu dzi ḽiwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Vhudzulo hatsho ha mvelo vhu katela mathungo o omaho a milambwana ngei Little Karoo na fhethu ho omaho u fhira huṅwe ha Eastern Cape na tshipembe tsha Western Cape.

Dzi-pod (zwikwama zwa mbeu) dzi phaḓaladza mbeu musi dzo oma. Rekhodani u kuvhanganya na u shumisa mbeu ha henefho; izwi a si khaṋo yo kaliwaho ya zwiḽiwa.

Nyeletshedzo ya u renga: Humbelani Schotia afra var. afra, vhuvha ha tshimela ha botaniki ho khwaṱhisedzwa. Zwine tshiko tsha amba nga ha murunzi nga miṅwaha miṱanu a si vhukale ha zwiḽiwa zwa u thoma.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu athu u khwaṱhisedzwa tshifhinga tshithihi tsha lushaka lwa henefho. Khalaṅwaha dza ndaedzi dzi re afho fhasi a si mbonelaphanḓa.

Ndaedzi fhedzi — tshiko tsha botaniki tsha Little Karoo na Cape ho omaho nga maanḓa — ndaedzi ya vhubveledzi ha mbeu i ḽiwaho: Shun–Fulw.

Tshiko: SANBI PlantZAfrica, schotia-afra-var-afra

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Karee

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo ya berry yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: September–January describes fruit production; exact ripe-picking dates are not stated. Do not count leaves used as livestock fodder as household food.

SANBI i rekhoda muri uyu wa murunzi u konḓelelaho murotholo wa gwanda na gomelelo kha nyimele dza Karoo na Kalahari.

U ya nga SANBI, maṱari atsho manzhi a nga tshinya muthetshelo wa mafhi a zwifuwo zwa mafhi; izwi a si themendelo ya mukovhe wa zwiḽiwa.

Nyeletshedzo ya u renga: Vhudzisani nga zwimela zwa tshiduna na tshisadzi arali mitshelo i yone yo dzudzanywaho. U phaḓalala ha mvelo kha tshiko a hu kateli KwaZulu-Natal; u konḓelela nyimele nnzhi a si khwaṱhisedzo ya mitshelo ya henefho.

U phaḓaladzwa ha muvhere: zwimela zwa tshinna na zwa tshisadzi zwo fhambanaho. Ṱolani ṱhoḓea dza lushaka lwo nangwaho na nesari.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica, searsia-lancea

Glossy currant

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Currant fruit food use and phenology: The primary account establishes edible ripe fruit and October–June fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Kaṋani mitshelo yo vhibvaho ni rekhode u omisa nga thungo.

Tshiṱaka itshi tshi tea ngade dza khunzi dzi songo tsireledzwaho na u khwaṱhisa zwikwara zwa muṱavha; khwaṱhisedzani vhubvo ha henefho.

Nyeletshedzo ya u renga: Humbelani neseḽi ya zwimela zwa henefho uri ni wane lushaka lwo teaho na zwimela zwa henefho. A hu na lushaka lwa zwiḽiwa lwa dzingu lu bveledzaho khaṋo nnzhi lwo khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Searsia lucida

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Natal currant

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Edible Natal currant fruit and local phenology: The primary account establishes edible ripe fruit and February–July fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Tendelani mitshelo i vhibve. Tshiṱaka itshi tshi konḓelelaho gomelelo tshi nga vha luhura; tshi tumuleni uri tshi tea tshikhala.

Nyeletshedzo ya u renga: Khwaṱhisedzani lushaka lwo teaho na neseḽi ya henefho ya zwimela zwa henefho; a hu na lushaka lwa dzingu lwo lingwaho kana nango ya khaṋo yo kaliwaho nga tshivhalo yo khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Searsia natalensis

Kuni bush

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo yo vhibvaho yo omiswaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Rekhodani u kaṋa mitshelo yo vhibvaho na u omisa nga thungo. A hu na tshifhinga tsha u vhulunga kana nḓila ya u vhila yo ṉewaho.

Nyeletshedzo ya u renga: Khwaṱhisedzani lushaka lwo teaho na vhubvo ha henefho; tsedzuluso ya SANBI i tshi vhea fhethu ho sendamaho ha matombo ha fynbos/karroid ha Northern Cape na Western Cape.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Tshiko: SANBI Red List: Searsia undulata

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

White milkwood

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo ya berry yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: SANBI qualifies the food statement; it is about ripe berries, not latex, bark or roots. Fruit presence February–September does not establish ripe picking throughout.

Species identification in KwaZulu-Natal: The primary study records confusion with Bridelia micrantha. It does not supply a safe harvest calendar or kg/tree.

Tsireledzani muri uyu wa khunzi kha muya wa lwanzhe wo ḓalaho muṋo u u vhudzulaho wo tou livha; SANBI i vhiga u swa nga muṋo naho vhudzulo hawo vhu ha khunzi.

Nyeletshedzo ya u renga: Humbelani neseḽi uri i khwaṱhisedze lushaka na vhubvo vhu tevhelaho mulayo. Ṱolani ṱhoḓea dza zwino dza u kuvhanganya musi ni sa athu u dzhia zwimela kana zwibveledzwa kha miri i re hone.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica, sideroxylon-inerme

Tshiko: University of KwaZulu-Natal, Phytochemical and elemental studies of two indigenous medicinal plants

Dune spinach

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: ṱhodzi ntswa dzi re na maṱari na tsinde.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Kaṋani ṱhodzi ntswa dzi re na maṱari; ṱanzwani muṱavha musi ni sa athu u bika.

Shumisani muṱavha wa khunzi u bvisaho maḓi zwavhuḓi ni dzule ni tshi ṉea zwimela zwo khethekanywaho muṋa u swika zwi tshi khwaṱha.

Nyeletshedzo ya u renga: Wanani zwimela zwa u ṱavha zwa khunzi zwo topolwaho zwavhuḓi; rekhodani u aluwa na miṅwedzi ya u kaṋa maṱari ya henefho ngauri zwifhinga zwa mvula zwi a fhambana u mona na South Africa.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Tetragonia decumbens

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Kinkelbossie

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: matavhi maṱuku, o bikwa.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Kaṋani matavhi maṱuku a u bika. Rekhodani khaṋo ya zwiḽiwa nga thungo na furu ya zwifuwo.

Shumisani fhethu ha ḓuvhani hu bvisaho maḓi zwavhuḓi ha khunzi, fynbos kana Karoo ho teaho tshimela itshi tsha u thoma u mela fhethu tshi shumisaho maḓi nga vhulondi.

Nyeletshedzo ya u renga: Wanani zwimela zwa u ṱavha zwa henefho zwo topolwaho zwavhuḓi; a hu na lushaka lwa zwiḽiwa lu re na dzina kana tshikalo tsha ṅwaha tsha kg/plant tshi bvaho tshikoni.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Tetragonia fruticosa

Veldkool

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maluvha a sa athu u vulea.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Rekhodani u kaṋa maluvha maṱuku a sa athu u vulea nga thungo na maluvha o vuleaho.

Fanyisani tshimela na nyimele dza muṱavha wa khunzi u bvisaho maḓi zwavhuḓi na u aluwa hatsho ha zwifhinga.

Nyeletshedzo ya u renga: Shumisani u topola lushaka lwo teaho ha neseḽi; a hu na vhukale ha khaṋo ya u thoma ya maluvha a sa athu u vulea kana tshikalo tsha zwiḽiwa tsha kg/plant tsho khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Trachyandra ciliata

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Pigeonwood

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari maṱuku a muroho wo bikwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Fanyisani tshimela itshi tsha u thoma u mela fhethu tshi aluwaho nga u ṱavhanya na nyimele dzi fhisaho dzi re na muṋa dza vhubvaḓuvha/devhula; tsireledzani zwimela zwiṱuku zwi re na muṋa zwi re murunzini.

Rekhodani khaṋo ya maṱari maṱuku nga thungo na mitshelo i ḽiwaho nga zwipuka zwa ḓaka.

Nyeletshedzo ya u renga: Shumisani zwimela zwa u ṱavha zwa henefho zwo topolwaho zwavhuḓi; a hu na lushaka lu re na dzina lwa maṱari a zwiḽiwa kana mbuelo ya zwiḽiwa ya ṅwaha yo kaliwaho nga tshivhalo yo khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica: Trema orientalis

Wild garlic

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: maṱari na maluvha a ḽiwaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

Tshi konḓelela gomelelo ḽa tshifhinga tshilapfu fhedzi tshi aluwa khwine nga maḓi a tshifhinga tshoṱhe. SANBI i rekhoda u phaḓalala hatsho vhubvaḓuvha ha South Africa.

Nyeletshedzo ya u renga: Humbelani Tulbaghia violacea, hu si tshimela tsho topolwaho fhedzi sa wild garlic. Rekhodani u kaṋa maṱari na maluvha ha henefho nga thungo.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu athu u khwaṱhisedzwa tshifhinga tshithihi tsha lushaka lwa henefho. Khalaṅwaha dza ndaedzi dzi re afho fhasi a si mbonelaphanḓa.

Ndaedzi fhedzi — tshiko tsha botaniki tsha vhubvaḓuvha ha South Africa — maluvha a ḽiwaho fhedzi: Phan–Lamb.

Tshiko: SANBI PlantZAfrica, tulbaghia-violacea

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Ginger

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: tsinde ntswa dzi aluwaho fhasi ha mavu; ginger yo omiswaho i rekhodiwa nga thungo.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Shumisani datumu ya u ṱavha u rekhoda mutevhe wa khaṋo ya rhizome, tsinde i aluwaho fhasi ha mavu. Tshikombo tsha mulingo wa Pretoria tshi na nyimele, a si khaḽenda i sa shandukiho kha kilima dzo fhambanaho.

Rekhodani zwileme zwa rhizome ntswa na tshinaki tsho omiswaho nga thungo; fhambanyani ginger i ḽiwaho na tshaka dza ginger dza u khavhisa kana dza African ginger.

Nyeletshedzo ya u renga: Humbelani muṋetshedzi wa henefho dzi-rhizome dza u ṱavha dza Zingiber officinale dzi re na mutakalo dzo topolwaho; a hu na girafu ya khaṋo ya lushaka lwo khetheaho lwa South Africa yo khwaṱhisedzwaho.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Tshiko: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Tshiko: University of Pretoria author manuscript: Gatabazi et al. (2022)

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Zwine zwa nga nangwa zwa bulasi - zwi khou bvela phanḓa

Shumisani zwe na zwi vhona bulasini na mitevhe mipfufhi ya ṱhoḓisiso. Khwaṱhisedzani zwine na nanga henefho musi ni sa athu u renga.

Mitshelo, nḓuhu na zwiḽiwa zwa hayani (zwi khou bvela phanḓa)

Buffalo thorn

Zwi re hone: 1; zwo dzudzanywaho: 1. Zwibveledzwa: mitshelo ya berry yo vhibvaho.

Tshifhinga tsha khaṋo ya u thoma: a tsho ngo wanala tshikoni; ni songo ḓitika nga zwimela zwi sa athu bva u ṱavhiwa uri ni wane zwiḽiwa nga u ṱavhanya.

Tshipiḓa tshi ḽiwaho na tshumiso ya henefho: SANBI describes dry, slightly sweet ripe pulp. Fruit remaining on leafless trees March–August is persistence, not a verified six-month picking season.

SANBI i rekhoda u phaḓalala hu angalalaho u bva khunzini i si na murotholo wa gwanda u ya madzinguni a murotholo wa gwanda wa vhuriha. Mipfa yatsho yo khotheaho na yo tou tswititi ndi ya ndeme musi hu tshi nangwa nḓila dza u swikelela.

Nyeletshedzo ya u renga: Humbelani tshimela tsha buffalo-thorn tsho topolwaho tshi teaho nyimele dza henefho; tshiko tsho ṱolwaho a tshi ṋei lushaka lwa zwiḽiwa lu re na dzina.

Khalaṅwaha ya u kaṋa yo lavhelelwaho: a hu na tshiko tsha miṅwedzi yo tiwaho ya dzingu. Rekhodani maḓuvha a u ṱavha na a u kaṋa henefho.

Tshiko: SANBI PlantZAfrica, ziziphus-mucronata

Tshaka dza zwiliṅwa dza u sedzulusa henefho

Vhulungani dzina ḽa lushaka lune na nanga zwa vhukuma. Mitevhe mipfufhi ya ṱhoḓisiso i ṱoḓa khwaṱhisedzo ya khalaṅwaha ya henefho, maḓi na mavu.

Tomatoes

Lushaka lwo rekhodwaho: a lu athu u nangwa.

Kilima ya fhethu hafha a i athu u ṱalutshedzea, ngauralo a hu buliwi mutevhe mupfufhi wa henefho. Vulani garaṱa ya tshiliṅwa u vhona zwiko na u rekhoda zwe na nanga.

A i athu ṱolwa nga muḓivhi wa zwa vhulimi — ndi tsumbanḓila ya u pulana, a si ndaela. I sumbedzeni muṱuṱuwedzi waṋu wa zwa vhulimi, ni ite ndingo ya mavu musi ni sa athu shumisa tshelede kha mavu.
Ṱhalutshedzelo ya mvetamveto: a i athu u sedzwa nga muthu a ambaho luambo zwavhuḓi. Sedzani tshiko tsha Luisimane phanḓa ha u tevhedza zwine zwa ambiwa.