SYNTHETIC GRAPH SOURCE {COUNT} $&

Before you buy or start

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Research climate group: Growing area not resolved. This grouping is not a validated recommendation for this farm.

What you reported about this farm

Water through the dry season

Not recorded. Complete the site survey and confirm water before choosing an intensive plan.

Frost on this farm

Local frost observations are not recorded. Satellite monthly averages cannot establish first and last frost dates.

Sun and drainage

Sun: not recorded.

Drainage: not recorded. Soil-test results and crop-specific needs still need checking.

Reporting-year records - not a forecast

Past reporting-year totals and months describe what was reported. They do not promise food in the coming plan.

Original eggs {year} $&

Farmer-reported annual total for 2025: 240 eggs.

Household use: 0 eggs. Sold: not recorded.

Reported harvest months: Jun, Jul. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Fish

Farmer-reported annual total for 2025: not recorded.

Household use: not recorded. Sold: not recorded.

Reported harvest months: not recorded. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

Other vegetables

Farmer-reported annual total for 2025: 12 kg.

Household use: 7 kg. Sold: 3 kg.

Reported harvest months: Jan, Dec. These are survey observations, not a new production forecast.

Check the recorded unit and totals before using this row.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods

Kriedoring

1 existing; 1 proposed. Products: food product not verified.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Exact identity; no confirmed household food part: The checked primary assessment establishes exact identity and South African distribution only. It does not establish edible parts, food-use preparation, picking dates or yield; claims for a different Lycium species are not transferred.

Record the exact identity and obtain local expert food-use advice before counting it as household food.

Buying advice: Confirm exact species with a qualified indigenous nursery. Do not substitute genus-level berry claims for identification.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI Red List: Lycium cinereum

Kriedoring

1 existing; 1 proposed. Products: berries - food use requires specialist review.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Documented berry-use caution: This explicit caution prevents routine household-food forecasting. Sporadic all-year fruiting is not promoted to an edible-food availability calendar.

Keep this as an identified hedge/inventory plant. Seek expert local food-use review before treating berries as household food.

Buying advice: Confirm the exact species and intended hedge use with a local indigenous nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Lycium ferocissimum

Ice plant

1 existing; 1 proposed. Products: leaves and fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Annual edible leaf and fruit crop: This is an annual, not a fruit tree. The account describes boiled leaves as a spinach substitute and pickling. It supplies no food harvest duration, kg/plant or exact leaf calendar.

Record planting batches and harvested leaves/fruit separately.

Match this frost-sensitive plant to dry saline/coastal conditions. Its accumulated salts can return to the soil as plants die.

Buying advice: Obtain exact-species seed or plants and assess the bed soil; no regional food cultivar or annual food yield is verified.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Mesembryanthemum crystallinum

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Coastal red milkwood

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The source describes bright orange-red mature fruit. It does not give a picking calendar.

SANBI records this tree in east-coast dune forest from Eastern Cape through KwaZulu-Natal, with tolerance of salty sea spray.

Buying advice: Confirm Mimusops afra and lawful nursery provenance. Check current protected-tree collection requirements before taking plants or products from existing trees.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, mimusops-afra

Nkaka

1 existing; 1 proposed. Products: traditionally prepared leaves/fruit - specialist review needed.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible-part and safety review: SANBI reports culinary uses but also toxic constituents and poisoning concerns. A preparation report is not a complete safe household recipe.

Retain this climber in mapped inventory, but do not count it as normal household food until a qualified local specialist confirms identity, edible part and safe preparation.

SANBI records it across South Africa except Western Cape, in varied habitats.

Buying advice: Resolve food-use and preparation with an experienced local food-plant specialist before buying it as a household crop.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, momordica-balsamica

Waxberry

1 existing; 1 proposed. Products: berry fat - traditional processed product.

Food-use review: confirm identity, edible part and appropriate preparation with a qualified local adviser before including this plant as household food. Held out of ordinary food totals and picking jobs.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Historical processed fat, not a fresh-fruit forecast: Historical food use concerns the fatty berry coating, not a generic fresh sweet berry. No validated household extraction method, recovered food yield or storage life is established. Source coating/oil percentages are not assigned as processing conversion factors.

Treat edible-fat preparation as a specialist process requiring local validation.

Use well-drained coastal sand with sun or semi-shade; fruit production needs correctly sexed planting material.

Buying advice: Confirm the exact species and male/female stock with an indigenous nursery. Keep landscape use separate from an unverified household food process.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica: Morella cordifolia

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Black mulberry

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Exact fruit identity and variability: The institutional species reference distinguishes fruiting forms and variable breeding systems. Ripe fruits are used as food; a black-fruited tree must not silently inherit this identity or the legal status of another Morus species.

South African taxonomic cultivation record: The museum lists M. nigra as cultivated in southern Africa and distinguishes naturalised Morus taxa. It gives no ripe-harvest months or quantified yield.

Fruit maturity and water context: Institutional management reference links dry roots to early fruit drop. It contains a non-local 15-year substantial-bearing statement; that is not a South African first-crop age and is not used as a forecast.

Confirm the exact mulberry species and that the chosen plant is a fruiting form. Fruit colour alone cannot identify it.

Watch water stress during fruit development and record the actual ripe picking months.

Buying advice: Ask for identified Morus nigra fruiting material with a locally observed season. Do not use a generic black-fruited mulberry or fruitless ornamental as proof of this product.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: Orwa et al. (2009), World Agroforestry, Morus nigra profile, p. 1

Source: Iziko Museums, Biodiversity Explorer, Morus

Source: Orwa et al. (2009), World Agroforestry, Morus nigra profile, p. 4

Wild olive

1 existing; 1 proposed. Products: ripe wild-olive fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Wild olive is distinct from a named commercial olive cultivar. March-July describes fruit following flowers; no exact ripe-picking start/end is established.

SANBI records this slow-growing tree in summer and winter rainfall areas, with frost, drought and wind resistance.

Keep its roots away from walls, patios and pools; SANBI notes that they can be aggressive.

Buying advice: Ask for the wild-olive taxon in the catalogue, rather than substituting a commercial olive cultivar. Record actual local picking.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, olea-europaea-subsp-africana

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Olive

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Cultivar and product choice: Frantoio is described as an oil and cross-pollen candidate. Mission suits black table olives and oil; Kalamata black table, Manzanilla and Barouni green table. Climate fit and pollen needs depend on cultivar.

Regional orchard budget is not per-tree yield evidence: This is a historical one-hectare financial model; it does not provide matched per-tree kg or a measured biological maturity curve. No financial values or pesticide recipes are copied.

Match the cultivar to table olives or oil. Check drainage, irrigation and winter conditions; humid summer rainfall can bring disease problems.

Pick at the intended green, black or oil stage. Processing and safe kept-stock conditions must be recorded separately.

Buying advice: Mission is a source-backed black-table and oil candidate; Kalamata is for black table olives; Manzanilla and Barouni for green table olives; Frantoio for oil and compatible pollen. Confirm regional fit through SA Olive's registered nurseries.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Western Cape - cultivar and end-use dependent olive harvest: Feb-Jul.

Source: SA Olive Association, Olive Growing

Source: Western Cape Department of Agriculture, Oil Olives, Garden/Little Karoo budget, 13 March 2021, p. 1

Bietou

1 existing; 1 proposed. Products: ripe fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: Food is the ripe fruit. Main flowering in autumn/winter and fruit shortly afterwards do not establish named picking months.

SANBI describes a spreading pioneer shrub for sun and well-drained soil, including coastal and inland habitats. Leave room for its spread.

Buying advice: Confirm the exact species and local provenance; the checked source gives no named fruit cultivar.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, osteospermum-moniliferum

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Date palm

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Kakamas phenology is not a picking calendar: Table is fruit-set-to-maturation duration, not months of ripe harvest. Its separate table lists July-August flowering. Neither becomes a food-availability bar.

Southern Hemisphere reference, awaiting local confirmation: Broad hemispheric context, not a measured cultivar/site calendar. Kept outside the structured local harvest windows.

Western Cape Medjool ripening and spoilage trial: 2022 trial at Kleinplasie near Hermon concerns Medjool. Abstract treatment dates label ripening stages inconsistently; no harvest months are inferred. Full university handle returned HTTP 403 during checking.

Female palms need suitable male pollen. Record sex, cultivar and pollination management before forecasting fruit.

Check heat, water and rain during ripening. The Hermon trial found uneven ripening and spoilage, so regional success alone does not establish a local crop.

Buying advice: Medjool is documented in the Hermon study. Obtain known fruiting female material and compatible pollen advice; a seedling's sex and eventual fruit quality are not established by its appearance.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: FAO Date Palm Cultivation (2002), Chapter VIII

Source: FAO Date Palm Cultivation (2002), Chapter IV, Table 23

Source: FAO Date Palm Cultivation (2002), Chapter IX

Source: Peddie (2023), Stellenbosch MScAgric thesis, university-provided abstract in FAO AGRIS

Pistachio

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Producer's own cultivars: The same questionnaire confirms the nuts are grown on the producer's farm. Cultivar names are reproduced as written; spelling is not a taxonomy or catalogue change.

Prince Albert establishment and maturity expectation: An expectation for the newly established Platberg project, not a measured universal age curve. Its planted cultivars are Shufra, Ariyeh and Sirora; irrigated Karoo context retained.

Female trees bear nuts and need pollen from suitable male trees. The Prieska reference comes from a dry, irrigated Karoo farm.

Confirm male and female planting material, irrigation and local winter/summer conditions; a February producer reference does not validate a humid coast.

Buying advice: Discuss Shufrah and Ariyeh from the Prieska producer evidence, or Shufra, Ariyeh and Sirora from the Prince Albert project, with a specialist nursery. Select a compatible male and confirm local fit.

Pollination: separate male female plants. Check the chosen cultivar's requirements with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Orange River farm near Prieska, Northern Cape - Karoo Pistachios producer observation: Feb.

Source: Karoo Pistachios, first-person producer questionnaire on FoodHub

Source: Prince Albert Municipality, Pistachio Pioneers, 20 October 2025

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Spekboom

1 existing; 1 proposed. Products: edible leaves.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Edible part and local use: The food is the tart leaf; flowers do not define leaf picking months.

Primary food-processing research: The study analyses processed spekboom leaves. Its laboratory method and composition are not a farm harvest rate, storage life or household food-safety recipe.

Protect young plants from heavy browsing. SANBI reports warm rocky and semi-arid habitats; ground-level goat browsing can prevent vegetative spread and reduce regeneration.

Buying advice: Ask for correctly identified spekboom and record whether it is planted for food, browse or restoration. No measured household leaf yield is supplied.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: SANBI PlantZAfrica, portulacaria-afra

Source: Mahlanza et al., University of Johannesburg research record, International Journal of Food Science and Technology (2026)

Apricot

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Solitaire winter and pollen context: Solitaire is listed as self-compatible. Its chilling units are named-cultivar requirements and are not a numeric range for every apricot.

Irrigated Robertson trial context: The field experiment used managed organic orchards and found trade-offs in mulch effects; it does not justify a universal yield improvement from mulch.

Keep each cultivar's picking reference separate. The Robertson example used summer irrigation.

Confirm winter chilling, water and the actual planting material before using a reference as a local plan.

Buying advice: Solitaire has a sourced November first-pick reference and named chilling context at Simondium; discuss its local fit with the nursery.

Buying advice: Imperial has a December reference at Robertson. This is a local trial candidate, not advice for frost-free coastal KZN.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Bien Donne, Simondium, Western Cape - Solitaire first pick: Nov.

Reference only - Tierhoek Organic Farm, Robertson, Breede River Valley, Western Cape - Imperial trial: Dec.

Source: Culdevco / ARC Infruitec-Nietvoorbij, Solitaire apricot cultivar sheet

Source: Birkhofer et al. (2019), Effects of Ground Cover Management in Organic Deciduous Fruit Orchards in South Africa, Table 1

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Sweet cherry

1 existing; 1 proposed. Products: Ripe sweet cherries.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Production at age six: The guide reports 4-6 kg/tree in year 6 and warns that yields are difficult to predict because of pollination and fruit set. The separate good result at year 8-9 is 9 t/ha. Do not connect them into a made-up growth curve or convert the later hectare result to arbitrary tree spacing.

Chilling and pollen partners: Conventional commercial cherry reference: more than 1,000 Positive Utah Chill Units, not 1,000 chill hours. Low-chill candidates need their own verified requirements. Most cultivars need compatible cross-pollination; the named self-fertile choices differ. The September flowering reference is not a picking-month array.

Observed Ceres season start: Ceres growers reported the start of the 2025 cherry harvest in late October, seven to ten days earlier than the prior year. This is an observed season in a named region, not a fixed future duration, KZN calendar or all-variety season.

Check measured winter chilling, spring frost, drainage and rain near picking. The older KZN guide regarded commercial suitability there as still under trial; a broad climate-zone label does not replace current local advice.

Pollination depends on the selected cultivar. Compatible flowering and pollen sources are separate from early or late fruit ripening.

Buying advice: Regional guide candidates include Bing, Van, Rainier, Burlat and Vista, which generally need compatible pollen partners. It explicitly lists Lapins, Stella and Sweetheart as self-fertile. Minnie Royal, Royal Lee, Royal Dawn and Early Sweet are lower-chill candidates. Confirm current stock, exact rootstock, compatibility and measured local chilling before purchase; this is not a warm-coast recommendation.

Expected picking season: no fixed regional months sourced. Record planting and picking dates locally.

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Yield, p. 31

Source: KZN DARD, Fruit and Nut Production in KZN (2006), Sweet Cherry: Climate and Cultivars, p. 30

Source: Hortgro / Tru-Cape growers, Cherry season begins: Ideal conditions, 29 October 2025

Almond

1 existing; 1 proposed. Products: nuts.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Texas Mission pollen, chill and weight context: Texas Mission is classified high chill and names Nonpareil as pollinator. Its 5 kg/tree production entry does not state tree age, management, trial region or whether weight is kernel or in-shell; it is not a mature annual food forecast.

Texas Mission named production entry: The Texas Mission table's Production (kg/tree) row gives 5 kg/tree. It does not state tree age, reporting period, trial region or whether this means kernels or nuts in shell. Retained as a cultivar source claim to investigate; numeric mature annual kg remains null.

Check the named cultivar's winter needs and compatible pollen before planting. Texas Mission is self-incompatible.

Record whether nut weights mean kernels or nuts in shell; these are different quantities.

Buying advice: Texas Mission is a documented SAPO cultivar; its sheet names Nonpareil as pollen partner and describes a high chilling requirement. Confirm local suitability with the nursery.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - South African SAPO cultivar sheet; trial location not stated - Texas Mission first pick: Feb.

Source: SAPO Trust, Texas Mission almond cultivar information

SYNTHETIC GRAPH SOURCE {COUNT} $&

Farm choices - continued

Use your farm observations and the research shortlists. Confirm choices locally before buying.

Fruit, nuts and indigenous foods (continued)

Pear

1 existing; 1 proposed. Products: fruit.

Time to first crop: not sourced; do not rely on newly planted stock for immediate food.

Rosemarie Select pollen partners: These are cultivar-specific compatible-pollen candidates, not a guarantee that any two pear trees pollinate one another.

Packham's Triumph pollen partners: SAPO describes chilling qualitatively as low and production as high; neither supplies a numeric age or kg/tree.

Record the named cultivar and actual first and last picking months. January and February references are alternatives, not a mixed-orchard calendar for every tree.

Buying advice: Rosemarie Select is a source-backed Western Cape candidate to discuss with the nursery. Ask about December or Packham's Triumph as compatible pollen partners.

Buying advice: For Packham's Triumph, ask about Clapp's Favourite, December or Starkrimson and check local winter conditions.

Expected picking season: no single local cultivar window established. Reference seasons below are not a forecast.

Reference only - Ceres, Koue Bokkeveld, Western Cape - Rosemarie Select first pick: Jan.

Reference only - Wolseley, Western Cape - Rosemarie Select first pick: Jan.

Reference only - South African cultivar sheet; test location not stated - Packham's Triumph first pick: Feb.

Source: Culdevco / ARC Infruitec-Nietvoorbij, Rosemarie Select pear cultivar sheet

Source: SAPO Trust, Packham's Triumph pear cultivar sheet

Crop varieties to check locally

Keep the named variety you actually choose. Research shortlists need confirmation for your local season, water and ground.

Tomatoes

Recorded variety: not chosen yet.

This site's climate has not resolved, so no local shortlist is claimed. Open the crop card to see sources and record your choice.