SYNTHETIC GRAPH SOURCE {COUNT} $&

Pele o reka kapa o qala

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Sehlopha sa tlelaemete sa dipatlisiso: Sebaka sa temo ha se eso hlake. Sehlopha sena ha se keletso e netefaditsweng bakeng sa polasi ena.

Seo o se tlalehileng ka polasi ena

Metsi nakong ya komello

Ha ho a rekotwa. Qetella phuputso ya sebaka mme o netefatse metsi pele o kgetha moralo o hlokang disebediswa tse ngata.

Serame polasing ena

Se bonweng ka serame sebakeng ha se a rekotwa. Dikarolelano tsa kgwedi tsa sathelaete di ke ke tsa tiisa matsatsi a pele le a ho qetela a serame.

Letsatsi le phallo ya metsi mobung

Letsatsi: ha ho a rekotwa.

Phallo ya metsi: ha ho a rekotwa. Diphetho tsa teko ya mobu le ditlhoko tsa sejalo ka seng di ntse di hloka ho hlahlojwa.

Direkoto tsa selemo sa tlaleho - eseng ponelopele

Dikakaretso le dikgwedi tsa selemo sa tlaleho se fetileng di hlalosa se tlalehilweng. Ha di tshepise dijo moralong o tlang.

Original eggs {year} $&

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 240 eggs.

Tshebediso ya lelapa: 0 eggs. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: Php, Phu. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Fish

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: ha ho a rekotwa.

Tshebediso ya lelapa: ha ho a rekotwa. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: ha ho a rekotwa. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Meroho e meng

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 12 kg.

Tshebediso ya lelapa: 7 kg. Ho rekisitsweng: 3 kg.

Dikgwedi tsa kotulo tse tlalehilweng: Phe, Tsh. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae

Dogwood

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: tatso ya makgasi ha ho ritela — tshebediso ya setsebi.

Tlhahlobo ya tshebediso e le dijo: netefatsa hore ke semela sefe, karolo e jewang le tokiso e nepahetseng le moeletsi wa sebaka ya nang le tsebo pele o se kenya dijong tsa lelapa. Se tlohetswe dikakaretsong tsa dijo tse tlwaelehileng le mesebetsing ya kotulo.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Documented food-processing use is leaf bittering: The Ethiopian gesho-leaf brewing context establishes a specialist processed flavour use, not ordinary fresh fruit nutrition or a South African household recipe. No leaf kg/plant, food storage life or local harvest date is assigned.

South African berry phenology concerns wildlife food: December–June berries are documented in the ecology account as wildlife food; this is not a verified human berry-picking window.

Netefatsa hore ke semela sefe hantle le tokiso ya dijo e rerilweng le moeletsi wa lehae ya nang le boiphihlelo.

Tlaleho ya South Africa e hlalosa kgolo mabopong a melatswana le matsoapong a meru le ho mamella serame sa leqhwa; monokotshwai wa sona o hlaloswa e le dijo tsa dinonyana.

Keletso ya ho reka: Reka dimela tsa mofuta o nepahetseng bakeng sa tshebediso e rerilweng e netefaditsweng. O se ke wa nka bopaki ba tshebediso ya moriana kapa ba ditholwana tsa diphoofolo tse hlaha e le ponelopele ya dijo tse foreshe tsa lelapa.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: Negash et al. (2021), primary study of Rhamnus prinoides leaves

Mohlodi: SANBI PlantZAfrica: Rhamnus prinoides

Baboon grape

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: menokotshwai e butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: The source gives a ripe-berry interval. This does not establish stored jam months or safe shelf life.

Fana ka tshehetso ya ho nama mme o laole ho ata ha sona moo sebaka se haellang. SANBI e se rekota ditutulung tsa lehlabathe tsa lebopong le mathokong a meru ho tloha Western Cape ho ya KwaZulu-Natal le Mpumalanga.

Keletso ya ho reka: Netefatsa Rhoicissus digitata; mefuta e meng ya wild-grape e na le ditlhaloso tse fapaneng mme e ke ke ya fana ka matsatsi a yona.

Nako ya kotulo e lebeletsweng: ha ho nako e le nngwe e tiisitsweng ya mofuta wa sebaka. Dinako tsa ditshupiso tse ka tlase ha se ponelopele.

Tshupiso feela — tlaleho ya botaniki ya Western/Eastern Cape, KwaZulu-Natal le Mpumalanga — monokotshwai o butswitseng: Hlb–Pha.

Mohlodi: SANBI PlantZAfrica, rhoicissus-digitata

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Bushman's grape

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: monokotshwai o jewang; metso ha e kenyeletswe.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Food is the berry; the same SANBI account reports poisonous tuberous roots. September–June describes berry occurrence, not a measured ripe-picking interval.

Fana ka foreime ya ho nama kapa tshehetso e nngwe. Dimela tse ntjha di hloka metsi ho fihlela di tsitsitse; SANBI e hlalosa ho mamella komello hamorao le tshebediso letsatsing ho ya moriting o sa fellang.

Keletso ya ho reka: Kopa Rhoicissus tridentata mme o rekote dikgwedi tsa ditholwana tse butswitseng sebakeng seo. O se ke wa sebedisa tshebediso ya metso e le moriana kapa bakeng sa diphoofolo e le bopaki ba dijo.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica, rhoicissus-tridentata

Rosemary

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi a foreshe kapa a omisitsweng bakeng sa ho noka dijo.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Food-leaf product distinct from essential oil: The brochure uses the older botanical name Rosmarinus officinalis; catalogue identity is preserved. It documents food seasoning leaves and separate essential oil uses, without a food-leaf yield, first cut age or local picking calendar.

Sebedisa mobu o ntshang metsi hantle le metsi a lekaneng nakong ya ho tiya ha dimela; qoba ho tlala metsi le ho bola ha metso.

Rekota makgasi a foreshe, senoko se omisitsweng le oli ya monko wa semela (essential oil) e le dihlahiswa tse fapaneng.

Keletso ya ho reka: Fumana dimela tsa rosemary tse phetseng hantle tse hlwauweng le keletso ya lehae ya ho di hodisa; ha ho mofuta wa sebaka wa tlhahiso ya makgasi a ho pheha o netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: DAFF: Rosemary production brochure (2012), p. 2

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Karoo boer-bean

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dipeo tse jewang.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Edibility is recorded for this exact variety; do not apply the common name boer-bean to every species as food evidence.

Tikoloho ya sona ya tlhaho e kenyelletsa mabopo a ommeng a melatswana Little Karoo le dibakeng tse ommeng ho feta tsa Eastern Cape le borwa ba Western Cape.

Li-pod (mekotlana ya peo) di hasanya dipeo ha di omme. Rekota ho bokella le ho sebedisa peo sebakeng sa heno; sena ha se kotulo ya dijo e lekantsweng.

Keletso ya ho reka: Kopa Schotia afra var. afra, ho netefaditswe hore ke semela sefe hantle ka thuto ya dimela. Tlaleho ya mohlodi ya moriti ka dilemo tse hlano ha se dilemo tsa dijo tsa pele.

Nako ya kotulo e lebeletsweng: ha ho nako e le nngwe e tiisitsweng ya mofuta wa sebaka. Dinako tsa ditshupiso tse ka tlase ha se ponelopele.

Tshupiso feela — tlaleho ya botaniki ya Little Karoo le Cape e ommeng ho feta — tshupiso ya tlhahiso ya peo e jewang: Mot–Php.

Mohlodi: SANBI PlantZAfrica, schotia-afra-var-afra

Karee

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: menokotshwai e butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: September–January describes fruit production; exact ripe-picking dates are not stated. Do not count leaves used as livestock fodder as household food.

SANBI e rekota sefate sena sa moriti se mamellang serame sa leqhwa le komello maemong a Karoo le Kalahari.

Ho ya ka SANBI, makgasi a sona a mangata a ka senya tatso ya lebese la diphoofolo tsa lebese; sena ha se kgothaletso ya kabo ya dijo.

Keletso ya ho reka: Botsa ka dimela tse tona le tse tshehadi haeba ho rerilwe ditholwana. Kabo ya tlhaho mohloding ha e kenyelletse KwaZulu-Natal; ho mamella maemo a mangata ha se tiisetso ya ditholwana sebakeng seo.

Ho tsamaiswa ha peo ya dipalesa: dimela tse arohaneng tse tona le tse tshehadi. Hlahloba ditlhoko tsa mofuta o kgethilweng le sebaka sa kgodiso ya dimela.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica, searsia-lancea

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Glossy currant

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: ditholwana tse butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Currant fruit food use and phenology: The primary account establishes edible ripe fruit and October–June fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Kga ditholwana tse butswitseng mme o rekote ho omisa ka thoko.

Sehlahla sena se tshwaneleha dirapeng tsa lebopong tse pepesehileng le ho tsitsisa ditutulu tsa lehlabathe; netefatsa tshimoloho ya lehae.

Keletso ya ho reka: Kopa setsi sa dimela tsa tlhaho bakeng sa mofuta o nepahetseng le dimela tsa lehae. Ha ho mofuta wa dijo wa sebaka o fanang ka kotulo e ngata o netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Searsia lucida

Natal currant

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: ditholwana tse butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Edible Natal currant fruit and local phenology: The primary account establishes edible ripe fruit and February–July fruit appearance. That interval includes developing fruit and is retained only as phenology, not a ripe picking calendar. No annual food kg or bearing age is supplied.

Dumella ditholwana ho butswa. Sehlahla sena se mamellang komello se ka etsa lekhoakhoa; se pome ho lekana sebaka.

Keletso ya ho reka: Netefatsa mofuta o nepahetseng le setsi sa lehae sa dimela tsa tlhaho; ha ho mofuta wa sebaka o lekilweng kapa kgetho ya kotulo e lekantsweng ka palo e netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Searsia natalensis

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Kuni bush

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: ditholwana tse butswitseng tse omisitsweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Named traditional dried-fruit food use: The discussion explicitly includes S. undulata among edible dried ripe currant fruits and traditional fermentation uses. Its host article species calendar is not transferred to S. undulata.

Exact species and habitat: Assessment identifies the catalogue species and Northern/Western Cape distribution; it does not provide a harvest calendar or food yield.

Rekota ho kga ditholwana tse butswitseng le ho omisa ka thoko. Ha ho nako ya polokelo kapa risepe ya ho bedisa e fanweng.

Keletso ya ho reka: Netefatsa mofuta o nepahetseng le tshimoloho ya lehae; tekolo ya SANBI e se beha matsoapong a majwe a fynbos/karroid a Northern Cape le Western Cape.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Searsia burchellii, food-use discussion naming S. undulata

Mohlodi: SANBI Red List: Searsia undulata

White milkwood

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: menokotshwai e butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: SANBI qualifies the food statement; it is about ripe berries, not latex, bark or roots. Fruit presence February–September does not establish ripe picking throughout.

Species identification in KwaZulu-Natal: The primary study records confusion with Bridelia micrantha. It does not supply a safe harvest calendar or kg/tree.

Sireletsa sefate sena sa lebopong moyeng o tobileng wa lewatle o tletseng letswai; SANBI e tlaleha ho tjheswa ke letswai leha tikoloho ya sona e le ya lebopong.

Keletso ya ho reka: Kopa setsi sa dimela ho netefatsa mofuta le tshimoloho e molaong. Hlahloba ditlhoko tsa hajwale tsa ho bokella pele o nka dimela kapa dihlahiswa difateng tse teng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica, sideroxylon-inerme

Mohlodi: University of KwaZulu-Natal, Phytochemical and elemental studies of two indigenous medicinal plants

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Dune spinach

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dintlha tse ntjha tse nang le makgasi le dikutu.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Leaf vegetable harvesting and preparation: New rainy-season growth gives the best tops; wash off coastal sand thoroughly and boil for the documented vegetable preparation. August–November flowering is not leaf availability.

Kotula dintlha tse ntjha tse nang le makgasi; hlatswa lehlabathe pele o pheha.

Sebedisa lehlabathe la lebopong le ntshang metsi hantle mme o boloke dimela tse arotsweng di le mongobo ho fihlela di tsitsitse.

Keletso ya ho reka: Fumana dimela tsa ho lema tsa lebopong tse hlwauweng hantle; rekota kgolo le dikgwedi tsa ho kga makgasi tsa lehae hobane dinako tsa pula di fapana ho pholletsa le South Africa.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Tetragonia decumbens

Kinkelbossie

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: mahlomela a matjha, a phehilweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Young-shoot vegetable use: The account describes boiling young shoots until tender. September–January flowering is not a shoot harvest calendar; fodder output is not household food kg.

Kotula mahlomela a matjha bakeng sa ho pheha. Rekota kotulo ya dijo ka thoko ho furu ya diphoofolo.

Sebedisa sebaka se nang le letsatsi se ntshang metsi hantle sa lebopong, fynbos kapa Karoo se loketseng semela sena sa pele se sebedisang metsi ka hloko.

Keletso ya ho reka: Fumana dimela tsa ho lema tsa lehae tse hlwauweng hantle; ha ho mofuta wa dijo o nang le lebitso kapa tekanyetso ya selemo ya kg/plant e nang le mohlodi.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Tetragonia fruticosa

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Veldkool

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dithunthung tsa dipalesa.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Edible bud identity: Coastal-sand perennial with edible congested flower buds. The stated July–September flowering interval does not identify exact unopened-bud picking months.

Rekota ho kga dithunthung tse ntjha tsa dipalesa ka thoko ho dipalesa tse bulehileng.

Bapisa semela le maemo a lehlabathe la lebopong le ntshang metsi hantle le kgolo ya sona ya dinako tsa selemo.

Keletso ya ho reka: Sebedisa tlhwayiso ya mofuta o nepahetseng setsing sa dimela; ha ho dilemo tsa kotulo ya pele ya dithunthung kapa tekanyetso ya dijo ya kg/plant e netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Trachyandra ciliata

Pigeonwood

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi a matjha bakeng sa moroho o phehilweng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Documented young-leaf food use: The human food product is young leaves. Berries described as bird/bat food are not promoted to a human fruit harvest. No leaf cutting age, exact months or household leaf kg is given.

Bapisa semela sena sa pele se holang ka potlako le maemo a futhumetseng a mongobo a botjhabela/leboya; sireletsa dimela tse nyane tse leng mongobong le moriting.

Rekota kotulo ya makgasi a matjha ka thoko ho ditholwana tse jewang ke diphoofolo tse hlaha.

Keletso ya ho reka: Sebedisa dimela tsa ho lema tsa lehae tse hlwauweng hantle; ha ho mofuta o nang le lebitso wa makgasi a dijo kapa tlhahiso ya dijo ya selemo e lekantsweng ka palo e netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica: Trema orientalis

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Wild garlic

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: makgasi le dipalesa tse jewang.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: Leaves and flowers are documented culinary parts. Medicinal bulb use does not make every part a food.

Se mamella komello e telele empa se hola hantle ho feta ka metsi a kamehla. SANBI e rekota kabo ya sona ka botjhabela ho South Africa.

Keletso ya ho reka: Kopa Tulbaghia violacea ho ena le semela se hlwauweng feela e le wild garlic. Rekota ho kga makgasi le dipalesa ha lehae ka thoko.

Nako ya kotulo e lebeletsweng: ha ho nako e le nngwe e tiisitsweng ya mofuta wa sebaka. Dinako tsa ditshupiso tse ka tlase ha se ponelopele.

Tshupiso feela — tlaleho ya botaniki ya botjhabela ba South Africa — dipalesa tse jewang feela: Phe–Mme.

Mohlodi: SANBI PlantZAfrica, tulbaghia-violacea

Ginger

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: dikutu tse foreshe tse melang ka tlasa mobu; ginger e omisitsweng e rekotwa ka thoko.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

South African trial harvest duration and native yields: Pretoria Hillcrest rain-shelter trial, 2015/16 and 2016/17, drip irrigation on sandy clay loam. Commercial ginger Z. officinale at eight months: 38.3 t/ha fresh and 8.8 t/ha dry rhizomes in the well-watered treatment; African ginger was a different species with different values. No matched plant density is supplied, so no kg/plant or household drying conversion is derived. The abstract says open field, while Methods specifies rain shelter; conditions are retained.

Confirmed culinary species and product: Extension identifies culinary ginger rhizomes used as spice, preserves and beverages. Florida planting and harvest months are not imported into the South African calendar.

Harvest recommendation: The author abstract recommends harvest seven to eight months after planting; it is a rhizome crop-cycle duration, not a full-bearing tree age.

Sebedisa letsatsi la ho lema ho rekota potoloho ya kotulo ya rhizome, e leng kutu e melang ka tlasa mobu. Tshupiso ya teko ya Pretoria e na le dipehelo, ha se khalendara e sa fetoheng ditlimaeteng tsohle.

Rekota boima ba rhizome e foreshe le senoko se omisitsweng ka thoko; boloka ginger e jewang e fapane le mefuta ya ginger ya mokgabiso kapa ya African ginger.

Keletso ya ho reka: Kopa mofani wa lehae di-rhizome tsa ho lema tsa Zingiber officinale tse phetseng hantle tse hlwauweng; ha ho kerafo ya kotulo ya mofuta o itseng wa South Africa e netefaditsweng.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: Gatabazi et al. (2022), Scientia Horticulturae 304:111353, p. 17

Mohlodi: UF/IFAS: Ginger, Galangal, and Turmeric Production in Florida

Mohlodi: University of Pretoria author manuscript: Gatabazi et al. (2022)

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Ditholwana, linate le dijo tsa lehae (e tswela pele)

Buffalo thorn

Tse teng: 1; tse rerilweng: 1. Dihlahiswa: menokotshwai e butswitseng.

Nako ya kotulo ya pele: ha e a fumanwa mohloding; o se ke wa itshetleha ka dimela tse sa tswa jalwa bakeng sa dijo hanghang.

Karolo e jewang le tshebediso ya sebaka: SANBI describes dry, slightly sweet ripe pulp. Fruit remaining on leafless trees March–August is persistence, not a verified six-month picking season.

SANBI e rekota kabo e pharaletseng ho tloha lebopong le se nang serame sa leqhwa ho ya dibakeng tsa serame sa leqhwa sa mariha. Meutlwa ya sona e kobehileng le e otlolohileng e bohlokwa ha ho kgethwa ditsela tsa phihlello.

Keletso ya ho reka: Kopa semela sa buffalo-thorn se hlwauweng se loketseng maemo a lehae; mohlodi o hlahlobilweng ha o fane ka mofuta wa dijo o nang le lebitso.

Nako ya kotulo e lebeletsweng: ha ho mohlodi wa dikgwedi tse tsitsitseng tsa sebaka. Rekota matsatsi a ho jala le kotulo sebakeng.

Mohlodi: SANBI PlantZAfrica, ziziphus-mucronata

Mefuta ya dijalo e lokelang ho hlahlojwa sebakeng seo

Boloka lebitso la mofuta oo o o kgethang ka nnete. Manane a makgutshwane a dipatlisiso a hloka netefatso bakeng sa nako ya sebaka, metsi le mobu.

Tomatoes

Mofuta o rekotilweng: ha ho eso kgethwe.

Maemo a lehodimo a sebaka sena ha a eso hlake, kahoo ha ho bolelwe lenane le lekgutshwane la sebaka. Bula karete ya sejalo ho bona mehlodi le ho rekota kgetho ya hao.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.