SYNTHETIC GRAPH SOURCE {COUNT} $&

Pele o reka kapa o qala

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Sehlopha sa tlelaemete sa dipatlisiso: Sebaka sa temo ha se eso hlake. Sehlopha sena ha se keletso e netefaditsweng bakeng sa polasi ena.

Seo o se tlalehileng ka polasi ena

Metsi nakong ya komello

Ha ho a rekotwa. Qetella phuputso ya sebaka mme o netefatse metsi pele o kgetha moralo o hlokang disebediswa tse ngata.

Serame polasing ena

Se bonweng ka serame sebakeng ha se a rekotwa. Dikarolelano tsa kgwedi tsa sathelaete di ke ke tsa tiisa matsatsi a pele le a ho qetela a serame.

Letsatsi le phallo ya metsi mobung

Letsatsi: ha ho a rekotwa.

Phallo ya metsi: ha ho a rekotwa. Diphetho tsa teko ya mobu le ditlhoko tsa sejalo ka seng di ntse di hloka ho hlahlojwa.

Direkoto tsa selemo sa tlaleho - eseng ponelopele

Dikakaretso le dikgwedi tsa selemo sa tlaleho se fetileng di hlalosa se tlalehilweng. Ha di tshepise dijo moralong o tlang.

Original eggs {year} $&

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 240 eggs.

Tshebediso ya lelapa: 0 eggs. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: Php, Phu. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Fish

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: ha ho a rekotwa.

Tshebediso ya lelapa: ha ho a rekotwa. Ho rekisitsweng: ha ho a rekotwa.

Dikgwedi tsa kotulo tse tlalehilweng: ha ho a rekotwa. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Meroho e meng

Kakaretso ya selemo e tlalehilweng ke molemi bakeng sa 2025: 12 kg.

Tshebediso ya lelapa: 7 kg. Ho rekisitsweng: 3 kg.

Dikgwedi tsa kotulo tse tlalehilweng: Phe, Tsh. Tsena ke tse bonweng phuputsong, eseng ponelopele e ntjha ya tlhahiso.

Hlahloba yuniti le dikakaretso tse rekotilweng pele o sebedisa mola ona.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo

Dinotshi: Honeybee (managed hives)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: mahe a dinotshi. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Kotulo e itshetlehile ka sehlopha sa dinotshi, dipalesa le pula ya sebaka. Hlahloba ho butswa mme o siele dinotshi dijo; ha ho nako e tsitsitseng ya naha ya mahe a dinotshi.

Nako ya sehlahiswa sa pele: ha ho mohlodi bakeng sa kgwebo ena.

Tshupiso ya tlhahiso: 25 kg ya mahe a dinotshi/ka ntlo ya dinotshi (hive)/ka selemo. From a 2008 national postal survey of 112 SABIO-member beekeepers (22.4% response rate) managing an estimated 19,520 hives at 2,094 apiary sites; figure is self-reported and un-weighted. Table 7 (same doc, p.13-14) shows wide variance by region/operation-focus/size for the same year: e.g. 'All' region, honey-focused, all-size operations = 25.1-26.9 kg/hive/yr across 3 survey sites; Western Cape honey-focused = 23.6-27.9 kg/hive/yr; Western Cape pollination-focused, small operations = only 7.0-7.5 kg/hive/yr. The same document also quotes an earlier, more rigorously validated study: 'Their average reported yield is very low, at just 13.4 kilograms per hive' for Western Cape hives (Allsopp & Cherry 2004, cited p.2 of this working paper), and historical national bee-census data of 8.3 kg/hive/annum (1911-1988 average) rising to 24 kg/hive in the 1974/5 census (same doc, p.1). No single number can be called definitive; 25 kg/hive/yr is used as the document's own headline national estimate.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Beha matlo a dinotshi hole le mabala a ho bapala, fana ka metsi a hlwekileng mme o sebedise sebaka se ommeng. Kopa morui wa dinotshi ya ngodisitsweng ho hlahloba ho bewa ho bolokehileng sebakeng sa heno pele o eketsa ntlo ya dinotshi.

Hlahloba sehlopha ka seng sa dinotshi kgafetsa bakeng sa matshwao a lefu la dinotshi le tlamehang ho tlalehwa le tshwaetso ya Cape-bee (Capensis).

Beha le ho sireletsa matlo a dinotshi kgahlanong le boshodu le tshenyo ya ka boomo, e leng e nngwe ya diphephetso tse kgolo ka ho fetisisa ho barui ba dinotshi ba Afrika Borwa.

Register with the Department before beekeeping. Registration lasts 24 months from registration or renewal and must be renewed when it expires.

A beekeeper who fails to renew registration at expiry will be deregistered and face penalties under the Act.

Beekeeping activities and hired beekeeping services require a valid registration certificate. Check the certificate before using a beekeeper.

Every beehive must be clearly and legibly marked with the beekeeper's registration number.

Keep written records of colonies, apiary locations, diseases, disease-control measures and colony losses with their reasons.

Do not move honeybees across the Cape bee line, or keep Cape bees north of it, without written approval.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dinotshi: Honeybee (managed hives) (e tswela pele)

Report any suspected notifiable disease (American foulbrood is listed); officials can order an infected colony destroyed.

Mohlodi: FAO: Beekeeping in Africa, Site selection (general African apiary guidance)

Mohlodi: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 6

Mohlodi: Gums & Bees: A roadmap for landowners in South Africa (South African National Biodiversity Institute, SANBI), p. 21

Mohlodi: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(1) and 2(4), p. 4

Mohlodi: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measure 2(5), p. 4

Mohlodi: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, control measures 2(8) and 2(9), p. 5

Mohlodi: Department of Agriculture: Control Measures relating to honey-bees R.858 of 2013, consolidated with R.1511 of 22 November 2019, p. 5

Mohlodi: Survey of Beekeeping in South Africa, CSSR Working Paper No. 221 (Conradie & Nortje, 2008, UCT Centre for Social Science Research), p. 14

Mohlodi: FAO, Beekeeping in Africa, honey harvesting

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Lesaka: Nguni / crossbred beef cattle (communal and extensive veld)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: nama. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Nako ya sehlahiswa sa pele: ha ho mohlodi bakeng sa kgwebo ena.

Tshupiso ya tlhahiso: 114.5–121.2 kg ya boima nakong ya ho kgweswa/ka namane. Table 5 gives weaning weight (WW) across four agro-ecological zones for the same nine Nguni communal herds: arid 114.51 kg (lowest), semi-arid 119.98 kg, dry sub-humid 117.91 kg, humid 121.21 kg (highest). The range recorded here spans this one study's own arid-to-humid zone means; it is not merged with any other paper's weaning-weight figures (other SA studies report different ranges for different systems - e.g. research-station or stud herds - and were deliberately excluded to avoid mixing systems). 826 Nguni calves from nine herds that the paper itself says were 'kept under extensive grazing system' in Limpopo Province - a genuine communal/extensive veld system matching this enterprise.

Tshupiso ya nako ya tlhahiso: dilemo tse 6.4. 2354 days / 365.25 = 6.44 years. 'Failure time' is the length of productive herd life for the 580 uncensored (fully-observed) records in the combined dataset; the dissertation itself defines productive herd life as 'the time interval between first calving date to culling date'. Combined data from three extensively-grazed Nguni stud herds, animals born 1968-2015.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Masaka a dikgomo ha a tshwanela ho tlala haholo: ho tshwanetse ho ba le sebaka se lekaneng sa fatshe hore phoofolo e nngwe le e nngwe e robale ka nako e le nngwe.

Nakong ya dipalangwang, kgomo ka nngwe e tshwanetse ho ba le bonyane 1.4 m^2 ya sebaka sa fatshe hore ho laiswa ho se ke ha petetsana haholo kapa ha ba le sebaka se seholo haholo.

Dikgomo ha di a tshwanela ho tsamaiswa dihora tse fetang 18 di sa theolwe mme tsa fuwa dijo, metsi le bodulo bo di sireletsang.

Ke tlolo ya molao ho bolaisa dikgomo tlala, ho di fepa hanyane haholo, kapa ho di hanela metsi kapa dijo.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, as part of the national bovine brucellosis control scheme under the Animal Diseases Act.

No animal may be slaughtered anywhere but a registered abattoir, with the exception below.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Lesaka: Nguni / crossbred beef cattle (communal and extensive veld) (e tswela pele)

Slaughter for the farmer's own household consumption, or for cultural or religious purposes, is exempt from the abattoir rule.

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.4.6.1, p. 7

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Mohlodi: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Mohlodi: Animal Identification Act 6 of 2002, s.7(1), p. 4

Mohlodi: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Mohlodi: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: Mpofu, Ginindza, Siwendu, Nephawe & Mtileni (2016). Effect of agro-ecological zone, season of birth and sex on pre-weaning performance of Nguni calves in Limpopo Province, South Africa. Tropical Animal Health and Production 49:187-194 (PMC5203849), Table 5

Mohlodi: Ngayo, M. (2017). Factors Affecting Productive Life and Fertility in Nguni Cows (MSc dissertation, University of the Free State), Table 5.1, p. 58

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Lesaka: Small-scale dairy cow (Jersey, Holstein or crossbred)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: lebese. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Rera ho ya ka ho tswala ha nnete ha manamane/madinyane le nako ya lebese. Diphoofolo tse sa hlahiseng lebese ha di fane ka lona.

Tshupiso ya sehlahiswa sa pele: dibeke tse 121.8–130.5. Table 1, AFC column, rows in the table's order Holstein-Friesland Reg./Grade, Jersey Reg./Grade, Ayrshire Reg./Grade: 28, 30, 25, 28, 29, 32 months. The Grade rows used here are Holstein-Friesland 30 and Jersey 28 (Grade Ayrshire 32 is not used: the enterprise is Jersey or Holstein type). 28 x 4.35 = 121.8 weeks; 30 x 4.35 = 130.5 weeks. Milk begins at first calving, so age at first calving is the time to first product.

Tshupiso ya tlhahiso: 3840–5422 kg ya lebese/ka kgomo e tshehadi/ka nako ya ho hlahisa lebese. Table 1 ('Comparison of number of cows, production and fertility in the main dairy breeds in South Africa'), Milk column, Grade rows: Jersey 3840 kg and Holstein-Friesland 5422 kg per lactation (Registered rows: 4468 and 6426 kg). The document says these are National Milk Recording Scheme herds, which 'have been shown to be better managed than those outside the scheme, with their performance per lactation being 50 % greater than that of the national herd', so a smallholder herd should expect less.

Tshupiso ya nako ya tlhahiso: dilemo tse 1.9–2. Holstein 739.33 and Jersey 696.81 days of productive life (first calving to culling) in South African milk-recorded herds, 2000-2019: 696.81/365.25 = 1.91 and 739.33/365.25 = 2.02 years.

Tshupiso ya dilemo/mokgwa: In hotter, humid KZN, KZN DARD describes Nguni × Jersey dairy trials at Makhathini to combine milk potential with local adaptation. Hand milking with the calf present highlights competition between household milk and calf growth. Discuss adapted stock and feeding locally; commercial milk-recording totals do not forecast this system. Record calving, lactation stage and milk available after calf needs.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Masaka a dikgomo ha a tshwanela ho tlala haholo: ho tshwanetse ho ba le sebaka se lekaneng sa fatshe hore phoofolo e nngwe le e nngwe e robale ka nako e le nngwe.

Dikgomo tse tshehadi ha di a tshwanela ho laiswa bakeng sa dipalangwang haeba matswele a tsona a ruruhile ka lebese.

Dikgomo ha di a tshwanela ho tsamaiswa dihora tse fetang 18 di sa theolwe mme tsa fuwa dijo, metsi le bodulo bo di sireletsang.

Ke tlolo ya molao ho bolaisa kgomo ya lebese tlala, ho e fepa hanyane haholo, kapa ho e hanela metsi kapa dijo.

Dikgomo tse tshehadi makgulong kapa dikampong tse bulehileng di hloka moriti: meaho ya moriti e tshwanetse ho fana ka bonyane 4 ho isa ho 5 m^2 ya moriti ka kgomo.

Foot-and-mouth disease (FMD) is a controlled animal disease under the Animal Diseases Act: state Veterinary Services enforce isolation and movement restrictions.

Every cattle owner must register an official identification mark and brand or mark their cattle with it, under the Animal Identification Act.

Dairy farmers are legally required to test their herds for bovine TB, mainly because of export requirements, so most dairy cattle in South Africa are tested.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Lesaka: Small-scale dairy cow (Jersey, Holstein or crossbred) (e tswela pele)

Any suspected or confirmed brucellosis-related abortion in cattle must be reported to the State Veterinary office, under the national bovine brucellosis control scheme.

A cull cow may be slaughtered only at an abattoir, except for the farmer's own consumption or for cultural or religious purposes.

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.1.2, p. 4

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.6.13, p. 9

Mohlodi: Livestock Welfare Coordinating Committee, Code of Practice: Handling and Transport of Livestock (2018), s.7.1, p. 9

Mohlodi: Animals Protection Act 71 of 1962, s.2(1)(c), p. 1

Mohlodi: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 129

Mohlodi: SAnews: KZN Foot and Mouth Disease boundaries extended (Dept of Agriculture statement)

Mohlodi: Animal Identification Act 6 of 2002, s.7(1), p. 4

Mohlodi: Bovine tuberculosis / dairy herd testing study, PMC10369683

Mohlodi: Marumo, Hlokwe & Kayoka-Kabongo, Seroprevalence of brucellosis in communal and smallholder cattle farming in North West Province, South Africa, Onderstepoort Journal of Veterinary Research (PMC10784229)

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: KZN DARD: Subsistence Dairying in KZN, Makhathini Research Station trials

Mohlodi: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), p. 10

Mohlodi: Muller, C.J.C., The Dairy Farming Handbook (2017), Western Cape Department of Agriculture (Elsenburg), Table 21.1, p. 122

Mohlodi: Gertenbach, W.D., Dairying in KwaZulu-Natal: Breeds of Dairy Cattle, Cedara Agricultural Development Institute, KZN Department of Agriculture and Rural Development, Table 1, p. 1

Mohlodi: Cow Longevity and Reasons and Risk Factors for Culling in South African Holstein and Jersey Dairy Herds, Animals 2025 (PMC12560872)

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Broilers (commercial)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: nama. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: kgetha mofuta wa nama le moralo wa dihlopha wa mofani; netefatsa kgodiso ya ditsuonyana ka mofuthu, dijo, metsi le ditokisetso tsa ho hlaba.

Rera dijo tse leka-lekaneng, metsi a hlwekileng, moriti, moya le tshireletso ho dibatana. Phepelo ya mahe e itshetlehile ka dinonyana tse behelang, dilemo tsa tsona, bophelo le tlhokomelo; hoko feela ha e fane ka mahe.

Tshupiso ya sehlahiswa sa pele: dibeke tse 4.6–4.7. 32 days / 7 = 4.57 weeks; 33 days / 7 = 4.71 weeks. This enterprise has a single grow-out cycle ending in slaughter, so 'time to first product' = age at slaughter. South Africa, commercial industry average.

Tshupiso ya tlhahiso: 1.8–1.9 kg/ka nonyana nakong ya ho hlajwa (matsatsi a 32–33). National commercial broiler industry average, South Africa, 2022. A separate breed-trial table (Table 4b, same document, p.39) gives named-strain 'Cumulative FI (kg)' and 'Body weight (kg)' at exactly 35 days: Arbor Acres Plus 2.29kg, Cobb 500 2.52kg, Indian River 2.30kg, Ross 308 2.30kg (cumulative feed intake 3.21-3.64kg) - a slightly older age (35 vs 32-33 days) and higher weight than the industry-wide average quoted above; not merged into the primary figure since it is a different (breed-standard trial) dataset, but used below for the feed-intake figure.

Tshupiso ya dilemo/mokgwa: Newer South African commercial reference: SAPA 2024 reports average slaughter at 30–31 days with bird weight 1.8–1.85 kg. The primary range above remains the earlier 2022 reference at 32–33 days. These are commercial bird weights, not dressed meat kg or a village flock target. Record hatch or purchase age, feeding system, actual harvest date and weighed product basis.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Dikgoho tsa nama tse dulang fatshe ka tlung: boima ba dikgoho bo se ke ba feta 40 kg ka mithara ya sekwere.

Madinyane a dikgoho a sa tswa qhotswa a tshwanetse ho fuwa dijo nakong ya dihora tse 24 ka mora ho qhotswa.

Madinyane a dikgoho a sa tswa qhotswa a tshwanetse ho fumana metsi nakong ya dihora tse 24 ka mora ho qhotswa, kapele ho feta moo ha ho tjhesa.

Metsi a hlwekileng le dijo tse nepahetseng di dule di fumaneha, hore dikgoho di se ke tsa lapa kapa tsa nyorwa.

Matlo a dikgoho tsa nama a hloka tekanyo ya moya o kenang le o phallang ya bonyane dimithara tse 5 tsa khubiki ka hora bakeng sa khilograma ka nngwe ya boima ba dikgoho tse ka tlung.

Tlhahisong ya dikgoho tsa nama tse itsamaelang kantle, palo e bolokwang ka tlung ha e a tshwanela ho feta dikgoho tse 15 ka mithara ya sekwere ya fatshe.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Broilers (commercial) (e tswela pele)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), 2.2 Housing of broilers in Floor Systems, p. 6

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), p. 11

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), p. 12

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), p. 7

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), p. 8

Mohlodi: SAPA Code of Practice for Broiler Production (May 2012), 2.4 Free Range Broiler Production, p. 8

Mohlodi: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Mohlodi: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Mohlodi: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: SAPA Industry Profile 2024, section 7.9 Performance efficiency, p. 82

Mohlodi: SAPA Industry Profile 2022, section 6.9 Performance efficiency, p. 79

Mohlodi: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Village chickens (indigenous, free-range)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: mahe. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Buying advice: Potchefstroom Koekoek is a dual-purpose option; Venda, Ovambo and Naked Neck are other ARC-conserved options. No province-specific winning breed is claimed.

Rera dijo tse leka-lekaneng, metsi a hlwekileng, moriti, moya le tshireletso ho dibatana. Phepelo ya mahe e itshetlehile ka dinonyana tse behelang, dilemo tsa tsona, bophelo le tlhokomelo; hoko feela ha e fane ka mahe.

Bokella mahe kgafetsa, boloka moo dihlaha di alwang ho hlwekile ho omme, mme o tshware/boloke mahe ka hloko bakeng sa tshebediso ya lelapa kapa thekiso.

Tshupiso ya sehlahiswa sa pele: dibeke tse 15.7–27.8. Free-range village chickens, KwaZulu-Natal; dry-environment birds started laying earlier (15.74 weeks) than wet-environment birds (27.78 weeks), difference flagged significant ('*') in the source table. Wide range reflects genuine dry/wet-season environmental variation reported by the source, not measurement uncertainty.

Tshupiso ya tlhahiso: 40.4–49.9 mahe/ka kgoho e tshehadi/ka selemo. Free-range/scavenging village chickens, KwaZulu-Natal: dry (40.43) vs wet (49.91) environment, difference not significant ('NS'). A managed-system ARC Irene trial of named indigenous breeds (Grobbelaar, Sutherland & Molalakgotla 2010, Animal Genetic Resources 46: 25-32) reports far higher counts, now separately verified and retained in productionReferences; managed conditions cannot replace this village range.

Tshupiso ya dilemo/mokgwa: The ARC Irene trial compared Potchefstroom Koekoek, Venda, Ovambo and Naked Neck under commercial feed, water and controlled lighting. Mean eggs/hen in a 52-week cycle were 195.9, 153.7, 125.5 and 138.9 respectively; first-egg ages were 138.5, 139.0, 134.5 and 129.0 days from hatch, with no significant first-egg age difference among breeds. This is managed breed context, not the yield of scavenging hens in KZN. Match feeding and housing before using a benchmark.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Mekgweng ya metseng ya ho rua sebakeng se seholo se bulehileng, dikgoho di tshwanetse ho ipatlela metsi le dijo.

Dikgoho tsa metseng tse ipatlelang dijo di hola butle ho feta tse ruwang kgwebong mme di kotsing e kgolo ya mafu, dikokwana-hloko tse phelang ho tsona le dibatana.

Phuputsong ya Limpopo le KwaZulu-Natal, 94.9% ya balemi ba ne ba kenya dikgoho tsa bona ka matlung bosiu.

Dijo tseo dikgoho tse tshehadi tse itsamaelang di neng di ipatlela tsona di ne di lekana ho hlokomela mmele le kgolo empa di sa lekane bakeng sa ho behela mahe: protheine dihleng tsohle, calcium le phosphorus mariha.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Village chickens (indigenous, free-range) (e tswela pele)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Mohlodi: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 471

Mohlodi: ARC, Climate-Smart Agriculture Training Manual, Module 11: Poultry Production, p. 468

Mohlodi: Malatji, Tsotetsi, van Marle-Koster & Muchadeyi (2016), 'A description of village chicken production systems and prevalence of gastrointestinal parasites: Case studies in Limpopo and KwaZulu-Natal provinces of South Africa', Onderstepoort Journal of Veterinary Research 83(1)

Mohlodi: Raphulu, Jansen van Rensburg & van Ryssen (2015), 'Nutrient content of crop contents of free ranging indigenous chickens raised under village conditions', South African Journal of Animal Science 45(2), p. 1

Mohlodi: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Mohlodi: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Mohlodi: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: Grobbelaar, Sutherland & Molalakgotla (2010), Egg production potentials of certain indigenous chicken breeds from South Africa: methods pp27–28, first-egg p29, Table 4 p31, p. 31

Mohlodi: Tenza, Mhlongo & Chimonyo (2023), 'Village chicken production and egg quality in dry and wet, resource-limited environments in KwaZulu-Natal, South Africa', South African Journal of Animal Science 53(6), p. 5

Mohlodi: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Mohlodi: FAO: Small-scale poultry production, egg marketing

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Laying hens (commercial)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: mahe. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bakeng sa tlhahiso ya mahe, kopa dikgoho tse nyane tse entilweng tse haufi le ho behela ho mofani wa dikgoho tse behelang, ka direkoto tsa dilemo, phepo le ente.

Rera dijo tse leka-lekaneng, metsi a hlwekileng, moriti, moya le tshireletso ho dibatana. Phepelo ya mahe e itshetlehile ka dinonyana tse behelang, dilemo tsa tsona, bophelo le tlhokomelo; hoko feela ha e fane ka mahe.

Bokella mahe kgafetsa, boloka moo dihlaha di alwang ho hlwekile ho omme, mme o tshware/boloke mahe ka hloko bakeng sa tshebediso ya lelapa kapa thekiso.

Tshupiso ya nako: Ka letsatsi la kganya le sa fetoheng la dihora tse 14–16, dikgoho tsa mahe tsa kgwebo di tswela pele ho behela ka tekanyo e phahameng selemo sohle. Tshupiso ena ya taolo ya kgwebo ha e bolele phepelo ya mahe ya lelapa e lekanang ka tlhaho.

Tshupiso ya sehlahiswa sa pele: dibeke tse 21. Dilemo tsa nonyana ho tloha ho qhotswa; netefatsa mofuta le kgodiso le mofani. Dinonyana tse haufi le ho behela di qala hamorao potolohong ena.

Tshupiso ya tlhahiso: 407.1–428.6 mahe/ka kgoho e tshehadi e kentsweng ntlong (hen housed), potoloho ya ho behela ho fihlela dibeke tse 90. Table 4a's cumulative 'Eggs/hen housed' for five commercial strains; the same table gives 'Age at depop. weeks 90.0' for every strain. Commercial barn/cage average, not a free-range smallholder trial. Kept as the source gives it: a per-year rate would need a point-of-lay age from another document.

Tshupiso ya dilemo/mokgwa: SAPA 2024 commercial strain standards cumulative to 90 weeks: Dekalb Amberlink 429 eggs/hen housed, Dekalb Brown 428, Hy-Line Silver Brown 435, Hy-Line Brown 434 and Lohmann Brown Lite 413. These are full-cycle objectives, not eggs per calendar year or a smallholder forecast. Purchase age, actual onset of lay, surviving laying hens, feed, water and lighting need separate records.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Mekgweng ya ho rua e dumellang ho itsamaela kantle, dikgoho tse tshehadi di tshwanetse ho fumana sebaka sa kantle bonyane dihora tse 8 ka letsatsi ha ho kganya.

Moo ho sebediswang mabokose a dihlaha ka bonngwe, ho be le bonyane sehlaha se le seng bakeng sa dikgoho tse tshehadi tse 8.

Dikgoho tse tshehadi di hloka bonyane dihora tse 8 tsa lefifi le sa kgaotsweng potolohong e nngwe le e nngwe ya dihora tse 24 hore di phomole hantle.

Ho rua fatshe ka tlung ho hloka dintho tsa ho ala bonyane karolong e le nngwe ho tse tharo (33%) ya fatshe.

Sebaka sa kantle sa ho itsamaela ha se a tshwanela ho tlala haholo: ho se be le dikgoho tse fetang 5 ka mithara ya sekwere ya sebaka seo.

An owner or manager whose animals are, or may be, infected with a controlled animal disease must report it to the state veterinary director immediately.

Newcastle disease and notifiable avian influenza are on the state list of controlled and notifiable animal diseases.

No animal may be slaughtered anywhere but an abattoir, with the exceptions below.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Dikgoho: Laying hens (commercial) (e tswela pele)

Slaughter for your own household, or for cultural or religious purposes, is exempt from the abattoir rule.

Mohlodi: SAPA Abridged Code of Practice: Commercial Layers, p. 1

Mohlodi: Animal Diseases Act 35 of 1984, s.11(1)(b)(ii), p. 9

Mohlodi: Department of Agriculture, Forestry and Fisheries, Controlled and Notifiable Animal Diseases (factsheet), p. 1

Mohlodi: Meat Safety Act 40 of 2000, s.7(1)(a), p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.7(2)(a), p. 6

Mohlodi: SAPA Industry Profile 2024, Table 4: commercial standards cumulative to 90 weeks, p. 39

Mohlodi: SAPA Layer Notes, Commercial Layers (training module, November 2013), p. 13

Mohlodi: SAPA Industry Profile 2022, Table 4a (Breed standards in laying hens), p. 39

Mohlodi: ARC Climate-Smart Agriculture Training Manual, Poultry production, p. 8

Mohlodi: ARC Climate-Smart Agriculture, module 11, printed pp. 468–480

Mohlodi: FAO: Small-scale poultry production, egg marketing

Mohlodi: SAPA Layer Notes, Commercial Layers (training module, November 2013)

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Matata: Meat duck (Pekin / Muscovy, backyard)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: nama. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Tshupiso ya sehlahiswa sa pele: dibeke tse 7–12. P44 = Pekin (7 weeks to slaughter); MR71 = Muscovy (10 weeks for ducks/hens, 12 weeks for drakes). Treated as 'time to first product' = age to slaughter, since this is a single-cycle meat enterprise (no repeat product like eggs). Non-South-African trial by Warsaw researchers, published in SAJAS; journal location does not establish local applicability.

Tshupiso ya tlhahiso: 2.3–2.9 kg ya boima bo phelang/ka nonyana nakong ya ho hlajwa. Non-SA (Guyana study reporting a citation to Steklenev 1990); no SA-specific absolute slaughter live-weight figure was found (see gaps). 2.93 kg = Pekin at 8-10 weeks, 2.29 kg = Muscovy at 8-10 weeks - matches this enterprise's two breeds. The same LRRD paper's own experiment (Table 2, intensive system, Guyana) measured live weight at slaughter (8 weeks) as: Pekin drakes 2.53 kg, Pekin hens 2.29 kg, Muscovy drakes 2.47 kg, Muscovy hens 1.51 kg - broadly consistent with, but not merged into, the cited range above since it is a separate dataset/table in the same source. No sourced eggs/duck/year figure was found for either breed (see gaps); the LRRD and SAJAS sources used here studied ducks reared only for meat.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

Ke tlolo ya molao ho hanela letata dijo kapa metsi, kapa ho le boloka moo ho se nang sebaka se lekaneng, moya o lekaneng, kganya kapa tshireletso mochesong le serameng.

Sheba mehlapeng ho shwa ha tshohanyetso, masiba a hlakorileng, ho ruruha ha dinama tse leketlang tlasa molomo kapa dikama tsa hloohong, kapa ho theoha ha tshohanyetso ha tlhahiso ya mahe - ana ke matshwao a mokakallane wa dinonyana mme a tshwanetse ho tlalehwa hang-hang.

Boloka matata hole le dinonyana tse hlaha le metsi kapa dijo tse pepeneneng tse di hohelang, mme o laole ho kena ha baeti matlong a matata, ho fokotsa kotsi ya mokakallane wa dinonyana.

Ducks are one of the domesticated animal species listed in the Meat Safety Act, so its rules on registered abattoirs and hygiene apply to them.

Ducks slaughtered to be sold must be slaughtered at a registered abattoir; slaughter for the farmer's own consumption is exempt but that meat may not be sold.

At the abattoir, ducks must be handled humanely during loading, transport, offloading, housing, immobilising and killing, under the Animals Protection Act.

Ducks are birds/poultry, so an outbreak of avian influenza in a duck flock is a controlled animal disease under the Animal Diseases Act and must be reported to the state vet immediately.

Mohlodi: Animals Protection Act 71 of 1962 (consolidated text, s.2(1)(b)-(c)), p. 2

Mohlodi: Western Cape Department of Agriculture, Avian Influenza (Bird Flu) factsheet, p. 2

Mohlodi: Meat Safety Act 40 of 2000, Schedule 1 (Animals to which this Act applies), p. 16

Mohlodi: Meat Safety Act 40 of 2000, s.7, p. 6

Mohlodi: Meat Safety Act 40 of 2000, s.9, p. 7

Mohlodi: Damaziak et al. (2014), Influence of housing system on the growth and histological structure of duck muscles, S. Afr. J. Anim. Sci. 44(2): Poland trial

Mohlodi: Livestock Research for Rural Development 18(11) 2006 - Solomon et al. (Guyana; not an SA figure), citing Steklenev 1990

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Letangwana: Tilapia (Mozambique tilapia, small pond or tank)

Meaho e 1 e teng; e 2 e sisintsweng. Kgwebo e kgethilweng: tlhapi. Ha ho nahannwe palo ya diphoofolo kapa tlhahiso.

Keletso ya ho reka: bapisa mofuta/diphoofolo le sehlahiswa, dijo tsa sebaka, maemo a lehodimo le bokgoni ba tlhokomelo; netefatsa le moeletsi wa sebaka wa mehlape kapa dinotshi.

Tshupiso ya sehlahiswa sa pele: dibeke tse 43.5–52.2. 10 months x 4.35 = 43.5 weeks; 12 months x 4.35 = 52.2 weeks. This is Mozambique Tilapia (O. mossambicus) grow-out time to harvestable size; the source separately states Nile Tilapia (O. niloticus) reaches harvestable maturity faster, in 6-10 months (= 26.1-43.5 weeks) — not used here as it is a different species with a different legal status (see 'legal').

Tshupiso ya tlhahiso: 800 g ya boima nakong ya kotulo/ka tlhapi. Table 8-3 gives the Mozambique Tilapia's harvestable size as an approximate single figure ('+- 800g'), not a range, so [800, 800] is used rather than inventing a spread. The same table cell wraps '10-12' across a line break in the PDF ('10-\n12'), kept verbatim. The table's Nile Tilapia row gives the same approximate weight (+-800g) reached faster, in 6-10 months — not used here because this dossier is framed around the indigenous, legally unrestricted Mozambique Tilapia per task instruction; see 'legal' for why Nile Tilapia is treated separately. No SA source was found giving a feed-conversion-based or per-hectare yield figure specific to O. mossambicus that could be converted to this per-fish unit without stitching numbers across documents, so outputUnit follows the source's own unit (grams harvest weight per fish) rather than being converted.

Tshupiso ya dilemo/mokgwa: ARC makes tilapia suitability depend on water temperature: summer grow-out can be suitable anywhere in South Africa when water exceeds 20°C; growth is poor below 18°C and winter water below 13°C can kill farmed tilapia. The Highveld reference is a summer crop; tunnels can widen the range. Measure pond water and confirm stocking species and dates. Air climate or a mapped pond does not prove harvest months, fish numbers or food kg.

Hlahloba ditshupiso tse qaqileng tsa kgwebo bakeng sa dijo le bodulo. Bapisa dijo le sebaka le boemo ba phoofolo le mokgwa wa bodulo; fana ka metsi a hlwekileng a tshepahalang.

The indigenous Mozambique Tilapia (O. mossambicus) is not a listed invasive species and may be farmed without an alien-species permit; the faster-growing Nile Tilapia (O. niloticus) is.

Nile Tilapia (Oreochromis niloticus) is a nationally listed invasive fish species: Category 1b inside National Parks, Provincial Reserves, Mountain Catchment Areas and Forestry Reserves, and Category 3 in all other areas.

Outside protected areas Nile tilapia is Category 3: it may remain in prescribed areas or provinces, but further propagation or trade is prohibited.

Nile Tilapia has been listed under the NEMBA Alien and Invasive Species Regulations as requiring a permit to produce.

Producing an alien or invasive species (such as Nile Tilapia) is prohibited in areas designated under the Protected Areas Act and the Fish Sanctuary Act; a permit is required to produce it elsewhere.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.

SYNTHETIC GRAPH SOURCE {COUNT} $&

Dikgetho tsa polasi - di tswela pele

Sebedisa seo o se boneng polasing le manane a makgutshwane a dipatlisiso. Netefatsa dikgetho sebakeng seo pele o reka.

Dihlahiswa tsa diphoofolo le tlhokomelo (e tswela pele)

Letangwana: Tilapia (Mozambique tilapia, small pond or tank) (e tswela pele)

Producing tilapia in South Africa requires a water use licence under the National Water Act.

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), Table 8-3 (Mozambique vs. Nile Tilapia), p. 69

Mohlodi: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), List 7 (National List of Invasive Fresh-water Fish Species), p. 32

Mohlodi: Dept. of Environmental Affairs / invasives.org.za, National Listed Invasive Species (NEMBA AIS Regulations booklet), p. 3

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 69

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 74

Mohlodi: IDC, Research into the Potential for the Production, Processing and Export of Tilapia for the Southern African Market (2015), p. 75

Mohlodi: ARC Aquaculture Production App: General FAQ, Can I grow tilapia on the Highveld?

Mefuta ya dijalo e lokelang ho hlahlojwa sebakeng seo

Boloka lebitso la mofuta oo o o kgethang ka nnete. Manane a makgutshwane a dipatlisiso a hloka netefatso bakeng sa nako ya sebaka, metsi le mobu.

Tomatoes

Mofuta o rekotilweng: ha ho eso kgethwe.

Maemo a lehodimo a sebaka sena ha a eso hlake, kahoo ha ho bolelwe lenane le lekgutshwane la sebaka. Bula karete ya sejalo ho bona mehlodi le ho rekota kgetho ya hao.

Ha o a hlahlojwa ke setsebi sa temo — ke tataiso ya ho rera, ha se taelo. O bontshe moeletsi wa hao wa temo, mme o etse teko ya mobu pele o sebedisa tjhelete mobung.
Phetolelo ya moralo: ha e so hlahlojwe ke motho ya tsebang puo hantle. Hlahloba mohlodi wa Senyesemane pele o etsa se bolelwang.