

Table S2. Results of the linear mixed model (LMM) assessing fluctuating asymmetry (FA) and directional asymmetry (DA) into native and neonative populations. Statistical significance of Side suggest DA whereas the interaction Side:Individual suggest FA. HUL: humeral length; FAL: forearm length; FL: femur length; TL: tibia length; FTL: hindfoot length.

Neonative					
Trait	Model	numDF	denDF	F-value	p-value
FL	Side	1	32	16.603	0.0003*
	Side:Id	60	32	2.532	0.0027*
TL	Side	1	32	10.472	0.0028*
	Side:Id	0	32	1.782	0.0391*
FTL	Side	1	32	0.359	0.5535
	Side:Id	60	32	1.449	0.1281
HUL	Side	1	32	1.694	0.2024
	Side:Id	60	32	2.072	0.0137*
FAL	Side	1	32	9.991	0.0034*
	Side:Id	60	32	1.668	0.0591
Native					
Trait	Model	numDF	denDF	F-value	p-value
FL	Side	1	22	12.060	0.0022*
	Side:Id	40	22	1.836	0.0652
TL	Side	1	22	0.841	0.3691
	Side:Id	40	22	0.990	0.5251
FTL	Side	1	22	20.185	0.0002*
	Side:Id	40	22	3.195	0.0025*
HUL	Side	1	22	2.754	0.1112
	Side:Id	40	22	0.810	0.7248
FAL	Side	1	22	5.994	0.0228*
	Side:Id	40	22	1.771	0.0772

* values < 0.05.