

PRECISION MEASUREMENT OF THE HIGGS BOSON MASS AND SEARCH FOR  
DILEPTON MASS RESONANCES IN  $H \rightarrow 4\ell$  DECAYS USING THE CMS DETECTOR AT  
THE LHC

By

JAKE ROSENZWEIG

A DISSERTATION PRESENTED TO THE GRADUATE SCHOOL  
OF THE UNIVERSITY OF FLORIDA IN PARTIAL FULFILLMENT  
OF THE REQUIREMENTS FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY

UNIVERSITY OF FLORIDA

2022

© 2022 Jake Rosenzweig

This work is dedicated to the living and loving memory of Jacob Myhre.

## ACKNOWLEDGEMENTS

Without so many two- and four-legged blessings along the way, I could never have made it to this point in my academic career. Thus, I begin by giving my endless gratitude to my high energy physics mentors, Professors Andrey Korytov and Guenkah Mitselmakher, for granting me this one-of-a-kind opportunity to do real *science* at CERN.

To my wife, Suzanne Rosenzweig, for showing me that dreams *do* come true. To my mother and father, Vicki and John, who always reassured me that I could achieve anything I put my mind to. Sleep peacefully, Mom. To my siblings, Alex, Ryan, Devin, Jace, and Claudia who frequently and gently reminded me that life existed outside of grad school. To Auntie Rachel and Uncle Yuri, who just *get* me and have given to me everything I could have possible asked for. To my mentor, Sheldon Friedman, and his wife, Rita Friedman (Rosenzweig), who chose to invest in my success at a young age. I have only made it this far thanks to their undying encouragement, love, and optimism. To Sheldon's best friend, Dr. Bernard Khoury, whose reputation and has helped pave my own path. To the many moms who generously gave unconditional support during the darkest of times and unequivocal love during the brightest times: Cyndi Reilly-Rogers, Dawn Hood, Margaret Sherrill, and Silet Wiley.

To Dr. Filippo Errico for his focus, leadership, selflessness, and patience in leading the Higgs mass analysis. To Dr. Lucien Lo for showing me the simplicity and beauty of Python in his typical laid-back way and for leading the dilepton analysis. To Dr. Noah Steinberg for showing me how majestically physics can be communicated from mind to mind. To Dr. Darin Acosta, for spending many hours of physics discussion with me and the other students, who have helped us build our "CMS Office Hours" Ari Gonzalez, Cris Caballeros, Jeremiah Anglin, Sean Kent, Evan Koenig, Neha Rawal, Nik Menendez, John Rötter. To the gents who paved my way to and through the world of CMS, Brendan Regnery and Bhargav Joshi. To my mentee, Matthew Dittrich, for accepting the baton of knowledge and making everything come full circle.

To my comrades for showing me what it takes to survive the core courses, Dr. Atool Divakarla, Dr. Brien O'Brendan, Dr. Donyell Guerrero, and Dr. Vladinar Martinez. To my quondam friend, Adamya Goyal, for his gentleness, humility, patience, and tutorage. To my Polish roommates in Saint-Genis-Pouilly for showing me what home away from home feels like:

Bartoszek, Dziadzius, Karolina, and Sandruśa. To the boys who have been there since the beginning: Jish, Willis, The Shane, Zacman, Duck, and Marcus for their clever competition and continual camaraderie which has shaped me to this day.

To Big Tree who stood as a symbol of strength, beauty, and life for centuries before us. As Irma's wild whirring winds worsened, the cacophony of ripping roots resounded throughout the western corridor. There I stood in that frozen moment—awestruck, speechless—watching her *fall* helplessly towards the physics building. What could have been a catastrophe of cataclysmic proportions was instead a gentle grazing against the north windows of NPB where, there, she was gracefully laid to rest.

Finally, to Existence, for this unpredictable, unbelievable blip of an experience called Life.

## TABLE OF CONTENTS

	<u>page</u>
ACKNOWLEDGEMENTS .....	4
LIST OF TABLES.....	7
LIST OF FIGURES.....	8
ABSTRACT.....	9
CHAPTER	
1 INTRODUCTION .....	10
2 CONCLUSION.....	14
REFERENCES .....	14
BIOGRAPHICAL SKETCH .....	16

## LIST OF TABLES

Tables

page

## LIST OF FIGURES

<u>Figures</u>	<u>page</u>
1-1 The elementary particles described by the SM. ....	10
1-2 Theoretical stability regions of our universe.....	12



Abstract of Dissertation Presented to the Graduate School  
of the University of Florida in Partial Fulfillment of the  
Requirements for the Degree of Doctor of Philosophy

PRECISION MEASUREMENT OF THE HIGGS BOSON MASS AND SEARCH FOR  
DILEPTON MASS RESONANCES IN  $H \rightarrow 4\ell$  DECAYS USING THE CMS DETECTOR AT  
THE LHC

By

Jake Rosenzweig

August 2022

Chair: Andrey Korytov

Co-Chair: Guenakh Mitselmakher

Major: Physics

The mass of the Higgs boson is measured in the  $H \rightarrow ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ ) decay channel and is found to be  $m_H = 125.38 \pm 0.11$  GeV; the world's most precise measurement to date. The data for the measurement were produced from proton-proton (pp) collisions at the Large Hadron Collider with a center-of-mass energy of 13 TeV during Run 2 (2016–2018), corresponding to an integrated luminosity of  $137.1 \text{ fb}^{-1}$ , and were collected by the Compact Muon Solenoid experiment. This measurement uses an improved analysis technique in which the final state muon tracks are constrained to originate from the primary pp vertex. Using data sets from the same run, a search for low-mass dilepton resonances in Higgs boson decays to the  $4\ell$  final state is also conducted. No significant deviation from the Standard Model prediction is observed.

# CHAPTER 1

## INTRODUCTION

The Universe, while overwhelmingly vast, is built from a remarkably few kinds of elementary (i.e., indivisible) particles. As shown in Fig. 1-1, any elementary particle can be classified into 1 of these 3 categories: matter particles (*fermions*), force-carrying particles (*gauge bosons*), and Higgs bosons. There are 12 kinds of fermions which can be split evenly into 2 groups, depending on with which forces they interact: those that interact via the electromagnetic (EM) force and the weak nuclear force are classified as *leptons*, of which there are 6 kinds (“*flavors*”), whereas those that interact via the EM, weak, *and* strong nuclear forces are classified as *quarks*, of which there are also 6 flavors. There are 4 kinds of gauge bosons, each of which is a force carrier for a specific force (the gluon is said to *mediate* the strong force, the photon mediates the EM force, while both the  $W^\pm$  and Z bosons mediate the weak force). Thus, all the diversity and manifestations of reality come from only 17 kinds of “building blocks”. It is the mission of particle physicists to understand the underlying mathematical structure that describes Nature in as accurate—and hopefully *concise*—a theory as possible. The best theory that stands today is called the Standard Model (SM) of particle physics.




Figure 1-1. The elementary particles described by the SM.

The SM has bore witness to many triumphs over its approximately 70-year development: it predicted the existence of quarks which were experimentally confirmed in the mid-1970s; it predicted the existence of the tau neutrino which was experimentally confirmed in 2000; its most groundbreaking prediction was experimentally confirmed on July 4, 2012—almost exactly 10 years ago from the date of this dissertation writing—when the ATLAS and CMS collaborations announced the discovery of the Higgs boson [1, 2, 3].

Quantum Field Theory is the mathematical and conceptual backbone of the SM. Within its framework, all particles are *excitations* of their corresponding field so, e.g., *every* electron in the Universe is thought to be an excitation of the single electron field that permeates all of spacetime. The existence of the Higgs boson (H) suggests that its corresponding field exists—the *Higgs field*—and that H is the quantum excitation of that field. This all-pervasive Higgs field and its corresponding boson are generated mathematically via the Brout-Englert-Higgs (BEH) mechanism. Most SM particles—except for neutrinos, photons, and gluons—“acquire” their mass by interacting with the Higgs field. This is also how the Higgs boson itself acquires its mass ( $m_H$ ): by interacting with its own field!

The masses of SM particles *depend* on the value of  $m_H$  . . . *so what is its value?* Unfortunately,  $m_H$  is a free parameter of the SM, so theory is unable to provide a value for  $m_H$  based solely on other fundamental constants. Instead, the value of  $m_H$  must be measured by experiment. TODO:CITE LOTS OF PEEPS [?] Although the value of  $m_H$  was measured back in 2012, it is always important to improve the measurement by lowering the uncertainties on (i.e., increasing the precision on) the mass value. A more precise value of  $m_H$  improves theoretical limits on the masses of most of the other elementary particles and may very well determine the ultimate stability of our universe, as shown in Fig. 1-2.

Plus, as mentioned previously, Furthermore, the value of  $m_H$ .

The production of a Higgs boson is only feasible at conditions close to those thought to exist at the beginning of the Universe. This grand achievement is accomplished frequently by the Large Hadron Collider (LHC), located on the border of France and Switzerland. It is the largest and




Figure 1-2. (Left) Theoretical stability regions of our universe based on the pole masses of the top quark ( $m_t^{\text{pole}}$ ) and Higgs boson ( $m_h^{\text{pole}}$ ). (Right) A closeup of the SM region of the left plot. The contours represent the 68%, 95%, and 99% confidence levels based on the experimental uncertainties of  $m_t^{\text{pole}}$  and  $m_h^{\text{pole}}$ . Plots taken from [4] and units added to all axes.

most powerful proton-proton (pp) collider ever made. The LHC accelerates protons to near-light speeds and produces pp collisions with center-of-mass energies as high as 13 TeV, Old and newly produced particles spew out of the collisions points and are analyzed by detectors like the aforementioned ATLAS and CMS experiments. These enormous particle detectors have thousands of scientists performing dozens of analyses to look for hints of beyond Standard Model (BSM) physics, extra dimensions, miniature black holes, and more. This dissertation utilizes data collected by the CMS experiment during the LHC Run 2 (2016–2018) to perform a precision measurement of the Higgs boson mass ( $m_H$ ).

The measurement of  $m_H$  presented in this dissertation utilizes the following improvements compared to previous measurements:

- Utilizing nearly four times as much collected data from Run 2 ( $\mathcal{L}_{\text{int}} = 137.1 \text{ fb}^{-1}$ ) vs. the data used for the 2016 measurement ( $\mathcal{L}_{\text{int}} = 35.9 \text{ fb}^{-1}$ ).
- Four final-state categories are used:  $4\mu$ ,  $4e$ ,  $2e2\mu$ ,  $2\mu2e$ . In previous measurements, the last two final states (the mixed-flavor states) were combined, when truly they have different

kinematical properties (depending on into which lepton pair the  $Z_1$  decayed): different peak widths (instrumental resolutions), different signal efficiencies, and different relative levels of reducible background.

- Ultra-Legacy (UL) reconstruction is used for muon, electron, photon, and jet tracks. This significantly improves electron momenta and improves the other particle momenta, though to a lesser degree.
- The measurements of muon  $p_T$  are improved by constraining the muon tracks to originate from the interaction vertex (also called a *vertex constraint*).
- When extracting the value of  $m_H$  in past measurements, a 3D pdf  $\left(m_{4\ell}, \sigma_{m_{4\ell}}, \mathcal{D}_{\text{bkg}}^{\text{kin}}\right)$  was built into a factorized form  $f\left(m_{4\ell}, \sigma_{m_{4\ell}} \mid m_H\right) \cdot g\left(\mathcal{D}_{\text{bkg}}^{\text{kin}} \mid m_{4\ell}\right)$ , which was later found to contain an existing correlation between  $\sigma_{m_{4\ell}}$  and  $\mathcal{D}_{\text{bkg}}^{\text{kin}}$ . To account for this correlation, now the events are split into 9 categories based on the per-event *relative* mass uncertainty  $\left(\frac{\sigma_{m_{4\ell}}}{m_{4\ell}}\right)$  and, for each, a 2D pdf  $\left(m_{4\ell}, \mathcal{D}_{\text{bkg}}^{\text{kin}} \mid m_H\right)$  is built.
- The systematic uncertainties on electron and muon momentum scales ( $p_T^{e,\mu}$ ) are reduced, thanks to a more detailed analysis on the uncertainties. This has the additional effect of significantly reducing the uncertainty on the per-event four-lepton mass resolution.

The following chapters of this dissertation begin with a thorough description of the CMS Experiment and its composite subdetectors in Chapter ???. Next, the details of the precision measurement of the Higgs boson mass using the LHC Run 2 data is discussed in Chapter ??. Finally, the results of the Higgs boson mass measurement analysis is summarized in Chapter 2.

## CHAPTER 2

### CONCLUSION

We measured it, boys.

## REFERENCES

- [1] S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Physics Letters B* **716** (2012) 30.
- [2] S. Chatrchyan, V. Khachatryan, A.M. Sirunyan, A. Tumasyan, W. Adam, T. Bergauer et al., *Observation of a new boson with mass near 125 GeV in pp collisions at  $\sqrt{s} = 7$  and 8 TeV*, *Journal of High Energy Physics* **2013** (2013) 81.
- [3] G. Aad, T. Abajyan, B. Abbott, J. Abdallah, S. Abdel Khalek, A.A. Abdelalim et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Physics Letters B* **716** (2012) 1.
- [4] A. Andreassen, W. Frost and M.D. Schwartz, *Scale-invariant instantons and the complete lifetime of the standard model*, *Phys. Rev. D* **97** (2018) 056006.

## BIOGRAPHICAL SKETCH

This is the “biographical sketch” tex file, which should have been set in the main file using the command `\setBiographicalFile{Drive:/file/location/biographyFile}`.

Keep in mind this should be written in **third person** and should assume you have already completed your degree that you are writing this thesis or dissertation for. For example:

Jason is a person that wrote some code, did some research, and eventually got a PhD in mathematics for some stuff. He had to actually write a real biographical sketch because he had forgotten to do it until final submission.