



CNOT7 two-phases




Fit Residuals




Fit Residuals

Fit Residuals




RNA populations for CNOT7_two-phases


2.0 μM enzyme



3.0 μM enzyme




5.0 μM enzyme




[Enzyme] = 2.0 μ M

Dense phase:
15.8 μ M enzyme
424.9 nM RNA




Length polyA tail


[Total cleavable RNA]
in Dense phase



[A1]
in Dense phase




Dilute phase:
0.9 μ M enzyme
33.5 nM RNA




Length polyA tail

[Total cleavable RNA]
in Dilute phase




[A1]
in Dilute phase



Time (s)


[Enzyme] = 3.0 μ M

Dense phase:
25.3 μ M enzyme
437.0 nM RNA




Length polyA tail


[Total cleavable RNA]
in Dense phase



[A1]
in Dense phase




Dilute phase:
1.1 μ M enzyme
35.8 nM RNA




Length polyA tail

[Total cleavable RNA]
in Dilute phase




[A1]
in Dilute phase



Time (s)


[Enzyme] = 5.0 μ M

Dense phase:
42.4 μ M enzyme
462.7 nM RNA




Length polyA tail


[Total cleavable RNA]
in Dense phase



[A1]
in Dense phase




Dilute phase:
1.3 μ M enzyme
41.0 nM RNA




Length polyA tail

[Total cleavable RNA]
in Dilute phase



[A1]
in Dilute phase



Time (s)