



However, heavy blood loss is not common and the volume of blood resuscitation is often restricted in patients with isolated TBI, suggesting that these factors may not be critical in TBI-associated coagulopathy. This review discusses the unique biochemical and cellular characteristics of the brain that make it uniquely procoagulant and prone to coagulopathy.

### Platelets and Platelet-activating Factor

Hemostasis begins when platelets are captured to the sub-endothelial matrix exposed at sites of vascular injury to form a platelet-rich plug that is subsequently stabilized by fibrin cross-linking resulting from activating coagulation cascade and thrombin generation (Fig. 1). Platelet functions after TBI have not been systemically investigated, but circumstantial evidence, such as low platelet counts without heavy blood loss and spontaneous platelet aggregation,<sup>25,26</sup> suggests that TBI may result in platelet hyperactivity, a condition of platelets being partially activated or primed for activation. Although reported to be in normal range, platelet aggregation was often induced by agonists at maximal dosages that would not distinguish between normal and hyperactive platelets.<sup>28,29</sup> The presence of platelet hyperactivity is also supported by the finding of intravascular microthrombosis, now considered to be more common in TBI patients than clinically recognized.<sup>9,23,30,31</sup> Enhanced platelet aggregation and microthrombosis in the lungs

and brain have been verified in animal models of TBI.<sup>23,32,33</sup> These brain microthrombi are mostly detected in the peri-contusion cortex<sup>34</sup> and contain platelets,<sup>23,34</sup> fibrin,<sup>32</sup> and von Willebrand factor (VWF).<sup>34</sup>

Multiple factors could make platelets hyperactive. Among them, platelet activating factor (PAF, 1-O-hexadecyl-2-acetyl-sn-glycero-3-phosphocholine) may play a unique role. PAF was first identified as a basophil-derived molecule that induces aggregation and serotonin release from rabbit platelets.<sup>35,36</sup> It was subsequently found to be synthesized and secreted from many cell types through constitutive and induced pathways.<sup>37</sup> Cells synthesizing PAF are often also targeted by it, making PAF one of the classic autacoids. PAF acts on target cells through a G-protein-coupled seven trans-membrane receptor.<sup>37,38</sup> Platelets have two types of PAF receptors: one with a high affinity (kDa, 37, 10<sup>-13</sup> nM) and low abundance (1399/498 sites/platelet) and the other with a low affinity and high abundance. Upon binding to the high-affinity receptor, PAF initiates intracellular signals to activate phospholipases C and A2 that hydrolyze phosphoinositide to release arachidonic acid, mobilize cytosolic calcium, and activate protein kinase C in platelets.<sup>37</sup> Leukocytes<sup>39,40</sup> and vascular endothelial cells<sup>42,43</sup> also express one or both receptors for PAF. The brain and spinal cord are a richer source of PAF than other tissues<sup>44,45</sup> and response to PAF through a G protein-coupled neuronal PAF receptor that is sensitive to pertussis toxin.<sup>47,48</sup> Neural cells release PAF during cerebral




FIG. 1. Schematic of hemostasis and coagulation. Hemostasis is initiated by an interaction between von Willebrand factor (VWF) in the subendothelium and the GP Ib-IX-V complex on platelets that slows down the movement of platelets to engage other ligand-receptor interactions with slower on-rates. Platelets are rapidly activated and only adhere to the subendothelium through an interaction between collagen matrix and GP VI and the integrin  $\alpha 2\beta 1$ . Activated platelets aggregate by fibrinogen crosslinking the integrin  $\alpha IIb\beta 3$ . They also provide a phosphatidylserine (PS)-rich surface on which tissue factor forms a complex with coagulation factor VIIa to initiate the extrinsic pathway of coagulation. Thrombin generated through this pathway cleaves fibrinogen to fibrin that forms laterally associated fibrils to stabilize platelet aggregation. Several procoagulant molecules are enriched in the brain and key steps of hemostasis and coagulation are uniquely enhanced in TBI.

ischemia<sup>50,52</sup> and tissue hypoxia.<sup>53</sup> Lindsberg and colleagues<sup>54</sup> reported that the levels of PAF were increased by approximately 20-fold from baseline in rats with sustained ischemic and reperfusion injury to the spinal cord.

As an autacoid, extensive studies in the past have been focused on PAF effects on neurological functions and inflammation.<sup>55,56</sup> A role of PAF in developing TBI-associated coagulopathy remains largely unknown, but should be expected because it is a potent platelet agonist. In addition, PAF also contributes to the hypoxia-induced breakdown of the blood brain barrier,<sup>58</sup> potentially resulting in the release of PAF and other brain-derived prothrombotic molecules to the systemic circulation. Several PAF antagonists are reported to attenuate ischemic edema, early post-ischemic hyperemia, and intravascular microthrombosis in animal models.<sup>23,59,60</sup> It is, however, not known whether these PAF antagonists improve clinical or experimental outcomes by reducing PAF-associated neurotoxicity or coagulopathy. Further, the pre-injury use of aspirin, which blocks platelet aggregation induced by arachidonic acid, a polyunsaturated omega-6 fatty acid released from platelets and neural membrane, and the platelet ADP P2Y<sub>12</sub> receptor antagonist clopidogrel failed to improve clinical outcome of TBI.<sup>61,62</sup> These apparent conflicting data regarding the clinical efficacy of platelet antagonists beg for answers to a key question of whether globally inhibiting platelets improve clinical outcomes of TBI without aggravating the risk for intracranial bleeding. In this regard, specifically blocking PAF activity may prevent TBI-induced platelet hyperactivity without inhibiting systemic hemostasis.

#### Platelet-endothelial Interactions

Similar to platelet hyperactivity, a role played by major platelet ligands in TBI-associated coagulopathy is also not clear. For example, VWF is widely used as a marker for endothelial injury,<sup>63,64</sup> but its change in its adhesion activity and contribution to coagulopathy in TBI are not known. VWF mediates the initial tethering of platelets to perturbed endothelium or subendothelial matrix exposed at sites of vascular injury. By forming a high affinity com-

plex, it also protects coagulation factor VIII from being inactivated by proteolytic degradation in the plasma. VWF is exclusively synthesized in endothelial cells and megakaryocytes/platelets (Fig. 2). VWF multimers are constitutively secreted and stored in the Weibel-Palade bodies of endothelial cells and α-granules of megakaryocytes/platelets.<sup>65,66</sup> The stored VWF multimers are enriched in ultra-large (UL) and prothrombotic forms<sup>65,67</sup> that are released upon stimulation to endothelial cells and platelets. These prothrombotic ULVWF multimers are rapidly and partially cleaved by the zinc metalloprotease ADAMTS-13 (a disintegrin and metalloprotease with thrombospondin type 1 repeats) into smaller multimers<sup>68,69</sup> that are active in hemostasis, but no longer prothrombotic. In the absence of this proteolytic activity, ULVWF multimers form rope-like long strings that tether platelets and leukocytes to endothelium,<sup>71,72</sup> leading to in situ thrombosis and thromboembolism in the downstream microvasculature.

There is no report on the contribution of VWF and ADAMTS-13 to TBI-associated coagulopathy, but several lines of evidence support this possibility. First, the secretion of ULVWF is significantly increased in traumatic injury, including TBI.<sup>63,64</sup> Second, ADAMTS-13 activity is reduced in traumatic and surgical injury. Third, intravascular thrombosis developed in the lesion boundary zone contained a substantial amount of VWF and platelets in a rat model of controlled cortical injury.<sup>30,34</sup> Fourth, endothelial cells in cerebral vascular beds, especially small vessels and microvasculature, synthesize significantly more VWF than vascular beds from other organs.<sup>73</sup> Fifth, VWF/FVIII infusion reduces regional blood flow after controlled ischemia in dogs.<sup>74</sup> The enhanced secretion of ULVWF multimers, coupled with a partial inhibition of their cleavage by ADAMTS-13, may allow accumulation of hyperactive ULVWF multimers on endothelium to promote leukocyte transmigration through the vessel wall and in plasma to agglutinate platelets, leading to thrombosis and tissue ischemia.

In addition to their role in forming initial clots, platelets also provide a phosphatidylserine (PS)-rich surface to initiate and propagate the extrinsic coagulation pathway by forming a complex composed of tissue factor, coagulation factor VIIa, and




FIG. 2. Synthesis and multimerization of von Willebrand factor (VWF). VWF is initially synthesized as a monomeric glycoprotein that then forms a C-terminal disulfide-linked homodimer. VWF dimers subsequently form multimers through N-terminal disulfide linkages after proteolytic removal of a large 714 amino acid propeptide. Once secreted, ULVWF (ultra-large von Willebrand factor) multimers are subjected to cleavage at the A2 domain by ADAMTS-13. The cleaved VWF multimers have variable sizes, with the largest multimers being hemostatically most active.

calcium on a PS surface to activate factor IX and X (Fig. 1).<sup>75</sup> Two key molecules for the extrinsic coagulation pathway are tissue factor and phosphatidylserine, which are richly expressed in cerebral tissue.

#### Tissue Factor and Tissue Factor-Bearing Microparticles

Tissue factor is an integral membrane protein that is enriched in the central nerve system.<sup>76,78</sup> Human tissue factor is a lipoprotein of 263 amino acids containing an extracellular domain, a transmembrane domain, and a short cytoplasmic tail.<sup>79,80</sup> The extracellular domain contains two fibronectin type III domains with two potential disulfide bonds of Cys49-Cys59 and Cys186-Cys209<sup>81</sup> and can form a complex with factor VIIa, preferentially on a PS-rich membrane.<sup>82</sup> The amino acid Cys245 in the cytoplasmic tail is linked to a palmitate fatty acyl chain,<sup>83</sup> suggesting that tissue factor is selectively concentrated in lipid microdomains of the cell membrane.<sup>84,86</sup>

Tissue factor is not normally exposed to circulating blood because it is primarily expressed on fibroblasts and smooth muscle cells in the vessel wall and non-vascular cells such as astrocytes, epidermal cells, and renal glomeruli. Monocytes<sup>87,89</sup> and endothelial cells<sup>90,92</sup> express tissue factor, but only after induction by stimulating factors such as injury and inflammation. There is debatable presence of soluble tissue factor in blood at approximately 100 ng/mL,<sup>93,94</sup> but this blood-born tissue factor does not appear to trigger coagulation under physiological conditions because it may be either encrypted or in an insufficient amount. The latter can be overcome by concentrating tissue factor at a site of vascular injury or in a freshly formed thrombus.<sup>95</sup> A long-standing question is how tissue factor is integrated into the membrane surface of activated platelets, which do not express tissue factor or have an extremely low level,<sup>96,97</sup> to activate the extrinsic coagulation

pathway. One potential mechanism is for tissue factor to be delivered by membrane microparticles.

In response to injury, cells such as platelets, erythrocytes, monocytes/macrophages, and endothelial cells shed membrane fragments through active microvesiculation<sup>98,99</sup> or apoptosis.<sup>100,102</sup> Intracellular signals that increase calcium influx activate the cysteine protease calpain, which then cleaves cytoskeletal proteins  $\alpha$ -actinin,  $\beta$ -tubulin, adducins, spectrin, talin, and actin<sup>103</sup> to disrupt membrane-cytoskeleton and produce microparticles.<sup>104</sup> Inhibiting calpain therefore blocks microparticle production.<sup>104,105</sup> Platelets generate microparticles during storage<sup>106</sup> in response to agonists and changes in hydrodynamic conditions.<sup>107,108</sup> These microparticles are primarily composed of membrane fragments with a limited amount of cytoplasm and measure 0.5–1  $\mu$ m.<sup>109</sup> They are enriched in lipid microdomains (Fig. 3) where cholesterol and sphingolipids, adhesion receptors, and tissue factor are highly concentrated.<sup>84,110</sup> Adhesion receptors on these microparticles could bring tissue factor to cells expressing counter-receptors or ligands for these receptors. For example, P-selectin glycoprotein ligand 1 (PSGL-1) expressed on monocytes could bring tissue factor-bearing microparticles to activated platelets and endothelial cells that express P-selectin.<sup>87,90,95</sup> Immunoelectron microscopy has indeed detected tissue factor-positive membrane vesicles in large clusters near the surface of platelets.<sup>95</sup> Falati and colleagues<sup>111</sup> further showed that fluorescently labeled tissue factor-bearing microparticles derived from mouse monocytes accumulate in the leading edge of a developing thrombus when they were infused into mice prior to vascular injury. The accumulation was drastically reduced when these microparticles were infused into mice that were P-selectin deficient.

Brain tissue is a rich source of tissue factor.<sup>76,78,112,114</sup> In fact, brain homogenate was not used to induce blood coagulation as




FIG. 3. Lipid microdomains in cell membrane. These microdomains (rafts) are cholesterol- and sphingolipid-rich compartmentalized regions of cell membrane where signaling molecules, receptors, and phosphatidylserine (PS) are selectively concentrated. These domains are orderly and tightly packed and can float freely in the membrane bilayer. Upon activation, platelets generate microparticles that are enriched in these lipid microdomains with concentrated adhesion receptors and PS. The latter could carry coagulation factors. Other cells may also produce microparticles that are PS-rich and contain signature molecules from the parental cells that deliver microparticles to target cells.

early as 1939. This rich source of tissue factor may account in part for a higher incidence of coagulopathy in patients with penetrating head injury as compared to those with blunt injury. Further, neural cells are highly sensitive to apoptosis to potentially produce tissue factor-bearing neural microparticles. Microglial cells have indeed been shown to produce microparticles upon ATP binding to the ligand-gated ATP receptor P2X7.<sup>115,117</sup> The microvesiculation is blocked by inhibiting acid sphingomyelinase and found to be extremely low in the culture of glial cells from acid sphingomyelinase null mice.<sup>118</sup> Platelet and endothelial cell microparticles were also detected in blood and cerebral-spinal fluid (CSF) samples from healthy subjects and TBI patients.<sup>119</sup> A high level of microparticles is associated with a low GCS score, a larger hematoma volume, and death in CSF.<sup>120</sup>

There are at least two reasons for determining the potential presence and procoagulant activity of neural cell-derived and tissue factor-bearing microparticles. First, these microparticles could deliver more tissue factor to platelets to enhance coagulation, a direct mechanism for enhancing coagulopathy. Second, they carry a different set of receptors and counter-receptors that may fuse with,<sup>121</sup> or be endocytosed by, other target cells.<sup>122,123</sup> As a result, the coagulation process may spread to cells that could not normally initiate coagulation, such as neurons. Fibrin deposition has indeed been detected in the brain.<sup>124</sup> The process of membrane fusion between microparticles and target cells could also allow otherwise difficult cell-cell interactions (such as between neurons and leukocytes or platelets) to occur.

#### Phosphatidylserine and Brain Phospholipids

Another group of molecules critical for initiating and propagating coagulation is anionic phospholipids. PS serves as a catalyst that links platelet activation to thrombin generation by providing a highly negatively charged surface and by binding to discrete and specific sites on prothrombin, factor Xa, and factor Va.<sup>125</sup> Platelet-derived microparticles are PS-enriched and have a 50- to 100-fold higher procoagulant activity than platelets.<sup>126</sup> PS is also abundantly detected on the surface of microparticles from ATP-stimulated microglial cells and injured neurons.<sup>115</sup>

Anionic phospholipids such as PS and phosphatidylethanolamine (PE) are normally present on the inner leaflet, whereas neutral phospholipids such as phosphatidylcholine (PC) and sphingomyelin are on the outer leaflet of a membrane bilayer. This asymmetrical distribution of membrane phospholipids is maintained by enzymatic transporters,<sup>127,128</sup> but is lost during apoptosis and/or cell activation, leading to the exposure of anionic PS on the outer membrane.<sup>129</sup>

The central nervous system has the highest lipid content next only to adipose tissue,<sup>130</sup> constituting more than half of the dry weight in a human brain.<sup>131</sup> The brain contains all three major categories of lipids: cholesterol, sphingolipids (sphingomyelin, cerebrosides, sulfatides, gangliosides), and phospholipids (phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositols).<sup>132,133</sup> Phospholipids account for approximately 25% of the dry weight of an adult rat brain compared to about 10% in other tissues. Injury to cerebral tissue could therefore release a large quantity of phospholipids and PS-bearing neural microparticles. Both membrane-anchored and soluble PS promotes thrombin generation.<sup>125,136,138</sup> In addition to PS, the anionic PE has also been shown to bind coagulation factor Va with a high affinity ( $K_d$  10 nM) and promote thrombin generation.<sup>139</sup>

Despite strong circumstantial clinical and laboratory evidence, the role of brain-derived phospholipids in the development of TBI-

associated coagulopathy remains unknown. Phospholipids released from injured neural cells, are also highly susceptible to oxidation.<sup>140,141</sup> Whether oxidized PS is more or less active in initiating and propagating coagulation remains a virgin field, not only to TBI-coagulopathy but also to understanding the coagulation state under conditions of oxidative stress. This is the case despite the finding that the lipid peroxidation markers F2-isoprostane and malondialdehyde are increased in brain tissue, serum, and CSF after experimental and clinical TBI.<sup>142,144</sup> More importantly, this increase in lipid peroxidation is associated with worse clinical outcomes in patients with severe TBI.<sup>145,146</sup> Also unique is the fact that phospholipids in different regions of the brain differ in their fatty acid compositions and side chains,<sup>132,133</sup> but association among types of phospholipids, regions of brain injury, and rates of coagulopathy has not been examined. TBI may be the most suitable model system to study how different types of phospholipids affect the initiation and propagation of coagulation. Information thus learned could help developing treatment strategies to not only improve TBI-associated coagulopathy, but also help to understand coagulation deficiencies and thrombosis in other clinical settings.

#### Conclusion

In summary, TBI creates a coagulopathic state in part because cerebral tissue is a rich source of potent platelet activating and procoagulant molecules, but their contribution to the development of TBI-associated coagulopathy remains largely unknown. This lack of understanding has greatly slowed the pace of developing effective therapeutics targeting this potentially lethal pathological state. As a result, clinical trials on the efficacy of anti-thrombin III,<sup>147</sup> low-molecular weight heparin,<sup>148,150</sup> factor VIIa,<sup>151,152</sup> and prostacyclin<sup>153,154</sup> have yielded variable outcomes in the setting of TBI and associated pathologies. The transfusion of blood products, such as platelets, fresh frozen plasma, and cryoprecipitate, has been a standard treatment. However, they are not only time-consuming, but their efficacy remains to be investigated in large trials. Transfusion may also cause fluid volume overload to the patients with isolated TBI and transfusion-related complications.<sup>155,156</sup> Extensive clinical and basic science research is therefore needed to study the pathogenesis of TBI-associated coagulopathy, to understand its impact on clinical outcomes, and to develop effective therapies.

#### Acknowledgment

This study was supported by grants from the National Institute of Health (nos. R01HL71895 and R01HL085769 to J.F.D. and K23GM086729 to T.W.) and a grant from the Chinese Natural Science Foundation (to J.Z., R.J., L.L.).

#### Author Disclosure Statement

No competing financial interests exist.

#### References

1. Niles, S.E., McLaughlin, D.F., Perkins, J.G., Wade, C.E., Li, Y., Spinella, P.C., and Holcomb, J.B. (2008). Increased mortality associated with the early coagulopathy of trauma in combat casualties. *J. Trauma* 64, 1459-1463.
2. Frith, D., and Brohi, K. (2010). The acute coagulopathy of trauma shock: clinical relevance. *Surgeon* 8, 159-163.
3. Mitra, B., Cameron, P.A., Mori, A., and Fitzgerald, M. (2010). Acute coagulopathy and early deaths post major trauma. *Injury* 43, 22-26.



4. Wafaisade, A., Wutzler, S., Lefering, R., Tjardes, T., Banerjee, M., Paffrath, T., Bouillon, B., and Maegele, M. (2010). Drivers of acute coagulopathy after severe trauma: a multivariate analysis of 1987 patients. *Emerg. Med. J.* 27, 934-939.
5. Maani, C.V., DeSocio, P.A., and Holcomb, J.B. (2009). Coagulopathy in trauma patients: what are the main influence factors? *Curr. Opin. Anaesthesiol.* 22, 255-260.
6. Mitra, B., Cameron, P.A., Mori, A., Maini, A., Fitzgerald, M., Paul, E., and Street, A. (2011). Early prediction of acute traumatic coagulopathy. *Resuscitation* 82, 1208-1213.
7. Mitra, B., Tullio, F., Cameron, P.A., and Fitzgerald, M. (2012). Trauma patients with the fear of death. *Emerg. Med. J.* 29, 622-625.
8. Thorsen, K., Ringdal, K.G., Strand, K., Soreide, E., Hagemo, J., and Soreide, K. (2011). Clinical and cellular effects of hypothermia, acidosis and coagulopathy in major injury. *Br. J. Surg.* 98, 894-907.
9. Stein, S.C., and Smith, D.H. (2004). Coagulopathy in traumatic brain injury. *Neurocrit. Care* 1, 479-488.
10. Hulka, F., Mullins, R.J., and Frank, E.H. (1996). Blunt brain injury activates the coagulation process. *Arch. Surg.* 131, 923-927.
11. Harhangi, B.S., Kompanje, E.J., Leebeek, F.W., and Maas, A.I. (2008). Coagulation disorders after traumatic brain injury. *Acta Neurochir. (Wien)* 150, 165-175.
12. Salehpour, F., Bazzazi, A.M., Porhomayon, J., and Nader, N.D. (2011). Correlation between coagulopathy and outcome in severe head trauma in neurointensive care and trauma units. *J. Crit. Care* 26, 352-356.
13. Sun, Y., Wang, J., Wu, X., Xi, C., Gai, Y., Liu, H., Yuan, Q., Wang, E., Gao, L., Hu, J., and Zhou, L. (2011). Validating the incidence of coagulopathy and disseminated intravascular coagulation in patients with traumatic brain injury: analysis of 242 cases. *Br. J. Neurosurg.* 25, 363-368.
14. Talving, P., Benoit, R., Hadjizacharia, P., Inaba, K., Chan, L.S., and Demetriades, D. (2009). Coagulopathy in severe traumatic brain injury: a prospective study. *J. Trauma* 66, 55-61.
15. Franschman, G., Boer, C., Andriessen, T., Van der Naalt, J., Horn, J., Haitsma, I., Jacobs, B., and Vos, P.E. (2012). Multicenter evaluation of the course of coagulopathy in patients with isolated traumatic brain injury: relation to CT characteristics and outcome. *J. Neurotrauma* 29, 128-136.
16. Cap, A.P., and Spinella, P.C. (2011). Severity of head injury is associated with increased risk of coagulopathy in combat casualties. *J. Trauma* 71, S78-S81.
17. Kim, Y.J. (2011). A systematic review of factors contributing to outcomes in patients with traumatic brain injury. *J. Clin. Nurs.* 20, 1518-1532.
18. Stein, S.C., and Spettell, C.M. (1995). Delayed and progressive brain injury in children and adolescents with head trauma. *Pediatr. Neurosurg.* 23, 299-304.
19. Stein, S.C., Young, G.S., Talucci, R.C., Greenbaum, B.H., and Ross, S.E. (1992). Delayed brain injury after head trauma: significance of coagulopathy. *Neurosurgery* 30, 160-165.
20. Kaufman, H.H., Moake, J.L., Olson, J.D., Miner, M.E., duCret, R.P., Pruessner, J.L., and Gildenberg, P.L. (1980). Delayed and recurrent intracranial hematomas related to disseminated intravascular clotting and fibrinolysis in head injury. *Neurosurgery* 7, 445-449.
21. Stein, S.C., Spettell, C., Young, G., and Ross, S.E. (1993). Delayed and progressive brain injury in closed-head trauma: radiological demonstration. *Neurosurgery* 32, 25-30.
22. Graham, D.I., and Adams, J.H. (1971). Ischaemic brain damage in fatal head injuries. *Lancet* 1, 265-266.
23. Maeda, T., Katayama, Y., Kawamata, T., Aoyama, N., and Mori, T. (1997). Hemodynamic depression and microthrombosis in the peripheral areas of cortical contusion in the rat: role of platelet activating factor. *Acta Neurochir. Suppl.* 70, 102-105.
24. Wafaisade, A., Lefering, R., Tjardes, T., Wutzler, S., Simanski, C., Paffrath, T., Fischer, P., Bouillon, B., and Maegele, M. (2010). Acute coagulopathy in isolated blunt traumatic brain injury. *Neurocrit. Care* 12, 211-219.
25. Awasthi, D., Rock, W.A., Carey, M.E., and Farrell, J.B. (1991). Coagulation changes after an experimental missile wound to the brain in the cat. *Surg. Neurol.* 36, 441-446.
26. Auer, L.M., and Ott, E. (1979). Disturbances of the coagulatory system in patients with severe cerebral trauma II. Platelet function. *Acta Neurochir. (Wien)* 49, 219-226.
27. Jacoby, R.C., Owings, J.T., Holmes, J., Battistella, F.D., Gosselin, R.C., and Paglieroni, T.G. (2001). Platelet activation and function after trauma. *J. Trauma* 51, 639-647.
28. Yee, D.L., Sun, C.W., Bergeron, A.L., Dong, J.F., and Bray, P.F. (2005). Aggregometry detects platelet hyperreactivity in healthy individuals. *Blood* 106, 2723-2729.
29. Yee, D.L., Bergeron, A.L., Sun, C.W., Dong, J.F., and Bray, P.F. (2006). Platelet hyperreactivity generalizes to multiple forms of stimulation. *J. Thromb. Haemost.* 4, 2043-2050.
30. Kaufman, H.H., Hui, K.S., Mattson, J.C., Borit, A., Childs, T.L., Hoots, W.K., Bernstein, D.P., Makela, M.E., Wagner, K.A., Kahan, B.D., et al. (1984). Clinicopathological correlations of disseminated intravascular coagulation in patients with head injury. *Neurosurgery* 15, 34-42.
31. Stein, S.C., Graham, D.I., Chen, X.H., and Smith, D.H. (2004). Association between intravascular microthrombosis and cerebral ischemia in traumatic brain injury. *Neurosurgery* 54, 687-691.
32. van der Sande, J.J., Emeis, J.J., and Lindeman, J. (1981). Intravascular coagulation: a common phenomenon in minor experimental head injury. *J. Neurosurg.* 54, 21-25.
33. Stein, S.C., Chen, X.H., Sinson, G.P., and Smith, D.H. (2002). Intravascular coagulation: a major secondary insult in nonfatal traumatic brain injury. *J. Neurosurg.* 97, 1373-1377.
34. Lu, D., Mahmood, A., Goussev, A., Qu, C., Zhang, Z.G., and Chopp, M. (2004). Delayed thrombosis after traumatic brain injury in rats. *J. Neurotrauma* 21, 1756-1766.
35. Henson, P.M. (1970). Release of vasoactive amines from rabbit platelets induced by sensitized mononuclear leukocytes and antigen. *J. Exp. Med.* 131, 287-306.
36. Benveniste, J., Henson, P.M., and Cochrane, C.G. (1972). Leukocyte-dependent histamine release from rabbit platelets. The role of IgE, basophils, and a platelet-activating factor. *J. Exp. Med.* 136, 1356-1377.
37. Chao, W., and Olson, M.S. (1993). Platelet-activating factor: receptors and signal transduction. *Biochem. J.* 292, 617-629.
38. Nakamura, M., Honda, Z., Izumi, T., Sakanaka, C., Mutoh, H., Minami, M., Bito, H., Seyama, Y., Matsumoto, T., Noma, M., et al. (1991). Molecular cloning and expression of platelet-activating factor receptor from human leukocytes. *J. Biol. Chem.* 266, 20400-20405.
39. Kunz, D., Gerard, N.P., and Gerard, C. (1992). The human leukocyte platelet-activating factor receptor. cDNA cloning, cell surface expression, and construction of a novel epitope-bearing analog. *J. Biol. Chem.* 267, 9101-9106.
40. Herbert, J.M., Castro-Faria-Neto, H.C., Barbosa-Filho, J.M., Cordeiro, R.S., and Tibirica, E. (1997). Pharmacological evidence for the putative existence of two different subtypes of PAF receptors on platelets and leukocytes; studies with yangambin. *J. Lipid Mediat. Cell Signal* 17, 1-14.
41. Korth, R.M. (1996). Specific high affinity binding of platelet activating factor to intact human blood neutrophils and eosinophils. *Int. Arch. Allergy Immunol.* 110, 124-131.
42. McIntyre, T.M., Zimmerman, G.A., Satoh, K., and Prescott, S.M. (1985). Cultured endothelial cells synthesize both platelet-activating factor and prostacyclin in response to histamine, bradykinin, and adenosine triphosphate. *J. Clin. Invest.* 76, 271-280.
43. Zimmerman, G.A., McIntyre, T.M., and Prescott, S.M. (1985). Production of platelet-activating factor by human vascular endothelial cells: evidence for a requirement for specific agonists and modulation by prostacyclin. *Circulation* 72, 718-727.
44. Kumar, R., Harvey, S.A., Kester, M., Hanahan, D.J., and Olson, M.S. (1988). Production and effects of platelet-activating factor in the rat brain. *Biochim. Biophys. Acta* 963, 375-383.
45. Goracci, G., and Francescangeli, E. (1991). Properties of PAF-synthesizing phosphocholinesterase and evidence for lysoPAF acetyltransferase activity in rat brain. *Lipids* 26, 986-991.
46. Yue, T.L., and Feuerstein, G.Z. (1994). Platelet-activating factor: a putative neuromodulator and mediator in the pathophysiology of brain injury. *Crit. Rev. Neurobiol.* 8, 11-24.
47. Clark, G.D., Zorumski, C.F., McNeil, R.S., Happel, L.T., Ovella, T., McGuire, S., Bix, G.J., and Swann, J.W. (2000). Neuronal platelet-activating factor receptor signal transduction involves a pertussis toxin-sensitive G-protein. *Neurochem. Res.* 25, 603-611.
48. Yue, T.L., Stadel, J.M., Sarau, H.M., Friedman, E., Gu, J.L., Powers, D.A., Gleason, M.M., Feuerstein, G., and Wang, H.Y. (1992). Pla-

- telet-activating factor stimulates phosphoinositide turnover in neurohybrid NCB-20 cells: involvement of pertussis toxin-sensitive guanine nucleotide-binding proteins and inhibition by protein kinase C. *Mol. Pharmacol.* 41, 281-289.
49. Wang, H.Y., Yue, T.L., Feuerstein, G., and Friedman, E. (1994). Platelet-activating factor: diminished acetylcholine release from rat brain slices is mediated by a Gi protein. *J. Neurochem.* 63, 1720-1725.
  50. Satoh, K., Imaizumi, T., Yoshida, H., Hiramoto, M., and Takamatsu, S. (1992). Increased levels of blood platelet-activating factor (PAF) and PAF-like lipids in patients with ischemic stroke. *Acta Neurol. Scand.* 85, 122-127.
  51. Nishida, K., and Markey, S.P. (1996). Platelet-activating factor in brain regions after transient ischemia in gerbils. *Stroke* 27, 514-518.
  52. Pettigrew, L.C., Meyer, J.J., Craddock, S.D., Butler, S.M., Tai, H.H., and Yokel, R.A. (1995). Delayed elevation of platelet activating factor in ischemic hippocampus. *Brain Res.* 691, 243-247.
  53. Rink, C., and Khanna, S. (2011). Significance of brain tissue oxygenation and the arachidonic acid cascade in stroke. *Antioxid. Redox. Signal* 14, 1889-1903.
  54. Lindsberg, P.J., Yue, T.L., Frerichs, K.U., Hallenbeck, J.M., and Feuerstein, G. (1990). Evidence for platelet-activating factor as a novel mediator in experimental stroke in rabbits. *Stroke* 21, 1452-1457.
  55. Yue, T.L., Gleason, M.M., Hallenbeck, J., and Feuerstein, G. (1991). Characterization of platelet-activating factor-induced elevation of cytosolic free-calcium level in neurohybrid NCB-20 cells. *Neuroscience* 41, 177-185.
  56. Lindsberg, P.J., Yue, T.L., Frerichs, K.U., Hallenbeck, J.M., and Feuerstein, G. (1990). Evidence for platelet-activating factor as a novel mediator in experimental stroke in rabbits. *Stroke* 21, 1452-1457.
  57. Musto, A.E., and Samii, M. (2011). Platelet-activating factor receptor antagonism targets neuroinflammation in experimental epilepsy. *Epilepsia* 52, 551-561.
  58. Deng, Y., Fang, W., Li, Y., Cen, J., Fang, F., Lv, P., Gong, S., and Mao, L. (2009). Blood-brain barrier breakdown by PAF and protection by XQ-1H due to antagonism of PAF effects. *Eur. J. Pharmacol.* 616, 43-47.
  59. Faden, A.I., and Tzengzalian, P.A. (1992). Platelet-activating factor antagonists limit glycine changes and behavioral deficits after brain trauma. *Am. J. Physiol.* 263, R909-R914.
  60. Tokutomi, T., Sigemori, M., Kikuchi, T., and Hirohata, M. (1994). Effect of platelet-activating factor antagonist on brain injury in rats. *Acta Neurochir. Suppl. (Wien)* 60, 508-510.
  61. Wong, D.K., Lurie, F., and Wong, L.L. (2008). The effects of clopidogrel on elderly traumatic brain injured patients. *J. Trauma* 65, 1303-1308.
  62. Ivascu, F.A., Howells, G.A., Junn, F.S., Bair, H.A., Bendick, P.J., and Janczyk, R.J. (2008). Predictors of mortality in trauma patients with intracranial hemorrhage on preinjury aspirin or clopidogrel. *J. Trauma* 65, 785-788.
  63. De Oliveira, C.O., Reimer, A.G., Da Rocha, A.B., Grivicich, I., Schneider, R.F., Roisenberg, I., Regner, A., and Simon, D. (2007). Plasma von Willebrand factor levels correlate with clinical outcome of severe traumatic brain injury. *J. Neurotrauma* 24, 1331-1338.
  64. Yokota, H., Naoe, Y., Nakabayashi, M., Unemoto, K., Kushimoto, S., Kurokawa, A., Node, Y., and Yamamoto, Y. (2002). Cerebral endothelial injury in severe head injury: the significance of measurements of serum thrombomodulin and the von Willebrand factor. *J. Neurotrauma* 19, 1007-1015.
  65. Sporn, L.A., Marder, V.J., and Wagner, D.D. (1986). Inducible secretion of large, biologically potent von Willebrand factor multimers. *Cell* 46, 185-190.
  66. Brouland, J.P., Egan, T., Roussi, J., Bonneau, M., Pignaud, G., Bal, C., Vaiman, M., Andre, P., Herve, P., Mazmanian, G.M., and Drouet, L. (1999). In vivo regulation of von willebrand factor synthesis: von Willebrand factor production in endothelial cells after lung transplantation between normal pigs and von Willebrand factor-deficient pigs. *Arterioscler. Thromb. Vasc. Biol.* 19, 3055-3062.
  67. Moake, J.L., and Chow, T.W. (1998). Increased von Willebrand factor (vWf) binding to platelets associated with impaired vWf breakdown in thrombotic thrombocytopenic purpura. *J. Clin. Apheresis* 13, 126-132.
  68. Furlan, M., Robles, R., and Lammle, B. (1996). Partial purification and characterization of a protease from human plasma cleaving von Willebrand factor to fragments produced by in vivo proteolysis. *Blood* 87, 4223-4234.
  69. Tsai, H.M. (1996). Physiologic cleavage of von Willebrand factor by a plasma protease is dependent on its conformation and requires calcium ion. *Blood* 87, 4235-4244.
  70. Levy, G.G., Nichols, W.C., Lian, E.C., Foroud, T., McClintick, J.N., McGee, B.M., Yang, A.Y., Siemieniak, D.R., Stark, K.R., Gruppo, R., Sarode, R., Shurin, S.B., Chandrasekaran, V., Stabler, S.P., Sabio, H., Bouhassira, E.E., Upshaw, J.D., Jr., Ginsburg, D., and Tsai, H.M. (2001). Mutations in a member of the ADAMTS gene family cause thrombotic thrombocytopenic purpura. *Nature* 413, 488-494.
  71. Dong, J.F., Moake, J.L., Nolasco, L., Bernardo, A., Arceneaux, W., Shrimpton, C.N., Schade, A.J., McIntire, L.V., Fujikawa, K., and Lopez, J.A. (2002). ADAMTS-13 rapidly cleaves newly secreted ultralarge von Willebrand factor multimers on the endothelial surface under flowing conditions. *Blood* 100, 4033-4039.
  72. Bernardo, A., Ball, C., Nolasco, L., Choi, H., Moake, J.L., and Dong, J.F. (2005). Platelets adhered to endothelial cell-bound ultra-large von willebrand factor strings support leukocyte tethering and rolling under high shear stress. *J. Thromb. Hemost.* 3, 562-570.
  73. Yamamoto, K., de, W., V, Fearn, C., and Loskutoff, D.J. (1998). Tissue distribution and regulation of murine von Willebrand factor gene expression in vivo. *Blood* 92, 2791-2801.
  74. Hallenbeck, J.M., Furlow, T.W., Jr., and Gralnick, H.R. (1981). Influence of factor VIII/von Willebrand factor protein (F VIII/vWF) and F VIII/vWF-poor cryoprecipitate on post-ischemic microvascular reperfusion in the central nervous system. *Stroke* 12, 93-97.
  75. Osterud, B., and Rapaport, S.I. (1977). Activation of factor IX by the reaction product of tissue factor and factor VII: additional pathway for initiating blood coagulation. *Proc. Natl. Acad. Sci. USA* 74, 5260-5264.
  76. Drake, T.A., Morrissey, J.H., and Edgington, T.S. (1989). Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. *Am. J. Pathol.* 134, 1087-1097.
  77. Fleck, R.A., Rao, L.V., Rapaport, S.I., and Varki, N. (1990). Localization of human tissue factor antigen by immunostaining with monospecific polyclonal anti-human tissue factor antibody. *Thromb. Res.* 59, 421-427.
  78. Eddleston, M., de la Torre, J.C., Oldstone, M.B., Loskutoff, D.J., Edgington, T.S., and Mackman, N. (1993). Astrocytes are the primary source of tissue factor in the murine central nervous system. A role for astrocytes in cerebral hemostasis. *J. Clin. Invest.* 92, 349-358.
  79. Spicer, E.K., Horton, R., Bloem, L., Bach, R., Williams, K.R., Guha, A., Kraus, J., Lin, T.C., Nemerson, Y., and Konigsberg, W.H. (1987). Isolation of cDNA clones coding for human tissue factor: primary structure of the protein and cDNA. *Proc. Natl. Acad. Sci. USA* 84, 5148-5152.
  80. Ahamed, J., and Ruf, W. (2004). Protease-activated receptor 2-dependent phosphorylation of the tissue factor cytoplasmic domain. *J. Biol. Chem.* 279, 23038-23044.
  81. Paborsky, L.R., Tate, K.M., Harris, R.J., Yansura, D.G., Band, L., McCray, G., Gorman, C.M., O'Brien, D.P., Chang, J.Y., Swartz, J.R., Fung, V.P., Thomas, J.N., and Vehar, G.A. (1989). Purification of recombinant human tissue factor. *Biochemistry* 28, 8072-8077.
  82. Ruf, W., Rehemtulla, A., Morrissey, J.H., and Edgington, T.S. (1991). Phospholipid-independent and -dependent interactions required for tissue factor receptor and cofactor function. *J. Biol. Chem.* 266, 2158-2166.
  83. Bach, R., Konigsberg, W.H., and Nemerson, Y. (1988). Human tissue factor contains thioester-linked palmitate and stearate on the cytoplasmic half-cystine. *Biochemistry* 27, 4227-4231.
  84. Davizon, P., Munday, A.D., and Lopez, J.A. (2010). Tissue factor, lipid rafts, and microparticles. *Semin. Thromb. Hemost.* 36, 857-864.
  85. Pendurthi, U.R., and Rao, L.V. (2008). Role of tissue factor disulfides and lipid rafts in signaling. *Thromb. Res.* 122, S14-S18.
  86. Awasthi, V., Mandal, S.K., Papanna, V., Rao, L.V., and Pendurthi, U.R. (2007). Modulation of tissue factor-factor VIIa signaling by lipid rafts and caveolae. *Arterioscler. Thromb. Vasc. Biol.* 27, 1447-1455.
  87. Semeraro, N., Biondi, A., Lorenzet, R., Locati, D., Mantovani, A., and Donati, M.B. (1983). Direct induction of tissue factor synthesis by endotoxin in human macrophages from diverse anatomical sites. *Immunology* 50, 529-535.

88. Rothberger, H., Zimmerman, T.S., Spiegelberg, H.L., and Vaughan, J.H. (1977). Leukocyte procoagulant activity: enhancement of production in vitro by IgG and antigen-antibody complexes. *J. Clin. Invest.* 59, 549-557.
89. Celi, A., Pellegrini, G., Lorenzet, R., De Blasi, A., Ready, N., Furie, B.C., and Furie, B. (1994). P-selectin induces the expression of tissue factor on monocytes. *Proc. Natl. Acad. Sci. USA* 91, 8767-8771.
90. Bevilacqua, M.P., Pober, J.S., Majeau, G.R., Fiers, W., Cotran, R.S., and Gimbrone, M.A., Jr. (1986). Recombinant tumor necrosis factor induces procoagulant activity in cultured human vascular endothelium: characterization and comparison with the actions of interleukin 1. *Proc. Natl. Acad. Sci. USA* 83, 4533-4537.
91. Bevilacqua, M.P., Pober, J.S., Majeau, G.R., Cotran, R.S., and Gimbrone, M.A., Jr. (1984). Interleukin 1 (IL-1) induces biosynthesis and cell surface expression of procoagulant activity in human vascular endothelial cells. *J. Exp. Med.* 160, 618-623.
92. Song, D., Ye, X., Xu, H., and Liu, S.F. (2009). Activation of endothelial intrinsic NF- $\kappa$ B pathway impairs protein C anticoagulation mechanism and promotes coagulation in endotoxemic mice. *Blood* 114, 2521-2529.
93. Ray, B., Chetter, I.C., Lee, H.L., Ettelaie, C., and McCollum, P.T. (2007). Plasma tissue factor is a predictor for restenosis after femoropopliteal angioplasty. *Br. J. Surg.* 94, 1092-1095.
94. Butenas, S., Bouchard, B.A., Brummel-Ziedins, K.E., Parhami-Seren, B., and Mann, K.G. (2005). Tissue factor activity in whole blood. *Blood* 105, 2764-2770.
95. Giesen, P.L., Rauch, U., Bohrmann, B., Kling, D., Roque, M., Fallon, J.T., Badimon, J.J., Himber, J., Riederer, M.A., and Nemerson, Y. (1999). Blood-borne tissue factor: another view of thrombosis. *Proc. Natl. Acad. Sci. USA* 96, 2311-2315.
96. Zillmann, A., Luther, T., Muller, I., Kotzsch, M., Spannagl, M., Kauke, T., Oelschlagel, U., Zahler, S., and Engelmann, B. (2001). Platelet-associated tissue factor contributes to the collagen-triggered activation of blood coagulation. *Biochem. Biophys. Res. Commun.* 281, 603-609.
97. Camera, M., Frigerio, M., Toschi, V., Brambilla, M., Rossi, F., Cottell, D.C., Maderna, P., Parolari, A., Bonzi, R., De, V.O., et al. (2003). Platelet activation induces cell-surface immunoreactive tissue factor expression, which is modulated differently by antiplatelet drugs. *Arterioscler. Thromb. Vasc. Biol.* 23, 1690-1696.
98. Heemskerk, J.W., Vuist, W.M., Feijge, M.A., Reutelingsperger, C.P., and Lindhout, T. (1997). Collagen but not kinogen surfaces induce bleb formation, exposure of phosphatidylserine, and procoagulant activity of adherent platelets: evidence for regulation by protein tyrosine kinase-dependent  $\text{Ca}^{2+}$  responses. *Blood* 90, 2615-2625.
99. Siljander, P., Farndale, R.W., Feijge, M.A., Comfurius, P., Kos, S., Bevers, E.M., and Heemskerk, J.W. (2001). Platelet adhesion enhances the glycoprotein VI-dependent procoagulant response: involvement of p38 MAP kinase and calpain. *Arterioscler. Thromb. Vasc. Biol.* 21, 618-627.
100. Shcherbina, A., and Remold-Donnell, E. (1999). Role of caspase in a subset of human platelet activation responses. *Blood* 93, 4222-4231.
101. Dale, G.L., and Friese, P. (2006). Bax activators potentiate coated-platelet formation. *J. Thromb. Haemost.* 4, 2664-2669.
102. Brown, S.B., Clarke, M.C., Magowan, L., Sanderson, H., and Savill, J. (2000). Constitutive death of platelets leading to scavenger receptor-mediated phagocytosis. A caspase-independent cell clearance program. *J. Biol. Chem.* 275, 5987-5996.
103. Fox, J.E., Austin, C.D., Reynolds, C.C., and Steffen, P.K. (1991). Evidence that agonist-induced activation of calpain causes the shedding of procoagulant-containing microvesicles from the membrane of aggregating platelets. *J. Biol. Chem.* 266, 13289-13295.
104. Fox, J.E., Austin, C.D., Boyles, J.K., and Steffen, P.K. (1990). Role of the membrane skeleton in preventing the shedding of procoagulant-rich microvesicles from the platelet plasma membrane. *J. Cell Biol.* 111, 483-493.
105. Basse, F., Gaffet, P., and Bienvenue, A. (1994). Correlation between inhibition of cytoskeleton proteolysis and anti-vesiculation effect of calpeptin during A23187-induced activation of human platelets: are vesicles shed by pod fragmentation? *Biochim. Biophys. Acta* 1190, 217-224.
106. Bode, A.P., Orton, S.M., Frye, M.J., and Udis, B.J. (1991). Vesiculation of platelets during in vitro aging. *Blood* 77, 887-895.
107. Miyazaki, Y., Nomura, S., Miyake, T., Kagawa, H., Kitada, C., Taniguchi, H., Komiyama, Y., Fujimura, Y., Ikeda, Y., and Fukuhara, S. (1996). High shear stress can initiate both platelet aggregation and shedding of procoagulant containing microparticles. *Blood* 88, 3456-3464.
108. Chow, T.W., Hellums, J.D., and Thiagarajan, P. (2000). Thrombin receptor activating peptide (SFLLRN) potentiates shear-induced platelet microvesiculation. *J. Lab. Clin. Med.* 135, 66-69.
109. Owens, A.P., III, and Mackman, N. (2011). Microparticles in hemostasis and thrombosis. *Circ. Res.* 108, 1284-1297.
110. Biro, E., Akkerman, J.W., Hoek, F.J., Gorter, G., Pronk, L.M., Sturk, A., and Nieuwland, R. (2005). The phospholipid composition and cholesterol content of platelet-derived microparticles: a comparison with platelet membrane fractions. *J. Thromb. Haemost.* 3, 2754-2763.
111. Falati, S., Liu, Q., Gross, P., Merrill-Skoloff, G., Chou, J., Vandendries, E., Celi, A., Croce, K., Furie, B.C., and Furie, B. (2003). Accumulation of tissue factor into developing thrombi in vivo is dependent upon microparticle P-selectin glycoprotein ligand 1 and platelet P-selectin. *J. Exp. Med.* 197, 1585-1598.
112. Astrup, T. (1965). Assay and content of tissue thromboplastin in different organs. *Thromb. Diath. Haemorrh.* 14, 401-416.
113. Keimowitz, R.M., and Annis, B.L. (1973). Disseminated intravascular coagulation associated with massive brain injury. *J. Neurosurg.* 39, 178-180.
114. Fleck, R.A., Rao, L.V., Rapaport, S.I., and Varki, N. (1990). Localization of human tissue factor antigen by immunostaining with monospecific polyclonal anti-human tissue factor antibody. *Thromb. Res.* 59, 421-437.
115. Bianco, F., Pravettoni, E., Colombo, A., Schenk, U., Moller, T., Matteoli, M., and Verderio, C. (2005). Astrocyte-derived ATP induces vesicle shedding and IL-1 beta release from microglia. *J. Immunol.* 174, 7268-7277.
116. Ferrari, D., Chiozzi, P., Falzoni, S., Dal, S.M., Collo, G., Buell, G., and Di, V.F. (1997). ATP-mediated cytotoxicity in microglial cells. *Neuropharmacology* 36, 1295-1301.
117. Duan, S., and Neary, J.T. (2006). P2X(7) receptors: properties and relevance to CNS function. *Glia* 54, 738-746.
118. Bianco, F., Perrotta, C., Novellino, L., Francolini, M., Riganti, L., Menna, E., Saglietti, L., Schuchman, E.H., Furlan, R., Clementi, E., Matteoli, M., and Verderio, C. (2009). Acid sphingomyelinase activity triggers microparticle release from glial cells. *EMBO J.* 28, 1043-1054.
119. Morel, N., Morel, O., Petit, L., Hugel, B., Cochard, J.F., Freyssinet, J.M., Szark, F., and Dabadie, P. (2008). Generation of procoagulant microparticles in cerebrospinal fluid and peripheral blood after traumatic brain injury. *J. Trauma* 64, 698-704.
120. Huang, M., Hu, Y.Y., and Dong, X.Q. (2009). High concentrations of procoagulant microparticles in the cerebrospinal fluid and peripheral blood of patients with acute basal ganglia hemorrhage are associated with poor outcome. *Surg. Neurol.* 72, 481-489.
121. Del Conde, I., Shrimpton, C.N., Thiagarajan, P., and Lopez, J.A. (2005). Tissue-factor-bearing microvesicles arise from lipid rafts and fuse with activated platelets to initiate coagulation. *Blood* 106, 1604-1611.
122. Terrisse, A.D., Puech, N., Allart, S., Gourdy, P., Xuereb, J.M., Payrastra, B., and Sie, P. (2010). Internalization of microparticles by endothelial cells promotes platelet/endothelial cell interaction under flow. *J. Thromb. Haemost.* 8, 2810-2819.
123. Faille, D., El-Asaad, F., Mitchell, A.J., Alessi, M.C., Chimini, G., Fusai, T., Grau, G.E., and Combes, V. (2011). Endocytosis and intracellular processing of platelet microparticles by brain endothelial cells. *J. Cell Mol. Med.* 10-4934.
124. Bardos, H., Molnar, P., Csecsei, G., and Adany, R. (1996). Fibrin deposition in primary and metastatic human brain tumours. *Blood Coagul. Fibrinolysis* 7, 536-548.
125. Lentz, B.R. (2003). Exposure of platelet membrane phosphatidylserine regulates blood coagulation. *Prog. Lipid Res.* 42, 423-438.
126. Sinauridze, E.I., Kireev, D.A., Popenko, N.Y., Pichugin, A.V., Panteliev, M.A., Krymskaya, O.V., and Ataullakhanov, F.I. (2007). Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thromb. Haemost.* 97, 425-434.
127. Devaux, P.F. (1992). Protein involvement in transmembrane lipid asymmetry. *Annu. Rev. Biophys. Biomol. Struct.* 21, 417-439.



128. Suzuki, J., Umeda, M., Sims, P.J., and Nagata, S. (2010). Calcium-dependent phospholipid scrambling by TMEM16F. *Nature* 468, 834-838.
129. Zwaal, R.F., Comfurius, P., and van Deenen, L.L. (1977). Membrane asymmetry and blood coagulation. *Nature* 268, 358-360.
130. Han, X. (2007). Neurolipidomics: challenges and developments. *Front. Biosci.* 12, 2601-2615.
131. Piomelli, D., Astarita, G., and Rapaka, R. (2007). A neuroscientist's guide to lipidomics. *Nat. Rev. Neurosci.* 8, 743-754.
132. Svennerholm, L., and Vanier, M.T. (1972). The Distribution of lipids in the human nervous system. II Lipid composition of human fetal and infant brain. *Brain Res.* 47, 457-468.
133. Svennerholm, L. (1968). Distribution and fatty acid composition of phosphoglycerides in normal human brain. *J. Lipid Res.* 9, 570-579.
134. Benjamins, J.A., Hajara, A.K., and Agranoff, B.W. (2006). Chemistry and metabolism of brain lipids, in: *Basic Neurochemistry: Molecular, Cellular, and Medical Aspects*. G. Siegel, W.R. Albers, S. Brady, and D. Price (eds). Elsevier Academic Press: San Diego, pps. 33-60.
135. Phillis, J.W., Horrocks, L.A., and Farooqui, A.A. (2006). Cyclooxygenases, lipoxygenases, and epoxygenases in CNS: their role and involvement in neurological disorders. *Brain Res. Rev.* 52, 201-243.
136. Nesheim, M.E., and Mann, K.G. (1983). The kinetics and cofactor dependence of the two cleavages involved in prothrombin activation. *J. Biol. Chem.* 258, 5386-5391.
137. Majumder, R., Weinreb, G., and Lentz, B.R. (2005). Efficient thrombin generation requires molecular phosphatidylserine, not a membrane surface. *Biochemistry* 44, 16998-17006.
138. Majumder, R., Weinreb, G., Zhai, X., and Lentz, B.R. (2002). Soluble phosphatidylserine triggers assembly in solution of a prothrombin-activating complex in the absence of a membrane surface. *J. Biol. Chem.* 277, 29765-29773.
139. Majumder, R., Liang, X., Quinn-Allen, M.A., Kane, W.H., and Lentz, B.R. (2011). Modulation of prothrombinase assembly and activity by phosphatidylethanolamine. *J. Biol. Chem.* 285, 35535-35542.
140. Bayir, H., Tyurin, V.A., Tyurina, Y.Y., Viner, R., Ritov, V., Amoscato, A.A., Zhao, Q., Zhang, X.J., Janesko-Feldman, K.L., Alexander, H., Basova, L.V., Clark, R.S.B., Kochanek, P.M., and Kagan, V.E. (2007). Selective early cardiolipin peroxidation after traumatic brain injury: an oxidative lipidomics analysis. *Ann. Neurol.* 62, 154-69.
141. Tyurin, V.A., Tyurina, Y.Y., Kochanek, P.M., Hamilton, R., DeKosky, S.T., Greenberger, J.S., Bayir, H., and Kagan, V.E. (2008). Oxidative lipidomics of programmed cell death. *Methods Enzymol.* 442, 375-403.
142. Hoffman, S.W., Roof, R.L., and Stein, D.G. (1996). A reliable and sensitive enzyme immunoassay method for measuring 8-iso-prostaglandin F2 alpha: a marker for lipid peroxidation after experimental brain injury. *J. Neurosci. Methods* 68, 133-136.
143. Seifman, M.A., Adamides, A.A., Nguyen, P.N., Vallance, S.A., Cooper, D.J., Kossmann, T., Rosenfeld, J.V., and Morganti-Kossmann, M.C. (2008). Endogenous melatonin increases in cerebrospinal fluid of patients after severe traumatic brain injury and correlates with oxidative stress and metabolic disarray. *J. Cereb. Blood Flow Metab.* 28, 684-696.
144. Sparvero, L.J., Amoscato, A.A., Kochanek, P.M., Pitt, B.R., Kagan, V.E., and Bayir, H. (2010). Mass-spectrometry based oxidative lipidomics and lipid imaging: applications in traumatic brain injury. *J. Neurochem.* 115, 1322-1336.
145. Pilitsis, J.G., Coplin, W.M., O'Brien, M.H., Wellwood, J.M., Diaz, F.G., Fairfax, M.R., Michael, D.B., and Phillis, J.W. (2003). Free fatty acids in cerebrospinal fluids from patients with traumatic brain injury. *Neurosci. Lett.* 349, 136-138.
146. Kasprzak, H.A., Wozniak, A., Drewa, G., and Wozniak, B. (2001). Enhanced lipid peroxidation processes in patients after brain contusion. *J. Neurotrauma* 18, 793-797.
147. Grenander, A., Bredbacka, S., Rydval, A., Aroch, R., Edner, G., Koskinen, L.O., and Olivecrona, M. (2001). Antithrombin treatment in patients with traumatic brain injury: a pilot study. *J. Neurosurg. Anesthesiol.* 13, 49-56.
148. Wahl, F., Grosjean-Piot, O., Bareyre, F., Uzan, A., and Stutzmann, J.M. (2000). Enoxaparin reduces brain edema, cerebral lesions, and improves motor and cognitive impairments induced by a traumatic brain injury in rats. *J. Neurotrauma* 17, 1055-1065.
149. Pahatouridis, D., Alexiou, G.A., Zigouris, A., Mihos, E., Drosos, D., and Voulgaris, S. (2010). Coagulopathy in moderate head injury. The role of early administration of low molecular weight heparin. *Brain Inj.* 24, 1189-1192.
150. Dudley, R.R., Aziz, I., Bonnici, A., Saluja, R.S., Lamoureux, J., Kalmovitch, B., Gursahaney, A., Razek, T., Maleki, M., and Marcoux, J. (2010). Early venous thromboembolic event prophylaxis in traumatic brain injury with low-molecular-weight heparin: risks and benefits. *J. Neurotrauma* 27, 2165-2172.
151. Brown, C.V., Sowery, L., Curry, E., Valadka, A.B., Glover, C.S., Grabarkewitz, K., Green, T., Hail, S., and Admire, J. (2012). Recombinant factor VIIa to correct coagulopathy in patients with traumatic brain injury presenting to outlying facilities before transfer to the regional trauma center. *Am. Surg.* 78, 57-60.
152. Solomon, D., Kim, B., Scultetus, A., Arnaud, F., Auken, C., Freilich, D., and McCarron, R. (2011). The effect of rFVIIa on pro- and anti-inflammatory cytokines in serum and cerebrospinal fluid in a swine model of traumatic brain injury. *Cytokine* 54, 20-28.
153. Grande, P.O., Moller, A.D., Nordstrom, C.H., and Ungerstedt, U. (2000). Low-dose prostacyclin in treatment of severe brain trauma evaluated with microdialysis and jugular bulb oxygen measurements. *Acta Anaesthesiol. Scand.* 44, 886-894.
154. Naredi, S., Olivecrona, M., Lindgren, C., Ostlund, A.L., Grande, P.O., and Koskinen, L.O. (2001). An outcome study of severe traumatic head injury using the fluid therapy with low-dose prostacyclin. *Acta Anaesthesiol. Scand.* 45, 402-406.
155. Popovsky, M.A. (2001). Transfusion and lung injury. *Transfus. Clin. Biol.* 8, 272-277.
156. Winter, J.P., Plummer, D., Bottini, A., Rockswold, G.R., and Ray, D. (1989). Early fresh frozen plasma prophylaxis of abnormal coagulation parameters in the severely head-injured patient is not effective. *Ann. Emerg. Med.* 18, 553-555.

Address correspondence to:  
 Jing-fei Dong, M.D., Ph.D.  
 Puget Sound Blood Center  
 921 Terry Avenue  
 Seattle, WA 98104  
 E-mail: jfdong@psbc.org